

## Letter to the Editor NIFTP: A Critical Pathologist View

### \*Corresponding author

**Sydney Correia Leão, MD**  
Medical Pathology Assistant  
Federal University of São Paulo  
São Paulo, SP, Brazil  
E-mail: [sydneyleao@hotmail.com](mailto:sydneyleao@hotmail.com)

Volume 3 : Issue 1

Article Ref. #: 1000OTLOJ3137

### Article History

Received: February 19<sup>th</sup>, 2017

Accepted: February 23<sup>rd</sup>, 2017

Published: February 24<sup>th</sup>, 2017

### Citation

Leão SC. NIFTP: A critical pathologist view. *Otolaryngol Open J.* 2017; 3(1): 27. doi: [10.17140/OTLOJ-3-137](https://doi.org/10.17140/OTLOJ-3-137)

**Sydney Correia Leão, MD\***

Medical Pathology Assistant, Federal University of São Paulo, São Paulo, SP, Brazil

In the last two years, there has been an interesting discussion on NIFTP (Non-invasive follicular Thyroid neoplasm papillary-like nuclear features), the previous encapsulated follicular variant of papillary thyroid carcinoma (EFVPTC). The study of Nikiforov et al,<sup>1</sup> laid the foundations of this new discovery, which are:

- a. The morphological features, i.e., the follicular growth pattern and nuclear features of papillary thyroid carcinoma (PTC);
- b. Lack of invasion, which separates this tumor from invasive FVPTC;
- c. Clonal origin determined by finding a driver mutation, which indicates that the lesion is biologically a neoplasm; and
- d. A very low risk of adverse outcome when the tumor is non-invasive.

However, some practical pathological problems arose in relation to this new pathological entity:

1. During freezing procedure of thyroidectomy, it would be necessary to freeze the entire capsule to discard microscopic foci of invasion, that would change the classification of the lesion to FVPTC;
2. The NIFTP cannot be diagnosed only by fine needle aspiration cytology (FNAC).<sup>2</sup> In this case, it is necessary to correlate the observations with the imaging findings (Thyroid US with encapsulated lesion);
3. During histological processing, it is interesting to note that the lesion is fully represented, because exclusion criteria for NIFTP such as necrosis and vascular invasion may appear focally;
4. In some developing countries, it is very difficult to perform molecular testing to confirm the mutation that is leading to this lesion.

Due to the aforementioned facts, I believe that NIFTP is a diagnosis of exclusion in patients with encapsulated lesions showing papillary nuclear characteristics and follicular architecture. I also believe that further studies should be done with an attempt to characterize this new entity, definitively separating it from the follicular variant of papillary carcinoma.

### REFERENCES

1. Nikiforov YE, Seethala RR, Tallini G, et al. Nomenclature revision for encapsulated follicular variant of papillary thyroid carcinoma: A paradigm shift to reduce overtreatment of indolent tumors. *JAMA Oncol.* 2016; 2(8): 1023-1029. doi: [10.1001/jamaoncol.2016.0386](https://doi.org/10.1001/jamaoncol.2016.0386)
2. Maletta F, Massa F, Torregrossa L, et al. Cytological features of “noninvasive follicular thyroid neoplasm with papillary-like nuclear features” and their correlation with tumor histology. *Hum Pathol.* 2016; 54: 134-142. doi: [10.1016/j.humpath.2016.03.014](https://doi.org/10.1016/j.humpath.2016.03.014)

### Copyright

©2017 Leão SC. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.