

## Review

### \*Corresponding author

**Léo Guimarães Soares**  
PhD at UERJ  
Rua Ministro Viveiros de Castro  
32/902 -Copacabana, Brazil  
Tel. +55 24 8818.9470  
Fax: 22021-010  
E-mail: [dr\\_leog@hotmail.com](mailto:dr_leog@hotmail.com)

Volume 1 : Issue 1

Article Ref. #: 1000DOJ1102

### Article History:

Received: September 16<sup>th</sup>, 2014

Accepted: October 28<sup>th</sup>, 2014

Published: November 18<sup>th</sup>, 2014

### Citation

Soares LG, Castagna L, Fischer RG, Tinoco EMB. Clinical efficacy of full-Mouth ultrasonic debridement against partial-mouth ultrasonic debridement in the treatment of chronic periodontitis: Full-vs partial-mouth debridement. *Dent Open J.* 2014; 1(1): 5-9. doi: [10.17140/DOJ-1-102](https://doi.org/10.17140/DOJ-1-102)

### Copyright

©2014 Soares LG. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

# Clinical Efficacy of Full-Mouth Ultrasonic Debridement against Partial-Mouth Ultrasonic Debridement in the Treatment of Chronic Periodontitis: Full-Vs Partial-Mouth Debridement

Léo Guimarães Soares<sup>1\*</sup>, Lisiane Castagna<sup>1</sup>, Ricardo Guimarães Fischer<sup>2</sup> and Eduardo Muniz Barreto Tinoco<sup>2</sup>

<sup>1</sup>PhD at UERJ, Rua Boulevard 28 de Setembro, 157 - Vila Isabel - Rio de Janeiro, Brazil

<sup>2</sup>Adjunct Professor, UERJ, UERJ, Rua Boulevard 28 de Setembro, 157 - Vila Isabel - Rio de Janeiro, Brazil

### ABSTRACT

**Aim:** To compare the clinical benefit of full-mouth ultrasonic debridement against partial-mouth ultrasonic debridement in patients with chronic periodontitis up to 3 months after therapy.

**Materials and Methods:** Sixty patients with probing pocket depth  $\geq 5$  mm were randomly assigned to evaluate the clinical efficacy of full-mouth ultrasonic debridement against partial-mouth ultrasonic debridement in the treatment of advanced chronic periodontitis.

**Results:** There was a reduction in the levels of probing pocket depth and relative attachment level but no statistically significant difference between treatment groups in intervals of 30, 60 and 90 days. For bleeding following pocket probing, was found statistical difference for 30 and 60 days, where the groups of full-mouth showed more reduction than conventional therapy. But in 90 days, there was no significant difference between groups.

**Conclusion:** At the evaluation 90 days after treatment, no statistical difference was found between the two periodontal therapies.

**Clinical Relevance:** This study tried to verify if full-mouth ultrasonic debridement provides clinically relevant improvements in the periodontal treatment.

**KEYWORDS:** Chronic periodontitis; Periodontal therapy; Periodontal disease; One-stage full-mouth disinfection.

**ABBREVIATIONS:** PPD: Probing Pocket Depth; SRP: Scaling Root and Planning; CHX: Chlorhexidine; OHI: Oral Hygiene Instruction.

### INTRODUCTION

Chronic periodontitis is the most common form of periodontitis with about 80% of prevalence.<sup>1-3</sup> The periodontal disease is an opportunistic infection associated with the formation of bacterial bio films on the tooth surfaces. This etiological factor acts through direct mechanisms: destruction caused by lytic enzymes and cytotoxins produced by bacteria, and indirect: periodontal destruction by the inflammation.<sup>4</sup>

The bio film is considered the primary agent in the aetiology of periodontitis. However, only the bio film is not enough to determine the disease, genetic and host (eg, oral hygiene

stress, diabetes, and smoking) may also be present.<sup>5</sup> periodontal infections have a multi factorial aetiology involving a susceptible host and the presence of periodontal pathogens.<sup>6</sup>

The chronic periodontitis has slowly progression and it can be classified in relation to the length (number of sites involved), as localized or generalized, and the severity (amount of insertion loss), as light, moderate and severe.<sup>5-7</sup>

The treatment aims to accomplish the removal of dental calculus and cement contaminated with toxins or microorganisms.<sup>3</sup> The scraping conventional instrumentation is performed by mechanical sextants or quadrants at intervals of one to two weeks, so that active treatment is complete in four to six weeks. In the periodontal therapy in a single session (full-mouth disinfection), a new approach suggested for the treatment of periodontal infections, Quirynen, et al.<sup>8</sup> proposed a model of disinfection treatment in one session performing Scaling Root and Planning (SRP) within 24 hours, supplemented with the use of Chlorhexidine (CHX) for two weeks. The dental calculus and cement contaminated can be removed by manual or ultrasonic debridement, and the studies show similar effectiveness in both approaches.<sup>9-10</sup> In a clinical evaluation for six months to evaluate manual and ultrasonic debridement the results were similar.<sup>11</sup> Other study demonstrated that a single session of full-mouth plus ultrasonic is a justified initial treatment approach that offers tangible benefits for the chronic periodontitis patient.<sup>12</sup>

So, the present study compared the clinical benefit of full-mouth ultrasonic debridement against partial-mouth ultrasonic debridement in patients with chronic periodontitis up to 3 months after therapy. This study tried to verify if full-mouth ultrasonic debridement provides clinically relevant improvements in the periodontal treatment.

## MATERIALS AND METHODS

### Selection of patients

This study had voluntary participants that according to inclusion criteria were patients with chronic periodontitis<sup>3</sup> with at least 20 teeth and Probing Pocket Depth (PPD)  $\geq 5$  mm in at least 6 sites; without systemic involvement; without periodontal treatment in the last 12 months; without use of antibiotics in the last 6 months; no pregnancy; no orthodontic therapy; non-smokers; no cardiac pace-makers users. All individuals involved were informed of the importance and purpose of the study and signed an informed consent form, previously approved by the Ethics Committee under number 0082.0.228.000-10.

### Study Design

This research was an intervention study adapted<sup>8,11</sup> to evaluate the effects of periodontal therapies, conventional in quadrants and full-mouth. It was performed between May 2010 and September 2011 with sixty patients, 38 women and 22 men,

aged between 33 and 66 (average  $54.55 \pm 2.60$ ). The power calculation was performed and the analysis indicated that with 30 individuals the study would have 80% power to detect a difference of 1 mm in PPD between the two groups (standard deviation of the error  $1.84 \pm$  and detectable difference of 2.43). So, the patients were included randomly (by coin toss) in two different groups:

**Group A:** Periodontal therapy in a single session (full-mouth ultrasonic debridement) plus mouthwash supplemented with the use of CHX for two weeks

**Group B:** Periodontal conventional therapy ultrasonic debridement in quadrants (partial mouth)

All the patients received Oral Hygiene Instruction (OHI) before the treatment. The CHX was used in the concentration of 0.2%, without alcohol, 15 ml, two times per day, during 60 seconds.

### Experimental phase

The periodontal examination in initial, 30, 60 and 90 days after treatment, was also performed following parameters of: (1) Probing Pocket Depth (PPD), characterized by the distance from the gingival margin to pocket, (2) Relative Attachment Level (RAL), characterized by distance from the cement enamel junction to the pocket, (3) Bleeding following pocket probing (Bop), presence or absence of bleeding 30 seconds after probing.<sup>12</sup> These measurements were performed at four points (mesial, buccal, distal and palatal / lingual) and it was performed by a single calibrated examiner (using a periodontal probe, University of North Carolina type, 15 mm - Hu-Friedy, USA). The single calibrated examiner was calibrated for intra examiner repeatability before the beginning of the study. Five patients with chronic periodontitis were examined at an interval of 48 hours and the reproducibility was 0.990, 0.897 and 0.760 for the mean PPD, RAL and Bop.

The patients were also evaluated by Visible Plaque Index (VPI) with dichotomous counting [presence (1) or absence (0)] and Gingival Bleeding Index (GBI), which was examined all surfaces of all teeth and the presence or absence of gingival bleeding was determined by soft inspection of gingival sulcus with a periodontal probe and the presence of bleeding indicated a positive score, expressed as a percentage of a total number of gingival margins examined.<sup>13</sup>

Then, the periodontal therapy was performed in all the teeth, with ultrasonic debridement (tip P10; Cavitron, Dentsply) and manual curettes type McCall, No. 13/14, 17/18 and type Gracey 5/6, 7/8, 11/12, 13/14 and Hirschfield #5-11 (Hu-Friedy, USA) for SRP supra and sub gingival under aesthetic blocking of the region.

In the conventional therapy, a total of four visits per patient with an interval between the visits about seven days were

The first re evaluation was performed 30 days after the end of treatment, where the teeth were evaluated in PPD, RAL, Bop, VPI and GBI<sup>12,13</sup>. The same measurements were made with 60 and 90 days after the end of treatment.

All the evaluations and treatments were performed by the same single examiner. After the initial periodontal treatment, the patients did not receive any supra or sub gingival treatment, only instruction of oral hygiene when necessary. No patients had to be included in a surgical phase after 90 days because non surgical treatment was successfully performed.

**Statistical analysis**

The data analyses were performed using Shapiro-Wilk, ANOVA (to PPD and RAL) and Kruskal Wallis (to Bop, VPI and GBI). All differences were considered significant at  $p < 0.05$ . Statistical analyses were performed using Sigma Plot<sup>®</sup> statistical software package (System Software Inc., San José, CA, USA).

**RESULTS**

Tables 1 show the means and standard deviations of PPD, RAL and BoP, respectively, in 30, 60 and 90 days of the two therapies, showing reduction of the clinical parameters in all periods, when compared with baseline measurements.

	Groups	Initial	30 days	60 days	90 days
PPD	A	6,37±1,12	4,81±1,43	4,96±1,42	4,77±1,43
	B	6,27±1,33	4,85±1,60	5,01±1,42	4,79±1,64
RAL	A	7,15±1,42	6,17±1,75	6,22±1,73	6,11±1,83
	B	7,08±1,61	6,27±2,11	6,25±1,82	6,10±2,22
BoP	A	85,49±21,44	29,44±22,75	31,40±28,21	14,80±22,30
	B	89,23±18,24	40,25±33,45	41,35±27,80	17,28±29,43

Table 1: Mean (standard deviation) of PPD (mm),RAL (mm) and Bop (%) at different time intervals for both groups

Table 2 shows the result of periodontal therapy in intervals of 30, 60 and 90 days. There was a reduction in the levels of PPD and RAL but without statistical difference between the therapies. For the parameter of Bop, was found a statistical

	30 days		60 days		90 days	
	A	B	A	B	A	B
PPD (mm)	1.56±1.46	1.42±1.47	1.41±1.41	1.26±1.38	1.60±1.45	1.48±1.74
p value	<b>0.879</b>		<b>0.875</b>		<b>0.830</b>	
RAL (mm)	0.98±1.30	0.81±1.26	0.93±1.65	0.83±1.77	1.04±1.28	0.98±1.88
p value	<b>0.790</b>		<b>0.806</b>		<b>0.841</b>	
BoP (%)	56.05	48.98	54.09	47.88	70.69	71.95
p value	<b>0.003</b>		<b>0.003</b>		<b>0.937</b>	

Table 2: Mean (standard deviation) of reduction to the PPD and RAL parameters of the periodontal therapies in 30, 60 and 90 days and median of reduction to Bop parameter (p value: PPD e RAL according ANOVA e Bop according Kruskal Wallis).

difference for 30 and 60 days, where the group of full-mouth showed more reduction of Bop than conventional therapy ( $p=0.003$ ). But in 90 days, there was no significant difference between groups ( $p=0.937$ ).

For VPI and GBI parameters (Table 3), there was not statistical difference between the groups ( $p > 0.05$ ) when comparing the reduction. VPI showed a statically significance ( $p=.007$ ) reduction 90 days after treatment when compared with the initial values.

	Groups	Initial	30 days	60 days	90 days
VPI	A	22.80 ±9.37	12.60 ±8,89	10.55 ±7.80	7.89 ±5.30
	B	20.30 ±7.44	14.45 ±7,67	11.21 ±7.67	8.64 ±3.99
GBI	A	39.44 ±19.45	30.88 ±12.67	23.86 ±16.67	17.37 ±10.22
	B	34.44 ±18.35	29.98 ±14.38	25.38 ±12.03	15.35 ±12.28

Table 3: Mean (standard deviation) of the VPI and GBI (both in %) at different time intervals.

**Discussion**

In the study, the results of different treatments at the end of 90 days of treatment (Table 2) showed that there was no statistical difference between the groups of non-surgical periodontal therapy: full-mouth and conventional therapy. Thus, the clinical results obtained with the non-surgical treatment alone showed efficacy in reducing PPD and RAL, consistent with the results of some studies<sup>9,14-15</sup> which confirmed the success of the therapy in periodontal disease control. Despite others studies<sup>9,14-15</sup> showed the same results, the present study have clinical relevance to the practice of community dentistry and strong interest to the international readership because the most of the researchers in their works asked to have more studies about the comparison of the periodontal therapies. With more studies, different protocols and different samples have about periodontal therapy, more will be possible to understand the disease and find the most effective treatment. So, it is always welcome to the community dentistry a study trying to verify if full-mouth ultrasonic debridement provides clinically relevant improvements in the periodontal treatment.

A systematic review<sup>16</sup> compared clinical effects of periodontal treatment modalities (full-mouth X quadrants) finding no significant differences between them. Several studies have found no significant difference between full-mouth therapy and quadrants,<sup>11,17-18</sup> agreeing with the results observed in this study, where there was a reduction of PPD and RAL levels, however there is still no statistically significant difference between treatment groups.

Another a systematic review<sup>19</sup> calculated an RAL average of 0.64 mm and PPD of 1.18 mm, the data from this study were gains of 0.99 mm and 1.46 mm, in line with the review cited. A study<sup>20</sup> showed six months after full-mouth ultrasonic debridement, the PPD reduction was 2.03 mm and 4.24 mm for the initially moderate and deep pockets, respectively. At the same time, RAL gain was 1.80 mm and 3.90 mm, respectively. And, according another study,<sup>21</sup> in also six months after full-mouth ultrasonic debridement the PPD changes were 1.93 and 3.44mm and RAL gain of 1.21 and 2.41mm for moderate and deep pockets, respectively.

A study<sup>22</sup> used three different treatments, full-mouth with CHX, without CHX and conventional without CHX in 25 patients and, at the end of the study, the authors found no difference between the groups and no difference in the presence or absence of CHX adjunct to treatment. Another study<sup>23</sup> also compared full-mouth against conventional therapy in 39 patients and found that, at the end of 90 days, no differences were found between the groups. Both aforementioned studies are consistent with the data found in the present study.

Some studies<sup>8,11,17,24</sup> compared full-mouth and conventional therapies with CHX groups which included placebo, and there were found some important effect adjunct of CHX within six months of follow up. In the present study, the groups also received full-mouth plus CHX; however the results showed no difference between groups, agreeing with the findings cited above. A systematic review<sup>25</sup> also showed that full-mouth with or without antiseptics not from significant clinical benefits in patients with chronic periodontitis. Anyway, the efficacy of CHX has been demonstrated in several studies<sup>26-28</sup> because of the antibacterial power and substantivity of this solution.

A study<sup>29</sup> evaluated 184 patients with moderate to severe periodontitis in four treatment groups: full-mouth + metronidazole, full-mouth + placebo, conventional + metronidazole and conventional + placebo and no differences were observed in the mean RAL and PPD values between the four experimental groups at baseline and 12 months post-treatment. A study<sup>30</sup> failed to demonstrate differences in the clinical, microbiological or immunological outcome between full-mouth and quadrant scaling and root planning. In the present study, there was no statistically significant difference between treatment groups at different time intervals.

## CONCLUSION

At the evaluation 90 days after treatment, no statistical difference was found between the two periodontal therapies, but at 30 and 60 days, the parameter BoP showed more reduction for the full-mouth therapy.

## REFERENCES

1. Roberts-Thomson K, Do L, Bartold P, Daniels J, Grosse A, Meihubers S. Prevalence, extent and severity of severe periodontal destruction in an urban Aboriginal and Torres Strait Islander population. *Aust Dent J.* 2014; 6. doi: [10.1111/adj.12138](https://doi.org/10.1111/adj.12138)
2. Brown LJ, Loe H. Prevalence, extent, severity and progression of periodontal disease. *Periodontol 2000.* 1993; 2: 57-71. doi: [10.1111/j.1600-0757.1993.tb00220.x](https://doi.org/10.1111/j.1600-0757.1993.tb00220.x)
3. AAP. American Academy of Periodontology. *Ann Periodontol.* 1999.
4. Lindhe J. *Tratado de Periodontia clínica e Implantodontia oral.* 5<sup>th</sup> ed. Rio de Janeiro, Brazil: Guanabara Koogan; 2010.
5. Flemming TF. Periodontitis. *Ann Periodontol.* 1999; 4(1): 32-38.
6. Bollen CML, Beikler T. Halitosis: The multidisciplinary approach. *Int J Oral Sci.* 2012; 4: 55-63. doi: [10.1038/ijos.2012.39](https://doi.org/10.1038/ijos.2012.39)
7. Borrell LN, Burt BA, Taylor GW. Prevalence and trends in periodontitis in the USA. *J Dent Res.* 2005; 84: 924-930. doi: [10.1177/154405910508401010](https://doi.org/10.1177/154405910508401010)
8. Quirynen M, Bollen CM, Vandekerckhove BN, Dekeyser C, Papaioannou W, Eyssen H. Full- vs. partial-mouth disinfection in the treatment of periodontal infections: Short-term clinical and microbiological observations. *J Dent Res.* 1995; 74(8): 1459-1467. doi: [10.1177/00220345950740080501](https://doi.org/10.1177/00220345950740080501)
9. Baderstein A, Nilveus R, Egelberg J. Effect of non-surgical periodontal therapy II. Severely advanced periodontitis. *J Clin Periodontol.* 1984; 11: 63-75. doi: [10.1111/j.1600-051X.1981.tb02024.x](https://doi.org/10.1111/j.1600-051X.1981.tb02024.x)
10. Leon LE, Vogel RI. A comparison of the effectiveness of hand scaling and ultrasonic debridement in furcations as evaluated by differential dark-field microscopy. *J Periodontol.* 1987; 58: 86-94. doi: [10.1902/jop.1987.58.2.86](https://doi.org/10.1902/jop.1987.58.2.86)
11. Wennström JL, Tomasi C, Bertelle A, Dellasega E. Full-mouth ultrasonic debridement versus quadrant scaling and root planning as an initial approach in the treatment of chronic periodontitis. *J Clin Periodontol.* 2005; 32: 851-859. doi: [10.1111/j.1600-051X.2005.00776.x](https://doi.org/10.1111/j.1600-051X.2005.00776.x)

12. Axelson P, Lindhe J. The significance of maintenance care in the treatment of periodontal disease. *J Clin Periodontol*. 1981; 8(4): 281-294. doi: [10.1111/j.1600-051X.1981.tb02039.x](https://doi.org/10.1111/j.1600-051X.1981.tb02039.x)
13. Ainamo J, Bay I. Problems and proposals for recording gingivitis and plaque. *Int Dent J*. 1975; 225-235.
14. Baderstein A, Niveus R, Egelberg J. 4-year observations of basic periodontal therapy. *J Clin Periodontol*. 1987; 14: 438-444. doi: [10.1111/j.1600-051X.1987.tb02248.x](https://doi.org/10.1111/j.1600-051X.1987.tb02248.x)
15. Cobb CM. Non-surgical pocket therapy: Mechanical. *Ann Periodontol*. 1996; 1: 443-490.
16. Eberhard J, Jervøe-Storm PM, Needleman I, Worthington H, Jepsen S. Full mouth treatment concepts for chronic periodontitis: A systematic review. *The Cochrane Library*. 2008. doi: [10.1111/j.1600-051X.2008.01239.x](https://doi.org/10.1111/j.1600-051X.2008.01239.x)
17. Apatzidou DA, Kinane DF. Quadrant root planning versus same-day full mouth root planning. I. Clinical findings. *J Clin Periodontol*. 2004; 31: 132-140. doi: [10.1111/j.0303-6979-2004.00461.x](https://doi.org/10.1111/j.0303-6979-2004.00461.x)
18. Koshy G, Kawashima Y, Kiji M, Nitta H, Umeda M, Nagasawa T, Ishikawa I. Effects of single-visit full-mouth ultrasonic debridement versus quadrant-wise ultrasonic debridement. *J Clin Periodontol*. 2005; 32(7): 734-743. doi: [10.1111/j.1600-051X.2005.00775.x](https://doi.org/10.1111/j.1600-051X.2005.00775.x)
19. Van der Weijden U, Timmerman MF. A systematic review on the clinical efficacy of subgingival debridement in the treatment of chronic periodontitis. *J Clin Periodontol*. 2002; 29: 55-71.
20. Balata ML, Andrade LP, Santos DB, et al. Photodynamic therapy associated with full-mouth ultrasonic debridement in the treatment of severe chronic periodontitis: A randomized-controlled clinical trial. *J Appl Oral Sci*. 2013; 21(2). doi: [10.1590/1678-775201302366](https://doi.org/10.1590/1678-775201302366)
21. Del Peloso Ribeiro E, Bittencourt S, Sallum EA, Nociti FH Jr, Gonçalves RB, Casati MZ. Periodontal debridement as a therapeutic approach for severe chronic periodontitis: A clinical, microbiological and immunological study. *J Clin Periodontol*. 2008; 35: 789-798. doi: [10.1111/j.1600-051X.2008.01292.x](https://doi.org/10.1111/j.1600-051X.2008.01292.x)
22. Swierkot K, Nonnenmacher CI, Mutters R, Flores-de-Jacoby L, Mengel R. One-stage full-mouth disinfection versus quadrant and full-mouth root planning. *J Clin Periodontol*. 2009; 36: 240-249. doi: [10.1111/j.1600-051X.2008.01368.x](https://doi.org/10.1111/j.1600-051X.2008.01368.x)
23. Zijnghe V, Meijer HF, Lie MA, et al. The recolonization hypothesis in a full-mouth or multiple-session treatment protocol: a blinded, randomized clinical trial. *J Clin Periodontol*. 2010; 37: 518-525. doi: [10.1111/j.1600-051X.2010.01562.x](https://doi.org/10.1111/j.1600-051X.2010.01562.x)
24. Quirynen M, Mongardini C, de Soete M, et al. The role of chlorhexidine in the one-stage full-mouth disinfection treatment of patients with advanced adult periodontitis. Long-term clinical and microbiological observations. *J Clin Periodontol*. 2000; 27: 578-589. doi: [10.1034/j.1600-051x.2000.027008578.x](https://doi.org/10.1034/j.1600-051x.2000.027008578.x)
25. Lang NP, Tan WC, Krähenmann MA, Zwahlen M. A systematic review of the effects of full-mouth debridement with and without antiseptics in patients with chronic periodontitis. *J Clin Periodontol*. 2008; 35(Supp 8): 8-21. doi: [10.1111/j.1600-051X.2008.01257.x](https://doi.org/10.1111/j.1600-051X.2008.01257.x)
26. Van Steenberghe D, Avontroodt P, Peeters W, et al. Effect of different mouth rinses on morning breath. *J Periodontol*. 2001; 72: 1183-1191. doi: [10.1902/jop.2000.72.9.1183](https://doi.org/10.1902/jop.2000.72.9.1183)
27. Winkel EG, Roldán S, Van Winkelhoff AJ, Herrera D, Sanz M. Clinical effects of a new mouth rinse containing chlorhexidine, cetylpyridinium chloride and zinc-lactate on oral halitosis. *J Clin Periodontol*. 2003; 30: 300-306. doi: [10.1034/j.1600-051X.2003.00342.x](https://doi.org/10.1034/j.1600-051X.2003.00342.x)
28. Young A, Jonski G, Rölla G. Inhibition of orally produced volatile sulfur compounds by zinc, chlorhexidine or cetylpyridinium chloride-effect of concentration. *Eur J Oral Sci*. 2003; 121: 400-404. doi: [10.1034/j.1600-0722.2003.00063.x](https://doi.org/10.1034/j.1600-0722.2003.00063.x)
29. Preus HR, Gunleiksrud TM, Sandvik L, Gjermo P, Baelum V. A randomized, double blind clinical trial comparing four periodontitis treatment strategies. One-year clinical results. *J Periodontol*. 2012. doi: [10.1902/jop.2012.120400](https://doi.org/10.1902/jop.2012.120400)
30. Kinane DF, Papageorgakopoulos G. Full mouth disinfection versus quadrant debridement: The clinician's choice. *J Int Acad Periodontol*. 2008; 10(1): 6-9.