

Editorial

*Corresponding author

Mohammad Javad Hosseinzadeh-Attar, MD, PhD
Associate Professor
Department of Clinical Nutrition
School of Nutritional Sciences and Dietetics
Tehran University of Medical Sciences
44, Hojjatdoost St, Naderi Ave, Keshavarz Bld
Tehran, Iran
Tel. +00982188993059
E-mail: mhosseinzadeh@tums.ac.ir

Volume 2 : Issue 1

Article Ref. #: 1000OROJ2e002

Article History

Received: April 16th, 2015

Accepted: April 16th, 2015

Published: April 17th, 2015

Citation

Hosseinzadeh-Attar MJ, Alipoor E.
New insight on Adipose tissue function in advanced renal failure. *Obes Res Open J*. 2015; 2(1): e5-e7. doi: [10.17140/OROJ-2-e002](https://doi.org/10.17140/OROJ-2-e002)

Copyright

©2015 Hosseinzadeh-Attar MJ. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

New Insight on Adipose Tissue Function in Advanced Renal Failure

Mohammad Javad Hosseinzadeh-Attar* and Elham Alipoor*

*These authors contributed equally to this work

Department of Clinical Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran

Adipose tissue has been considered simply as an exclusive fat storage organ for a long period. Later in the mid-1990s and after identification of leptin, a revolution occurred in understanding of white adipose tissue function. In fact, adipose tissue has been recognized as a dynamic endocrine organ, which could affect whole body homeostasis.¹ Thereafter, more than 50 secretory peptides, which are called “adipokines” or “adipocytokines”, with diverse metabolic functions have been identified² and the biological role of adipose tissue became more sophisticated and interesting.

The adipokines may act through an autocrine, paracrine and endocrine manner. They are supposed to be involved in a wide range of physiologic to pathologic processes including dietary intake and appetite regulation, energy expenditure, insulin resistance, lipid metabolism, immunity, inflammation and acute-phase responses, vascular homeostasis, endothelial function, angiogenesis and so on.^{2,3} Leptin, adiponectin, ghrelin, resistin, visfatin, vaspin, omentin, TNF- α , and IL-6 are among the most studied adipokines in health and disease. However, studies regarding the extensive biological and metabolic functions of adipokines are still ongoing.

Along with the recent advances in the field of obesity, adipose tissue and adipokines, a great interest has been emerged regarding their relationship with different pathophysiological conditions and their probable mechanism in metabolic disorders. Diabetes, metabolic syndrome, non-alcoholic fatty liver and cardiovascular diseases are among the common diseases, which have been studied. Moreover, pharmacological, nutritional and medical interventions have been done increasingly with the aim of altering the function of adipose tissue, and consequently the gene expression and circulating levels of adipokines in order to decline metabolic dysfunctions and improve the patients' outcomes. Investigating the role of adipose tissue and adipokines in renal insufficiencies especially in End Stage Renal Disease (ESRD) and patients under dialysis has been taken into consideration in recent years.

Adipokines could be destructive for kidney tissue due to the impairment of endothelial function, inducing oxidative stress and inflammation, and stimulating the sympathetic nervous system of kidneys.⁴ It has been proposed that most adipokines act as proinflammatory factors. In contrast, few adipokines act as anti-inflammatory agents and could be protective against the metabolic complications.⁵ Obesity could induce kidney disease through disrupting the balance between protective such as adiponectin and pathologic such as TNF- α adipokines.⁴ The changes of the adipokines' levels such as leptin, adiponectin, visfatin, resistin, IL-6 and TNF- α could result in the reduction of Glomerular Filtration Rate (GFR) and increasing albuminuria, which are among the major pathophysiological mechanisms in CKD. Although the exact mechanism of adipokines' action is still unclear, they may play their role by affecting various types of cells in kidney nephrons.⁶ Previous studies have shown that increased levels of leptin could lead to hypertrophy in glomerular mesangial cells that could activate fibrotic and inflammatory pathways, and also thickening of the basement membrane and consequently glomerulosclerosis.⁶ Moreover, it has been demonstrated that hyperleptinemia could alter the metabolic function of proximal tubular cells and lead to tubular apoptosis.⁷ These structural changes in the nephrons

could result in increased proteinuria, albuminuria and fibrotic pathways in tubular cells through altering the permeability of the cells.⁶ Adiponectin, another major adipokine, is recognized essentially as a protective peptide.⁶ It has been shown that in obesity, hypoadiponectinemia increases Reactive Oxygen Species (ROS) generation and thereby oxidative stress in podocytes, which could probably change the GFR.⁶

Regardless of the roles of adipocytokine in the pathophysiology of renal diseases, the gene expression and circulating levels of these peptides change significantly along with disease progression, particularly in ESRD and dialysis therapy.^{6,8,9} Considering the multiple effects of adipocytokines, their alterations may have considerable biologic, metabolic, and clinical outcomes, which are not well studied in uremic conditions so far.^{9,10} For example, the circulating levels of adiponectin increase in diabetic nephropathy and ESRD, which unexpectedly have been observed to be correlated with higher mortality¹¹⁻¹² and complicate the role of adiponectin in kidney failure.

Moreover, with the identification of new adipokines including Zinc alpha-2 glycoprotein (ZAG), apelin, lipocalin, adipoin, etc. investigating their probable roles in different metabolic aspects of advanced kidney insufficiencies are of major importance. For instance, ZAG is an adipokine that has been proposed to play a role in various metabolic disorders including lipid metabolism,¹³ insulin resistance,¹⁴⁻¹⁵ energy hemostasis,^{13,16-17} inflammation,^{18,19} and determining body composition.^{20,21} In addition, recent studies indicated that the circulating levels of ZAG are increased in hemodialysis.²²⁻²³ However, there are not enough data regarding the effect of ZAG elevation on mentioned disorders in ESRD or patients under regular hemodialysis and the probable effects on patients' prognosis.

In summary, it seems that adipose tissue and its products could be considered as a novel field of study, which may play a major role in the metabolic alteration of advanced renal failure. The limited and mainly descriptive data indicate that further studies are required, more specifically with the aim of identifying the precise roles and mechanisms of adipose tissue and its products in uremic patients, and their probable impact on patients' outcome.

CONFLICTS OF INTEREST

The authors declare that there is no conflict of interest.

ACKNOWLEDGMENTS

We should thank Tehran University of Medical Sciences for their full support.

REFERENCES

1. Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, Friedman JM. Positional cloning of the mouse obese gene and its human homologue. *Nature*. 1994; 372: 425-432. doi: [10.1038/372425a0](https://doi.org/10.1038/372425a0)
2. Trayhurn P, Wood IS. Adipokines: inflammation and the pleiotropic role of white adipose tissue. *Br J Nutr*. 2004; 92: 347-355. doi: [10.1079/BJN20041213](https://doi.org/10.1079/BJN20041213)
3. Lago F, Dieguez C, Gomez-Reino J, Gualillo O. The emerging role of adipokines as mediators of inflammation and immune responses. *Cytokine Growth Factor Rev*. 2007; 18: 313-325. doi: [10.1016/j.cytogfr.2007.04.007](https://doi.org/10.1016/j.cytogfr.2007.04.007)
4. Ruster C, Wolf G. Adipokines promote chronic kidney disease. *Nephrol Dial Transplant*. 2013; 28(Suppl 4): iv8-14. doi: [10.1093/ndt/gft191](https://doi.org/10.1093/ndt/gft191)
5. Ohashi K, Shibata R, Murohara T, Ouchi N. Role of anti-inflammatory adipokines in obesity-related diseases. *Trends in endocrinology and metabolism: TEM*. 2014; 25: 348-355. doi: [10.1016/j.tem.2014.03.009](https://doi.org/10.1016/j.tem.2014.03.009)
6. Briffa JF, McAinch AJ, Poronnik P, Hryciw DH. Adipokines as a link between obesity and chronic kidney disease. *Am J Physiol Renal Physiol*. 2013; 305: F1629-F1636. doi: [10.1152/ajprenal.00263.2013](https://doi.org/10.1152/ajprenal.00263.2013)
7. Briffa JF, Grinfeld E, McAinch AJ, Poronnik P, Hryciw DH. Short term exposure to elevated levels of leptin reduces proximal tubule cell metabolic activity. *Mol Cell Endocrinol*. 2014; 382: 38-45. doi: [10.1016/j.mce.2013.09.001](https://doi.org/10.1016/j.mce.2013.09.001)

8. Iglesias P, Diez JJ. Adipose tissue in renal disease: clinical significance and prognostic implications. *Nephrol Dial Transplant*. 2010; 25: 2066-2077. doi: [10.1093/ndt/gfq246](https://doi.org/10.1093/ndt/gfq246)
9. Axelsson J, Stenvinkel P. Role of fat mass and adipokines in chronic kidney disease. *Curr Opin Nephrol Hypertens*. 2008; 17: 25-31.
10. Jia T, Carrero JJ, Lindholm B, Stenvinkel P. The complex role of adiponectin in chronic kidney disease. *Biochimie*. 2012; 94: 2150-2156. doi: [10.1016/j.biochi.2012.02.024](https://doi.org/10.1016/j.biochi.2012.02.024)
11. Menon V, Li L, Wang X, et al. Adiponectin and mortality in patients with chronic kidney disease. *J Am Soc Nephrol*. 2006; 17: 2599-2606. doi: [10.1681/ASN.2006040331](https://doi.org/10.1681/ASN.2006040331)
12. Jorsal A, Tarnow L, Frystyk J, et al. Serum adiponectin predicts all-cause mortality and end stage renal disease in patients with type I diabetes and diabetic nephropathy. *Kidney Int*. 2008; 74: 649-654. doi: [10.1038/ki.2008.201](https://doi.org/10.1038/ki.2008.201)
13. Russell ST, Tisdale MJ. Antidiabetic properties of zinc-alpha2-glycoprotein in ob/ob mice. *Endocrinology*. 2010; 151: 948-957. doi: [10.1210/en.2009-0827](https://doi.org/10.1210/en.2009-0827)
14. Mracek T, Ding Q, Tzanavari T, et al. The adipokine zinc-alpha2-glycoprotein (ZAG) is downregulated with fat mass expansion in obesity. *Clin Endocrinol (Oxf)*. 2010; 72: 334-341. doi: [10.1111/j.1365-2265.2009.03658.x](https://doi.org/10.1111/j.1365-2265.2009.03658.x)
15. Ceperuelo-Mallafre V, Naf S, Escote X, et al. Circulating and adipose tissue gene expression of zinc-alpha2-glycoprotein in obesity: its relationship with adipokine and lipolytic gene markers in subcutaneous and visceral fat. *J Clin Endocrinol Metab*. 2009; 94: 5062-5069. doi: [10.1210/jc.2009-0764](https://doi.org/10.1210/jc.2009-0764)
16. Eckardt K, Schober A, Platzbecker B, et al. The adipokine zinc-alpha2-glycoprotein activates AMP kinase in human primary skeletal muscle cells. *Arch Physiol Biochem*. 2011; 117: 88-93. doi: [10.3109/13813455.2011.560950](https://doi.org/10.3109/13813455.2011.560950)
17. Bing C, Russell ST, Beckett EE, et al. Expression of uncoupling proteins-1, -2 and -3 mRNA is induced by an adenocarcinoma-derived lipid-mobilizing factor. *Br J Cancer*. 2002; 86: 612-618.
18. Gao D, Trayhurn P, Bing C. Macrophage-secreted factors inhibit ZAG expression and secretion by human adipocytes. *Mol Cell Endocrinol*. 2010; 325: 135-142. doi: [10.1016/j.mce.2010.05.020](https://doi.org/10.1016/j.mce.2010.05.020)
19. Selva DM, Lecube A, Hernandez C, Baena JA, Fort JM, Simo R. Lower zinc-alpha2-glycoprotein production by adipose tissue and liver in obese patients unrelated to insulin resistance. *J Clin Endocrinol Metab*. 2009; 94: 4499-4507. doi: [10.1210/jc.2009-0758](https://doi.org/10.1210/jc.2009-0758)
20. Hirai K, Hussey HJ, Barber MD, Price SA, Tisdale MJ. Biological evaluation of a lipid-mobilizing factor isolated from the urine of cancer patients. *Cancer Res*. 1998; 58: 2359-2365.
21. Todorov PT, McDevitt TM, Meyer DJ, Ueyama H, Ohkubo I, Tisdale MJ. Purification and characterization of a tumor lipid-mobilizing factor. *Cancer Res*. 1998; 58: 2353-2358.
22. Philipp A, Kralisch S, Bachmann A, et al. Serum levels of the adipokine zinc-alpha2-glycoprotein are increased in chronic hemodialysis. *Metabolism*. 2011; 60: 669-672. doi: [10.1016/j.metabol.2010.06.019](https://doi.org/10.1016/j.metabol.2010.06.019)
23. Leal VO, Lobo JC, Stockler-Pinto MB, et al. Zinc-alpha2-glycoprotein: is there association between this new adipokine and body composition in hemodialysis patients? *Ren Fail*. 2012; 34: 1062-1067. doi: [10.3109/0886022X.2012.712859](https://doi.org/10.3109/0886022X.2012.712859)