

Illustration

Corresponding author

Vineet Bhatia, MBBS, MD, DM, DNB, MNAMS, FESC
Senior Consultant Cardiologist
Department of Cardiology
Fortis Hospital
Noida, India
E-mail: vinmonica@yahoo.co.in

Volume 2 : Issue 1

Article Ref. #: 1000HROJ2109

Article History

Received: April 8th, 2015

Accepted: April 14th, 2015

Published: April 16th, 2015

Citation

Bhatia V, Arora P, Minocha A. Saddle thrombus seen on transthoracic echo: a rare feature in pulmonary embolism. *Heart Res Open J.* 2015; 2(1): 51-52. doi: [10.17140/HROJ-2-109](https://doi.org/10.17140/HROJ-2-109)

Saddle Thrombus seen on Transthoracic Echo: A Rare Feature in Pulmonary Embolism

Vineet Bhatia*, Parneesh Arora and Anil Minocha

Department of Cardiology, Fortis Hospital, Noida, India

The Saddle thrombus is a rare condition characterized by presence of a large thrombus which straddles the main pulmonary arterial trunk and its bifurcation leading to acute massive Pulmonary Embolism (PE). Its incidence among patients diagnosed with PE is approximately 2.6%.¹ Saddle PE frequently results in significant hemodynamic instability and signals the potential for imminent hemodynamic collapse. This is rarely seen on Trans-thoracic echocardiography (TTE). We recently came across a patient who presented with breathlessness following transatlantic air travel. His TTE (routine and 3D) revealed a saddle thrombus in the main pulmonary arterial trunk extending into its two main branches (Figures 1 and 2). A CT pulmonary angiogram confirmed the same (Figure 3). He was subsequently successfully thrombolized and made an uneventful recovery.




Figure 1: Echocardiographic Parasternal short axis view (PSAX) shows a saddle thrombus at the Main Pulmonary artery bifurcation extending into its two branches (marked by an arrow).

Copyright

©2015 Bhatia V. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Figure 2: 3 dimensional Echocardiographic Parasternal short axis view (PSAX) shows a saddle thrombus at the Main Pulmonary artery bifurcation extending into its two branches.



Figure 3: A CT pulmonary angiogram showing the saddle thrombus.

CONFLICTS OF INTEREST: None.

REFERENCES

1. Pruszczyk P, Pacho R, Ciurzynski M, et al. Short term clinical outcome of acute saddle pulmonary embolism. *Heart*. 2003; 89: 335-336. doi: [10.1136/heart.89.3.335](https://doi.org/10.1136/heart.89.3.335)