

Research

*Corresponding author

Paul H. Falcone, MS
MusclePharm Sports Science Institute
4721 Ironton St., Building A
Denver, CO 80239, USA
Tel. (303) 396-6105
Fax: (800) 490-7165
E-mail: paul@musclepharm.com

Volume 3 : Issue 1

Article Ref. #: 1000PCSOJ3120

Article History

Received: November 30th, 2016

Accepted: January 11th, 2017

Published: January 11th, 2017

Citation

Falcone PH, Tai C-Y, Carson LR, et al. The effect of mild dehydration induced by heat and exercise on cognitive function. *Psychol Cogn Sci Open J*. 2017; 3(1): 17-23. doi: [10.17140/PCSOJ-3-120](https://doi.org/10.17140/PCSOJ-3-120)

Copyright

©2017 Falcone PH. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The Effect of Mild Dehydration Induced by Heat and Exercise on Cognitive Function

Paul H. Falcone, MS^{1*}; Chih-Yin Tai, EdD¹; Laura R. Carson, BS¹; Jesse A. Gwinn, BA¹; Tyler R. Mccann, BA¹; Nick J. Loveridge, BA¹; Jordan R. Moon, PhD²

¹MusclePharm Sports Science Institute, 4721 Ironton St., Building A, Denver, CO 80239, USA

²American Public University System, School of Health Sciences, Charles Town, WV, USA

ABSTRACT

Background: Past studies have demonstrated cognitive impairment after dehydration, though results are not always consistent. Methodological differences may account for these discrepancies. Therefore, the purpose of the present study was to determine the effects of mild dehydration on various domains of cognitive function.

Methods: Twenty-seven men (n=14; 26.9±4.6 years; 176.3±6.6 cm; 79.1±9.0 kg) and women (n=13; 27.3±4.3 years; 171.0±6.5 cm; 66.5±4.5 kg) participated in the study. Subjects were dehydrated to approximately 2% body mass loss *via* treadmill running and sauna. Measurements were collected prior to and after dehydration. Cognitive function tests included: Finger tapping, symbol digit coding, Stroop, and shifting attention tests.

Results: Reaction time during the shifting attention test was significantly lower from pre-testing to post-testing (PRE: 882.67±126.59 ms; POST: 830.00±105.83 ms; $p=0.0012$). The Stroop reaction time was significantly lower from pre-testing to post-testing (PRE: 712.56±97.52 ms; POST: 671.63±97.21 ms; $p=0.02$). No significant changes were observed in any other measurements.

Conclusion: Dehydration may enhance cognitive functioning in the areas of Stroop reaction time and reaction time during the shifting attention test.

KEYWORDS: Reaction time; Executive function; Heat stress; Thermoregulatory; Amino acid.

INTRODUCTION

Water is an essential nutrient for many bodily systems and functions, such as regulation of body temperature, lubrication of joints, and optimal cardiovascular function.¹ Hydration is especially important in sports and physical activity, due to prodigious perspiration. However, dehydration is a common problem since many athletic individuals do not hydrate properly. Evidence supports the idea that dehydration leads to decrements in physical performance.²⁻⁴ Regarding cognitive function, many researchers contend that dehydration leads to reduced cognitive function,⁵⁻¹⁰ though the evidence is equivocal.¹¹⁻¹⁴

One possible reason for the mixed results in the literature is that the methods for testing cognitive function are inconsistent among studies. Because there are many cognitive domains that contribute to cognitive function, some studies focused on specific domains. In many of the studies on athletic populations such as golfers,¹⁰ soccer players,¹⁵ and jockeys,¹⁶ the testing methods were sport-specific to determine the effects of dehydration on the cognitive aspect of their specific sport. Task-specific outcomes are also seen among cognitive tests in military personnel.^{9,13} While sport-specific testing methods may be ideal for elite athletes, they may not serve recreational or amateur athletes who lack the necessary level of expertise.

Conversely, many studies that tested a wider range of domains chose cognitive tests that may not be ideal for athletic individuals. Though reaction time has been included in many studies to date, researchers often focused on short- and long-term memory^{5,6} or employed tests

that may not translate well to sport, such as perceptive discrimination *via* visual determination of line length.⁷ More recently, studies have tested areas such as executive function¹¹ and sustained attention,^{17,18} but a paucity of data exists in areas such as motor speed and psychomotor speed, which are essential for athletes and athletic individuals. More research needs to be conducted to establish a set of cognitive tests and/or cognitive domains that accurately measure the cognitive functions necessary for athletic performance.

Another reason for the mixed results in the literature is the variety of methods employed to achieve dehydration. Common interventions include: Water deprivation,^{10,19} passive heat stress,^{7,11,16} exercise,^{11,14,20} combined exercise and heat,^{5,6,12} and comparison of dehydration methods.^{7,8} Since dehydration can result from various and multiple insults, various dehydration methods must be tested and should reflect the population being studied. In a sport context, most dehydration occurs during exercise in a hot environment; therefore, both exercise and heat should be employed to elicit dehydration to measure cognitive function in athletes of all levels.

Therefore, the purpose of the present study was to determine the effects of mild dehydration in recreational athletes elicited by a dehydration protocol employing both exercise and heat on various domains of cognitive function. Cognitive domains were chosen to be sport-specific and include: Psychomotor speed, reaction time, cognitive flexibility, processing speed, executive function, and motor speed. Our hypothesis was that mild dehydration would cause a decrement in cognitive function in all tests.

METHODS

Subjects

Twenty-seven men ($n=14$; age: 26.9 ± 4.6 years; height: 176.3 ± 6.6 cm; weight: 79.1 ± 9.0 kg) and women ($n=13$; age: 27.3 ± 4.3 years; height: 171.0 ± 6.5 cm; weight: 66.5 ± 4.5 kg) participated in the study. All individuals were recreationally trained, defined as exercising two or more times per week for at least one year, as assessed by an exercise status questionnaire. Subjects were free of cardiovascular disease, diabetes, or any other diseases that would exclude them from participation in the study, as assessed by a health history questionnaire. No medications or supplements that could interfere with the study had been consumed for at least one month prior to testing. All protocols for this study were approved by an Institutional Review Board and written informed consent was provided by the subjects prior to testing.

Experimental Design

Subjects arrived to the laboratory in the morning in a euhydrated state after an overnight fast. Subjects were also required to abstain from exercise for 24 hours prior to testing so that fatigue was not a factor. A urine sample was provided and measured

for urine specific gravity *via* refractometry (Atago, Inc., Tokyo, Japan) to ensure hydration status. If the USG value was above 1.025, the subject was considered dehydrated and given 500 ml of water to drink in accordance with ACSM guidelines prior to testing to achieve euhydration at baseline. Urine color, body temperature and skin temperature were measured, followed by cognitive function testing. After baseline testing, subjects performed the dehydration protocol. Subsequently, all tests were repeated.

Dehydration

A combination of active dehydration *via* treadmill running and passive dehydration *via* sauna was employed to achieve dehydration. After baseline testing, subjects were required to run on a treadmill (Woodway, Inc., Waukesha, WI, USA) for 30 minutes at 80% maximum heart rate ($206.9 - (0.67 * \text{Age})$).²¹ Afterward, subjects sat in a sauna at 70 °C for 15 minute intervals until 2% body weight loss was attained, which is the value that performance decrements begin to occur.²² Nude, dry body weight was measured between sauna intervals *via* calibrated scale to ensure that the body weight loss was as close to 2% as possible. When 2% body weight loss was achieved, subjects showered and dried their bodies completely. Nude, dry body weight was measured again and this weight was used as the post-dehydration body weight in the present study.

Measurements

All measurements were taken prior to and immediately following the dehydration protocol. After USG was measured, urine color was measured using a color chart developed by Armstrong and colleagues and validated in young athletes.^{23,24} Body temperature *via* digital thermometer (American Diagnostic Corporation, Hauppauge, NY, USA) and skin temperature *via* forehead infrared thermometer (Telatemp Corp., Anaheim, CA, USA) were measured, immediately followed by cognitive function testing. Subjects sat at a computer and performed a variety of cognitive function tests (CNS Vital Signs Inc., Morrisville, NC, USA), including: Finger tapping test, symbol digit coding, Stroop test, and shifting attention test. The CNS Vital Signs test battery was chosen for its high reliability,²⁵ lack of a learning effect,²⁵ and the application of its tests to sports performance. The finger tapping test requires subjects to tap the space bar with their right hand as many times as they can in ten seconds. The test is repeated with the left hand, and two trials are performed with each hand. The symbol digit coding test involves a serial presentation of symbols and empty boxes below, along with a legend at the top that designates a number to correspond with each possible symbol. Subjects must enter the number that corresponds with the highlighted symbol. The stroop test consists of three parts. In the first part, the words RED, YELLOW, BLUE and GREEN appear at random on the screen and the subject responds as quickly as possible by hitting the space bar (simple reaction time). In the second part, the words RED, YELLOW, BLUE and GREEN appear in color on the screen randomly and the subject responds

only when the color matches the word (complex reaction time). In the third part, the words RED, YELLOW, BLUE and GREEN appear in color on the screen randomly and the subject responds only when the color does not match the word (Stroop reaction time). The shifting attention test involves three geometric figures on a screen – one on the top and two on the bottom. The top figure is either a square or a circle. The bottom figures are a square and a circle. The figures are either red or blue (color is randomized). Subjects must match one of the bottom figures with the top figure *via* arrow keys. The rules of matching are changed randomly (either by color or by shape). Each test started with a practice test to ensure familiarization with the test protocol. The testing room was free of distraction and extraneous noise. A total of 11 outcome variables were collected from each subject, which were then used to calculate 6 cognitive domains (Figure 1).

Statistics

All data were analyzed using SPSS software (Version 20.00, SPSS Inc., Chicago, IL, USA). Dependent *t*-tests were used to calculate time differences between pre- and post-dehydration cognitive function. Since multiple pairwise tests were being performed on the same set of data, a Bonferroni correction was used to adjust the *p* value. Therefore, significance was set at $p \leq 0.0029$.

RESULTS

Mild dehydration was achieved in all subjects (weight lost: 1.51 ± 0.27 kg; percent dehydration: $2.11 \pm 0.32\%$). Significant

differences were observed for urine specific gravity values (PRE: 1.008 ± 0.008 ; POST: 1.015 ± 0.009 ; $p = 0.00006$) and for urine color (PRE: 3.22 ± 1.27 ; POST: 4.17 ± 1.30 ; $p = 0.018$). No significant differences were observed for body temperature (PRE: 36.29 ± 0.21 °C; POST: 36.29 ± 0.19 °C; $p > 0.05$) or for skin temperature (PRE: 34.28 ± 0.60 °C; POST: 34.08 ± 1.30 °C; $p > 0.05$). All results for cognitive domains and specific test results can be found in Table 1. Significant differences were not demonstrated in any of the six cognitive domains measured: psychomotor speed (PRE: 203.74 ± 23.50 ; POST: 207.48 ± 27.29 ; $p > 0.05$); reaction time (PRE: 658.48 ± 63.38 ms; POST: 645.67 ± 95.85 ms; $p > 0.05$); cognitive flexibility (PRE: 53.22 ± 8.82 ; POST: 53.93 ± 7.82 ; $p > 0.05$); processing speed (PRE: 83.11 ± 14.20 ; POST: 86.04 ± 17.17 ; $p > 0.05$); executive functioning (PRE: 54.93 ± 8.32 ; POST: 55.74 ± 7.09 ; $p > 0.05$); motor speed (PRE: 119.33 ± 13.50 ; POST: 119.56 ± 16.26 ; $p > 0.05$). Regarding specific tests, two measurements demonstrated a significant difference from pre to post at $p \leq 0.05$, and one measurement demonstrated a significant difference according to the Bonferroni correction ($p \leq 0.0029$). The reaction time during the shifting attention test was significantly lower at post-testing, compared to pretesting (PRE: 882.67 ± 126.59 ms; POST: 830.00 ± 105.83 ms; $p = 0.0012$). The Stroop reaction time during the Stroop test was significantly lower at post-testing, compared to pretesting (PRE: 712.56 ± 97.52 ms; POST: 671.63 ± 97.21 ms; $p = 0.02$), but was not significant when the Bonferroni correction was applied. No significant changes were observed in any other measurements during specific tests.




Table 1: Summary Data for Cognitive Function Domains and Specific Test Results before and after Dehydration.

		Mean	SD	Mean Difference	p Value	Effect Size (d)
Psychomotor speed	Pre	203.74	23.50	3.74	0.16	0.14
	Post	207.48	27.29			
Reaction time†	Pre	658.48	63.38	-12.81	0.33	0.24
	Post	645.67	95.85			
Cognitive flexibility	Pre	53.22	8.82	0.70	0.49	0.36
	Post	53.93	7.82			
Processing speed	Pre	83.11	14.20	2.93	0.13	0.54
	Post	86.04	17.17			
Executive functioning	Pre	54.93	8.32	0.81	0.45	0.47
	Post	55.74	7.09			
Motor speed	Pre	119.33	13.50	0.22	0.89	0.04
	Post	119.56	16.26			
Finger tapping test: right taps average	Pre	62.63	7.60	-0.48	0.63	0.22
	Post	62.15	10.38			
Finger tapping test: left taps average	Pre	56.33	7.18	1.04	0.27	0.72
	Post	57.37	7.11			
Symbol digit coding: correct	Pre	84.41	14.69	2.52	0.19	0.44
	Post	86.93	17.58			
Symbol digit coding: errors†	Pre	1.30	1.81	0.22	0.54	2.59
	Post	1.52	1.31			
Stroop: simple reaction time†	Pre	345.93	41.87	-6.70	0.45	0.25
	Post	339.22	49.02			
Stroop: complex reaction time†	Pre	604.78	69.24	14.52	0.46	0.22
	Post	619.30	123.46			
Stroop: Stroop reaction time†	Pre	712.56	97.52	-40.93	0.02*	0.55
	Post	671.63	97.21			
Stroop: commission errors†	Pre	1.67	1.57	0.15	0.75	0.99
	Post	1.81	2.48			
Shifting attention test: correct	Pre	57.74	6.65	1.48	0.07	1.29
	Post	59.22	5.77			
Shifting attention test: errors†	Pre	2.81	2.56	0.67	0.08	3.75
	Post	3.48	2.05			
Shifting attention test: reaction time†	Pre	882.67	126.59	-52.67	0.00121**	0.53
	Post	830.00	105.83			

*indicates post significantly different from pre at $p \leq 0.05$. **indicates post significantly different from pre at $p \leq 0.0029$. †indicates tests in which a lower score is better.

DISCUSSION

In contrast to our hypothesis, the main findings of the present study indicated that Stroop reaction time during the Stroop test and reaction time during the shifting attention test were enhanced by a sport-specific dehydration protocol employing both heat and exercise, and reaction time during the shifting attention test was significantly different even after the Bonferroni correction was applied. Previous studies have generally found that dehydration caused a decrement in cognitive function⁵⁻¹⁰ or that no significant changes occurred.^{13-15,18,19,26} However, three studies did demonstrate cognitive improvement after dehydration.¹¹⁻¹³ It is possible that the differences among cognitive function as-

essment tools and domains contributed to the discrepancy in findings. For example, Tomporowski and associates¹¹ employed a category-switching task developed by Kramer²⁷ that is very similar to the shifting attention test in the present study. Reaction times were faster after exercise in the Tomporowski study, regardless of whether subjects were 3% dehydrated or if fluid replacement kept them hydrated, which supports the faster reaction times during the shifting attention test in the present study. Also, errors significantly increased after exercise during the category-switching task when the response required the subject to switch categories, i.e. when the task became more complex, as opposed to a non-switching response.¹¹ Similarly, there was a trend observed for errors ($p=0.08$) on the shifting attention test

in the present study. Therefore, consistency in testing methods is essential in order to compare results among studies.

In other studies that tested for reaction time, results were either positive or not significant. Leibowitz et al¹² induced either 2.5% or 5% dehydration using exercise performed in a heat chamber, which improved reaction time in response to peripheral stimuli but no change in response to central stimuli. In a marksmanship test in volunteer soldiers, response time between target presentation and firing of the rifle improved after exercise with no effect from 2.6% dehydration.¹³ Cullen et al¹⁶ observed no effect of 4% dehydration on choice reaction time in jockeys, which was supported in other studies employing water deprivation to achieve 1.5% and unreported dehydration, respectively.^{10,19} Though the literature is not conclusive, it is interesting to note that most of the studies that demonstrated cognitive decrements did not test for reaction time.^{5,6,9,10} In the study by Cian and associates where cognitive decrements were observed after 2.7% dehydration and choice reaction time was measured, cognitive function worsened in perceptual discrimination and short-term and long-term memory, but no changes were observed in choice reaction time.⁷ Therefore, it seems plausible that choice reaction time is a cognitive domain that is insensitive to dehydration or is improved in some cases of exercise-induced dehydration.

Since there are a variety of cognitive domains that contribute to cognitive function, it is important to test domains that are specific to the population and the outcomes of interest. In healthy, active individuals, there are many possible outcomes of interest, such as quality of life, decreased risk for chronic disease, and improvements in work output, school performance, and sports performance. Cognitive domains such as short- and long-term memory may be more applicable to quality of life and school performance, whereas choice reaction time and motor skills may be more applicable to sports performance. The domains chosen for the present study were derived from previous studies in athletes to translate to sports performance. Bandelow et al¹⁵ tested soccer players on the same motor skills test (finger tapping test) and with a choice reaction time test that tested the same cognitive domain as the complex reaction time section of the Stroop test.¹⁵ Triathletes were tested for simple and choice reaction time, which are tested in the Stroop test.²⁸ Track and field athletes were tested on a variety of executive and non-executive tasks to determine the effects of exercise on executive function, which was tested *via* the shifting attention test in the present study.²⁹

Interestingly, Stroop reaction time was the only reaction time test measured by the Stroop test that demonstrated improvements, though it is the most complex. Stroop reaction time measures the responses in the third part of the Stroop test, where a response must be given when the color of the word does not match the word written. This was a novel finding since it suggested that the subjects' ability to respond to rapidly changing and increasingly complex sets of directions was improved, and

other researchers have suggested that an attenuation of exercise-induced cognitive benefits occurs as complexity increases, because more errors were made in tasks that required subjects to shift decision-making rules.¹¹ This was not the case in the present study, in which the most robust effect was observed from the shifting attention test, which was significant even after the Bonferroni correction was applied.

Another area of discrepancy in the literature is the method of dehydration. Various methods of dehydration are employed in studies, and dehydration methods should reflect how dehydration occurs in specific populations. For example, fluid deprivation is probably the most common method by which dehydration occurs in everyday life. In most athletic situations involving dehydration, both heat and exercise play a role. In military, tactical and sport contexts, intense exercise is often performed in hot, humid conditions. Therefore, dehydration protocols may be more applicable to sport when heat and exercise are both employed, even though studies have not demonstrated significant differences between deprivation methods.^{7,8} Since exercise was employed as a method of dehydration in the present study, the cognitive improvements may have been caused by the exercise and not the dehydration. Multiple studies have demonstrated improved cognitive function after exercise.^{9,20,30-32} For example, Nanda et al³⁰ elicited improvements in memory, reasoning, and planning after 30 minutes of cycling at 70% of heart rate reserve.³⁰ Hillman and associates found that amplitude of event-related brain potential increased after 30 minutes of treadmill running,²⁰ which involved a similar exercise bout as employed by the present study. One possible mechanism is that exercise increases blood flow to the brain, which has been linked to improvements in cognitive function.³³ The exercise-induced dehydration may have elicited positive cholinergic effects, as demonstrated in an animal study by Fordyce et al.³⁴ Though exercise may be contributing to the cognitive benefits, it is important to note that not all exercise-induced dehydration studies demonstrated improvements.^{5,33}

There are limitations regarding interpretation of the findings of the present study. First, the combination of exercise and heat stress in the dehydration protocol limits the ability to determine whether exercise or dehydration is causing the improvements, though employing both dehydration methods was important to apply the dehydration to sport contexts. Second, our null results and trends may be non-significant due to small data set size. Third, the subjects were athletic, active individuals, but were not collegiate or professional athletes. Therefore, it is difficult to determine whether the findings would apply to highly athletic men and women. Future studies investigating dehydration and cognitive function need to enlist elite athletes in various sports to support the use of sport-specific cognitive testing methods and dehydration protocols.

CONCLUSION

In summary, mild dehydration caused by both exercise and heat

exposure enhanced Stroop reaction time during the Stroop test. Second, reaction time during the shifting attention test was significantly different after dehydration, even after the Bonferroni correction was applied. These findings have military, tactical, and sport applications, in which athletes and athletic individuals are often performing at various levels of dehydration caused by both heat and exercise, and require optimal cognitive functioning in multiple, specific areas. Also, weight-class athletes may employ both exercise and sauna to achieve weight loss, which may induce quicker reaction times when the dehydration is mild. More research is needed to elucidate the precise areas of cognitive function that improve or decline as a result of mild dehydration, and whether elite athletes will demonstrate the same improvements as athletic individuals.

ACKNOWLEDGEMENTS

We would like to thank MusclePharm Inc., for funding for this study. Citation of commercial organizations or products does not constitute endorsement by the authors.

CONFLICTS OF INTEREST

The authors declared that they have no conflicts of interest.

REFERENCES

- Hellwig JP, Otten JJ, Meyers LD. *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*. Washington DC, USA; National Academies Press; 2006.
- Institute of Medicine of the National Academies. *Electrolytes, and Water, DRI. Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate*. Washington DC, USA: National Academy Press; 2005.
- American College of Sports Medicine, Sawka MN, Burke LM, et al. Exercise and fluid replacement. *Med Sci Sports Exerc*. 2007; 39(2): 377-390. doi: [10.1249/mss.0b013e31802ca597](https://doi.org/10.1249/mss.0b013e31802ca597)
- Judelson DA, Maresh CM, Farrell MJ, et al. Effect of hydration state on strength, power, and resistance exercise performance. *Med Sci Sports Exerc*. 2007; 39(10): 1817-1824. doi: [10.1249/mss.0b013e3180de5f22](https://doi.org/10.1249/mss.0b013e3180de5f22)
- Gopinathan PM, Pichan G, Sharma VM. Role of dehydration in heat stress-induced variations in mental performance. *Arch Environ Health*. 1988; 43(1): 15-17. doi: [10.1080/00039896.1988.9934367](https://doi.org/10.1080/00039896.1988.9934367)
- Sharma VM, Sridharan K, Pichan G, Panwar MR. Influence of heat-stress induced dehydration on mental functions. *Ergonomics*. 1986; 29(6): 791-799. doi: [10.1080/00140138608968315](https://doi.org/10.1080/00140138608968315)
- Cian C, Barraud PA, Melin B, Raphel C. Effects of fluid ingestion on cognitive function after heat stress or exercise-induced dehydration. *Int J Psychophysiol*. 2001; 42(3): 243-251.
- Cian C, Koulmann N, Barraud PA, Raphel C, Jimenez C, Melin B. Influences of variations in body hydration on cognitive function: Effect of hyperhydration, heat stress, and exercise-induced dehydration. *J Psychophysiol*. 2000; 14(1): 29-36. doi: [10.1027//0269-8803.14.1.29](https://doi.org/10.1027//0269-8803.14.1.29)
- Lindseth PD, Lindseth GN, Petros TV, Jensen WC, Caspers J. Effects of hydration on cognitive function of pilots. *Mil Med*. 2013; 178(7): 792-798. doi: [10.7205/MILMED-D-13-00013](https://doi.org/10.7205/MILMED-D-13-00013)
- Smith MF, Newell AJ, Baker MR. Effect of acute mild dehydration on cognitive-motor performance in golf. *J Strength Cond Res*. 2012; 26(11): 3075-3080. doi: [10.1519/JSC.0b013e318245bea7](https://doi.org/10.1519/JSC.0b013e318245bea7)
- Tomprowski PD, Beasman K, Ganio MS, Cureton K. Effects of dehydration and fluid ingestion on cognition. *Int J Sports Med*. 2007; 28(10): 891-896. doi: [10.1055/s-2007-965004](https://doi.org/10.1055/s-2007-965004)
- Leibowitz HW, Abernethy CN, Buskirk ER, Bar-or O, Hennessy RT. The effect of heat stress on reaction time to centrally and peripherally presented stimuli. *Hum Factors*. 1972; 14(2): 155-160. Web site. <http://journals.sagepub.com/doi/abs/10.1177/001872087201400205>. Accessed November 29, 2016.
- Adam GE, Carter R 3rd, Chevront SN, et al. Hydration effects on cognitive performance during military tasks in temperate and cold environments. *Physiol Behav*. 2008; 93(4-5): 748-756. doi: [10.1016/j.physbeh.2007.11.028](https://doi.org/10.1016/j.physbeh.2007.11.028)
- Armstrong LE, Ganio MS, Casa DJ, et al. Mild dehydration affects mood in healthy young women. *J Nutr*. 2012; 142(2): 382-388. doi: [10.3945/jn.111.142000](https://doi.org/10.3945/jn.111.142000)
- Bandelow S, Maughan R, Shirreffs S, et al. The effects of exercise, heat, cooling and rehydration strategies on cognitive function in football players. *Scand J Med Sci Sports*. 2010; 20(Suppl 3): 148-160. doi: [10.1111/j.1600-0838.2010.01220.x](https://doi.org/10.1111/j.1600-0838.2010.01220.x)
- Cullen S, Dolan E, McGoldrick A, et al. The impact of making-weight on cognitive performance in apprentice jockeys. *J Sports Sci*. 2015; 33(15): 1589-1595. doi: [10.1080/02640414.2014.1002104](https://doi.org/10.1080/02640414.2014.1002104)
- Ganio MS, Armstrong LE, Casa DJ, et al. Mild dehydration impairs cognitive performance and mood of men. *Br J Nutr*. 2011; 106(10): 1535-1543. doi: [10.1017/S0007114511002005](https://doi.org/10.1017/S0007114511002005)
- Ely BR, Sollanek KJ, Chevront SN, Lieberman HR, Kenefick RW. Hypohydration and acute thermal stress affect mood state but not cognition or dynamic postural balance. *Eur J Appl Physiol*. 2013; 113(4): 1027-1034. doi: [10.1007/s00421-012-2506-6](https://doi.org/10.1007/s00421-012-2506-6)
- Neave N, Scholey AB, Emmett JR, Moss M, Kennedy DO, Wesnes KA. Water ingestion improves subjective alertness, but has no effect on cognitive performance in dehydrated healthy

- young volunteers. *Appetite*. 2001; 37(3): 255-256. doi: [10.1006/appe.2001.0429](https://doi.org/10.1006/appe.2001.0429)
20. Hillman CH, Snook EM, Jerome GJ. Acute cardiovascular exercise and executive control function. *Int J Psychophysiol*. 2003; 48(3): 307-314. doi: [10.1016/S0167-8760\(03\)00080-1](https://doi.org/10.1016/S0167-8760(03)00080-1)
21. Jackson AS. Estimating maximum heart rate from age: Is it a linear relationship? *Med Sci Sports Exerc*. 2007; 39(5): 821. doi: [10.1249/mss.0b013e318054d3ca](https://doi.org/10.1249/mss.0b013e318054d3ca)
22. Sawka MN, Chevront SN, Kenefick RW. Hypohydration and human performance: Impact of environment and physiological mechanisms. *Sports Med*. 2015; 45(Suppl 1): S51-S60. doi: [10.1007/s40279-015-0395-7](https://doi.org/10.1007/s40279-015-0395-7)
23. Armstrong LE, Soto JA, Hacker FT Jr, Casa DJ, Kavouras SA, Maresh CM. Urinary indices during dehydration, exercise, and rehydration. *Int J Sport Nutr*. 1998; 8(4): 345-355. Web site. <https://www.infona.pl/resource/bwmeta1.element.elsevier-edd78b3e-3323-3fbb-8d6e-050dae3adc03>. Accessed November 29, 2016.
24. Armstrong LE, Maresh CM, Castellani JW, et al. Urinary indices of hydration status. *Int J Sport Nutr*. 1994; 4(3): 265-279.
25. Gualtieri CT, Johnson LG. Reliability and validity of a computerized neurocognitive test battery, CNS Vital Signs. *Arch Clin Neuropsychol*. 2006; 21(7): 623-643. doi: [10.1016/j.acn.2006.05.007](https://doi.org/10.1016/j.acn.2006.05.007)
26. Szinnai G, Schachinger H, Arnaud MJ, Linder L, Keller U. Effect of water deprivation on cognitive-motor performance in healthy men and women. *Am J Physiol Regul Integr Comp Physiol*. 2005; 289(1): R275-R280. doi: [10.1152/ajpregu.00501.2004](https://doi.org/10.1152/ajpregu.00501.2004)
27. Rogers WA, Fisk AD. *Human Factors Interventions for the Health Care of Older Adults*. CRC Press; 2001.
28. Collardeau M, Brisswalter J, Verduyssen F, Audiffren M, Goubault C. Single and choice reaction time during prolonged exercise in trained subjects: Influence of carbohydrate availability. *Eur J Appl Physiol*. 2001; 86(2): 150-156. doi: [10.1007/s004210100513](https://doi.org/10.1007/s004210100513)
29. Luft CD, Takase E, Darby D. Heart rate variability and cognitive function: Effects of physical effort. *Biol Psychol*. 2009; 82(2): 164-168. doi: [10.1016/j.biopsycho.2009.07.007](https://doi.org/10.1016/j.biopsycho.2009.07.007)
30. Nanda B, Balde J, Manjunatha S. The acute effects of a single bout of moderate-intensity aerobic exercise on cognitive functions in healthy adult males. *J Clin Diagn Res*. 2013; 7(9): 1883-1885. doi: [10.7860/JCDR/2013/5855.3341](https://doi.org/10.7860/JCDR/2013/5855.3341)
31. Liu YF, Chen HI, Wu CL, et al. Differential effects of treadmill running and wheel running on spatial or aversive learning and memory: Roles of amygdalar brain-derived neurotrophic factor and synaptotagmin I. *J Physiol*. 2009; 587 (Pt 13): 3221-3231. doi: [10.1113/jphysiol.2009.173088](https://doi.org/10.1113/jphysiol.2009.173088)
32. Lin TW, Chen SJ, Huang TY, et al. Different types of exercise induce differential effects on neuronal adaptations and memory performance. *Neurobiol Learn Mem*. 2012; 97(1): 140-147. doi: [10.1016/j.nlm.2011.10.006](https://doi.org/10.1016/j.nlm.2011.10.006)
33. Critchley HD, Corfield DR, Chandler MP, Mathias CJ, Dolan RJ. Cerebral correlates of autonomic cardiovascular arousal: A functional neuroimaging investigation in humans. *J Physiol*. 2000; 523(Pt 1): 259-270. doi: [10.1111/j.1469-7793.2000.t01-1-00259.x](https://doi.org/10.1111/j.1469-7793.2000.t01-1-00259.x)
34. Fordyce DE, Farrar RP. Enhancement of spatial learning in F344 rats by physical activity and related learning-associated alterations in hippocampal and cortical cholinergic functioning. *Behav Brain Res*. 1991; 46(2): 123-133. doi: [10.1016/S0166-4328\(05\)80105-6](https://doi.org/10.1016/S0166-4328(05)80105-6)