

Research

*Corresponding author

Kingsley Badu, BSc, MPhil, PhD

Department of Theoretical and Applied Biology

Kwame Nkrumah University of Science and Technology, Kumasi, Ghana

E-mail: kingsbadu@gmail.com

Volume 4 : Issue 1

Article Ref. #: 1000HARTOJ4126

Article History

Received: November 17th, 2017

Accepted: December 11th, 2017

Published: December 13th, 2017

Citation

Tay SCK, Aryee ENO, Badu K. Intestinal parasitemia and HIV/AIDS co-infections at varying CD4⁺ T-cell levels. *HIV/AIDS Res Treat Open J*. 2017; 4(1): 40-48. doi: [10.17140/HARTOJ-4-126](https://doi.org/10.17140/HARTOJ-4-126)

Copyright

©2017 Badu K. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Intestinal Parasitemia and HIV/AIDS Co-infections at Varying CD4⁺ T-cell Levels

Samuel CK Tay, BSc, DVM, MSc, MPH¹; Eric Nii Okai Aryee, BSc, MSc²; Kingsley Badu, BSc, MPhil, PhD^{3*}

¹Department of Clinical Microbiology, School of Medical Sciences, College of Health Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana

²Madina Polyclinic- Kekele Ghana Health Service, Madina, Accra, Ghana

³Department of Theoretical and Applied Biology, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana

ABSTRACT

Background: Intestinal parasites, especially coccidian parasites, cause gastrointestinal symptoms such as severe diarrhoea which increases morbidity and mortality rates in people living with Human Immunodeficiency Virus/Acquired Immunodeficiency Syndrome (HIV/AIDS), particularly in Sub-Saharan Africa. We examined the prevalence of intestinal parasites in people living with AIDS at different CD4⁺ T-cell levels.

Method: Case-control studies were conducted over a four month period including a total of 672 participants, between the ages of 8 and 72 years. HIV screening and confirmatory tests were done. We examined stool samples by wet mount, followed by formol-ether concentration and staining with Modified Field's and Ziehl Neelsen techniques. We also carried out fluorescence-activated cell sorting (FACS) analyses to obtain their CD4⁺ T-cell levels.

Results: The prevalence of intestinal parasites were significantly higher (25.2%) among HIV seropositives than HIV seronegative individuals (13.3%), ($p < 0.001$). Coccidian parasites: *Cystoisospora belli* (formerly *Isoospora belli*), *Cryptosporidium* and the round worm *Strongyloides stercoralis* infections were found exclusively in HIV seropositives. *Cryptosporidium* infections were more frequently observed in the rural cohort ($p = 0.039$). *C. belli*, *Cryptosporidium*, *Giardia lamblia* and *Strongyloides stercoralis* infections were significantly higher in diarrhoeic stools. Microsporidia and *Cystoisospora belli* were found mostly in individuals with CD4⁺ T-cell levels of ≤ 200 cells/ μ L. Participants with CD4⁺ T-cell count of ≤ 50 cells/ μ L were associated with diarrhoea.

Conclusion: The prevalence of opportunistic coccidian parasites remains high in HIV-infected individuals with low CD4⁺ T-cell counts. Routine diagnosis is recommended to ensure comprehensive care for HIV patients.

KEY WORDS: HIV/AIDS; Intestinal parasites; *Diarrhoea*, CD4⁺ T-cell count.

ABBREVIATIONS: HIV/AIDS: Human Immunodeficiency Virus/Acquired Immunodeficiency Syndrome; NSAID: Non-Steroidal Anti-Inflammatory Drugs; ART: Antiretroviral Therapy; CI: Confidence Interval.

INTRODUCTION

Human Immunodeficiency Virus/Acquired Immunodeficiency Syndrome (HIV/AIDS) and intestinal parasites co-infections are linked in a vicious cycle which, results in a major public health burden for developing countries.¹ Currently, about 23.5 million people, an estimated 69% of all people living with HIV and AIDS, live in sub-Saharan Africa.^{2,3} Moreover, 90-92% of all pregnant women and children living with HIV are reportedly living on the continent.^{3,4}

In Ghana, the HIV prevalence is relatively low, but the rising trend in the last three years is a matter of concern.⁵ According to HIV sentinel survey report for 2016, the national

prevalence is estimated to be 2.4% representing “a second consecutive upsurge from the 2014 prevalence of 1.6 % and 1.8 % in 2015”.⁵ The survey also found that HIV prevalence is higher in urban areas (2.5%) than rural (1.9%). New infections remained unchanged at 1.1%. The prevalence of the disease was highest among the 45-49 age groups at 5.6%, followed by 35-39 year group at 3.5% whilst 15-19 being the lowest at 0.6%.⁵ The survey also found that the proportion of HIV subtype 1 is 98.5% compared to 1.5% for dual HIV type 1 and 2 infections but no subtype 2 sole infections.⁵

Although, antiretro viral therapy is available in Ghana, the coverage hardly reaches all the patients who need them. According to the Centers for Disease Control and Prevention (CDC), 66,366 adults were receiving Antiretroviral Therapy (ART) at the coverage of 62% by 2012 (<http://www.cdc.gov/globalaids/global-hiv-aids-at-cdc/countries/ghana/default.html>). In the 2015 UN Sustainable Development Goals, the world committed itself to stopping HIV/AIDS by 2030, it is therefore imperative that infections that aggravate disease progression, such as intestinal parasites are closely investigated.

HIV/AIDS infection is associated with high prevalence of gastrointestinal infections including parasite-associated diarrhoea due to apparent dysfunctional immunity.⁶⁻⁸ The progressive decline of the mucosal immunologic defense mechanisms in HIV/AIDS patients predisposes them to early, intermediary, or late gastrointestinal infections.⁹ Progressive HIV infection have been characterized by an increase in Th2-like responses, which may either be a consequence or a cause of the immune deterioration.¹⁰⁻¹⁵ This has been shown to be responsible for increasing host susceptibility to a myriad of intestinal opportunistic agents, such as *Cryptosporidium parvum*, *Cystoisospora belli* and *Microsporidia* species.¹⁶ Diarrhoea affects up to 90% of HIV patients, increasing in frequency and magnitude as the disease progresses.^{17,18} Helminths are known to cause T-cell dysfunction, thereby worsening the already compromised immune system.¹⁹ Diarrhoea, as a result, has been recognized as one of the most frequent causes of morbidity and death in people living with HIV.²⁰ At the national level, data on intestinal parasites in people living with HIV/AIDS is virtually non-existent since there are no specific guidelines that require standard investigation and diagnosis of intestinal parasitic infections in HIV patients. The objective of the study was to determine the patterns of intestinal parasitic infections in people living with HIV/AIDS, and its association to diarrhoea at different CD4⁺ T cell levels.

METHODS

Study Area

This study was based on two cross-sectional surveys conducted at two hospitals in the Ashanti region of Ghana which is known to have HIV prevalence of 2.6%.⁵ The surveys were conducted at HIV/AIDS Voluntary Counseling and Testing centers (VCT) at the two hospitals. The Nyinahin Government Hospital was

located in a rural area and St. Patrick’s Hospital at Offinso was located in a peri-urban area. The surveys were carried out between April and July, 2011.

The study was conducted in two districts of the Ashanti region (forest zone) in the middle belt of Ghana; the Atwima Mponua District and the Offinso Municipality. The Atwima Mponua District is located in the western part of Ashanti Region, and lies between longitude 2°00'W and 2°32'W and latitude 6°32'N and 6°75'N. The district is within the wet-semi equatorial zone and has two peaks of rainfall in the year. About 92% of the people reside in the rural countryside, with only about 8% living within an urban settlement. The Offinso Municipality, on the other hand, is located in the north-western stretch of Ashanti. It is about 40 km away from the regional capital of Kumasi, and lies between longitude 1°65'W and 1°45'E and latitudes 6°45'N and 7°25'S. The district covers an area of 1255 km². The high population growth rate in these localities can be attributed to high immigration and the spillover population from the Kumasi metropolis giving rise to about 40% of the populace in urban dwellings. The district represents a peri-urban settlement. Nyinahini and Offinso are district capitals of the Atwima Mponua District and Offinso Municipality, Ashanti Region, Ghana, respectively.

Sample Size

This was determined based on assumptions of the binomial distribution to estimate the confidence interval (CI) of HIV prevalence at 95% with an estimated population size of less than 3000, described earlier.^{21,22} $n = z^2 (pq)/d^2$, where z is the critical value of standard normal distribution, p is the baseline parasite prevalence and d is the level of precision.^{21,22} For 95% CI and precision level of 5%, the minimum number of participants required to achieve adequate statistical power was 323.

Inclusion and Exclusion Criteria

All patients attending the VCT clinic, who were at least one year old, regardless of their HIV status (seropositive or negative) and gender, and not on non-steroidal anti-inflammatory drugs (NSAID) were eligible to be included in the study. Informed consent was sought from all eligible patients, and participants were only recruited when consent was given.

Study Participants and Recruitment

The study took place in two hospitals, (The Nyinahin Government hospital and St. Patrick’s Hospital) specifically at HIV VCT clinic.

Researchers visited the clinics twice per week until desired sample size was realized. Researchers approached all persons attending the clinic including family members for consent and participation. The study was duly explained to them and participation was purely voluntary. Informed consent mainly adult

consent forms and parental/guardian/assent consent forms were administered.

A total of 341 HIV patients (seropositives) consented to the study and these constituted the case group. Another 331 HIV seronegative gave their consent to participate in the study; this group constituted the negative control group. Demographic data and antiretroviral drug usage information were obtained for each individual who consented to the study. Each participant was provided with sterile screw-capped containers to provide stool samples (collected in the morning) on 2 consecutive days during their scheduled visits. The stool samples were examined for ova, larvae and cyst of parasites regardless of the presence of diarrhoea. In addition, 3 mL of blood taken from participants were put in ethylenediaminetetraacetic acid (EDTA) anti-coagulated tube for CD4⁺ T-cell count.

HIV Testing

Screening: All consenting study participants regardless of the knowledge of their status were screened for the presence or otherwise of HIV-1 and HIV-2 or both, using the First Response HIV Card Test (PMC Ltd, Shree Indl Estate, India). Briefly, (according to manufacturer's instructions) 10 µl of serum was dropped into the sample well and 35 µl of assay diluent was added. The results were read and interpreted within 5-15 minutes. The presence of only one band in the control line in the result window indicates a negative result. However, two-color bands, one control and the other for HIV-1 indicated reactivity for antibodies to HIV-1. Two-color bands, one control and the other for HIV-2 indicated reactivity for antibodies to HIV-2. All three color bands indicated reactivity for antibodies to HIV-1 and HIV-2.

Confirmation: All seropositive individuals were re-tested with the Qualitative Immunoassay test for confirmation of their HIV status. The HIV-1/2 Oral Quick Rapid Test (OraSure Technologies, Inc., Bethlehem, PA 18015, USA) was used according to manufacturer's instructions. Briefly, the rapid test device was removed from its pouch and the padded-end was used to swab the upper and lower gums of participants. The padded-end of the rapid test device was inserted into the buffer, and the test result was read and interpreted after 20 minutes.

Stool Examination

A total of 1,344 stool samples were obtained from 672 participants. These were subjected to routine stool examination, which included saline and iodine mounts to screen for helminth ova and larvae, protozoan cyst, and trophozoites.

Direct wet mount of stool sample in normal saline (0.85% NaCl) was prepared immediately upon arrival in the laboratory and examined under light microscopy (x10 and x40 objectives) for the presence of vegetative forms, larvae, and ova of helminthes. Field's stain and Lugol's iodine staining was used

to detect *Giardia lamblia* flagellates and cysts of protozoa, respectively. The formol-ether concentration technique was employed to concentrate stool samples for further confirmatory microscopic examination. Examination of fecal smears after special staining (Modified Zhiel Neelsen and Modified Field's staining techniques for the detection of *Cryptosporidium*, *Isospora belli*, *Cyclospora cayetanensis* and Microsporidia spores, respectively) was done according to Chessbrough.²³

CD4⁺ T-cell Count

The fluorescence-activated cell sorting (FACS) count (Becton Dickinson Immunocytometry system, Singapore) was used for the immunophenotyping of lymphocytes. Briefly, CD4 reagent tubes were vortexed and opened with the coring station for 50 µl of whole blood to be added. These were vortexed and incubated for one hour in the dark at room temperature. The tubes were uncapped and 50 µl of fixative added. The tubes were recapped and vortexed for 5 seconds upright before subjecting it to the FACS Count instrument for the immunophenotyping of lymphocytes.

Data Analysis

Statistical analysis was carried out using SPSS (2007) version 17.0. Data was summarized using frequency tables. The proportions of parasites were compared between the CD4⁺ T-cell counts with chi-square test. The relationship between the CD4⁺ T-cell count and the presence of diarrhoea were assessed using the chi-square analysis with significance level set at 0.05.

Ethical Considerations

The study was conducted with the approval of the Committee on Human Research Publication and Ethics (CHRPE) of the School of Medical Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.

RESULTS

Demographic Characteristics

A total of 672 patients gave their consent to participate in the study; these comprised of 341 HIV seropositives and 331 seronegatives. Overall, there were 81.5% females (n=548) 18.5% males (n=124). The percentage of male and female participants under each study group (seropositive or seronegative) was similar, and followed the same trend as that mentioned for the overall percentages (Figure 1).

When the study population was stratified into age groups, it was realized that majority of the HIV seropositives 65% (n=225) belonged to the most productive age bracket 25-45 years. Only 3.2% (n=11) belonged to the paediatric population, that is less or equal to 14 years (Figure 2).

Parasite Prevalence among HIV Seropositive and Seronegative





Table 1: Parasite Prevalence among HIV Seropositive and Seronegative Participant.

Parasite	HIV seropositive N=341 (%)	HIV seronegative N=331 (%)	Total (N=672)	p-value
<i>Giardia lamblia</i>	39 (11.4)	39 (11.8%)	78 (11.6)	0.905
<i>Entamoeba histolytica</i>	4 (1.2)	0 (0)	4 (0.6)	0.124
<i>Cystoisospora belli</i> (formerly <i>Isospora belli</i>),	9 (2.6)	0 (0)	9 (1.3)	0.004
<i>Cryptosporidium</i> spp.	7 (2.1)	0 (0)	7 (1.0)	0.032
Microsporidia	1 (0.3)	0 (0)	1 (0.1)	1
<i>Cyclospora cayetanensis</i>	1 (0.3)	0 (0)	1 (0.1)	1
<i>Strongyloides stercoralis</i>	13 (3.8)	0 (0)	13 (1.9)	<0.001
<i>Enteriobius vermicularis</i>	1 (0.3)	1 (0.3)	2 (0.3)	1
<i>Ascaris lumbricoides</i>	0 (0)	1 (0.3)	1 (0.1)	0.493
Hookworm	3 (0.9)	2 (0.6)	5 (0.7)	1
<i>G. lamblia</i> and <i>E. coli</i>	1 (0.3)	0 (0)	1 (0.1)	1
<i>G. lamblia</i> and <i>E. histolytica</i>	4 (1.2)	1 (0.3)	5 (0.7)	0.373
<i>C. belli</i> and <i>S. stercoralis</i>	1 (0.3)	0 (0)	1 (0.1)	1
Microsporidia and <i>C. belli</i>	2 (0.6)	0 (0)	2 (0.3)	0.499

HIV+=HIV Positive; HIV=HIV Negative; N=Number of Participants, *Coccidian parasites: p-value of <0.001 with HIV positive participants.

Participants

The overall prevalence of intestinal parasites among the study subjects was 19.3%. A total of 130 participants had at least one parasite. *Giardia lamblia* was the most common parasite encountered, and had similar prevalence in HIV seropositives (11.4%)

and seronegatives (11.8%) (Table 1). Parasite prevalence was significantly higher in HIV seropositive participants (25.2%), than the control group (13.3%) [$\chi^2=15.3, p=0.000$ (Table 1)]. This confirms the susceptibility of HIV patients to opportunistic parasitic infections.

Among the HIV seropositive cohort, 21 (6.2%) were infected with the coccidian parasites (*Cystoisospora belli*, *Cryptosporidium*, *Cyclospora cayetanensis*) including the fungi-like unicellular intracellular parasite Microsporidia. There were single infections as well as co infections with more than one parasite. Parasites such as *Cystoisospora belli* (2.6%), *Cryptosporidium sp* (2.1%), Microsporidia (0.3%) and *C. cayetanensis* (0.3%) as well as *S. stercoralis* were single infections exclusively found with HIV seropositives. The predominant helminths was *S. stercoralis* (3.8%) followed by Hookworm (0.9%) and *E. vermicularis* (0.3%) among seropositive participants. Again, there were co-infections with *Giardia lamblia* and *E. coli*, *Giardia lamblia* and *Entamoeba histolytica*, *C. belli* and *S. stercoralis* as well as Microsporidia and *C. belli* albeit in low prevalence. *Ascaris lumbricoides* occurred in only 1 HIV negative participant (0.3%) Table 1.

CD4⁺ T-cell Levels among Parasite and HIV Co-infected Participants

Using CD4⁺ T-cell estimate as a marker of relative risk of developing HIV related opportunistic infections,²⁴ we observed 29.3% of HIV seropositives were in the acute (asymptomatic) infection stage, 44.3% were in the intermediate (symptomatic) stage, and

19.7% were in the late (symptomatic) disease stage. Still 6.7% were in the most advanced HIV disease stage (Table 2). *Giardia lamblia* infections were found at all CD4⁺ T-cell levels with lower prevalence (4.3%) among the participants with the least CD4⁺ T-cell count (<50 cells/μl), but this was not statistically significant ($p=0.852$) (Table 2). *C. belli*, Microsporidia, and *C. cayetanensis* were also found in patients with the least CD4⁺ T-cell count. Although, *S. stercoralis* occurred exclusively among HIV seropositives ($p<0.001$) (Table 1), its occurrence was not significantly associated with CD4⁺ T-cell level.

CD4⁺ T-Lymphocyte Levels and Diarrhoea

Out of 341 participants belonging to the HIV seropositive group, 110 presented diarrhoeic stools; representing 32.3%. Intestinal parasites were observed in 32.7% of the diarrhoeal stools. The highest incidence of diarrhoeic stools (78.3%) was observed among participants with CD4⁺ T-cell count <50 cells/μL (Table 3). Contrary, only 2% of participants with CD4⁺ T-cell count of ≥500 cells/μL presented diarrhoeal stools.

Coccidian parasites were detected more commonly in HIV seropositives with diarrhoea than in participants with hel-

Table 2: CD4⁺ T-cell Levels among Parasite and HIV Co-Infected Participants.

Parasite	CD4 ⁺ T- cell count (cells/ul)				p-value
	>500 (N=100) (29.3%)	200-500 (N=151) (44.3%)	50-200 (N=67) (19.6%)	<50 (23) (6.7%)	
<i>Giardia lamblia</i>	50 (11.9%)	16 (9.9%)	11 (16.4%)	1 (4.3%)	0.852
<i>Entamoeba histolytica</i>	0 (0%)	3 (1.9%)	1 (1.5%)	0 (0%)	0.079
<i>Cystoisospora belli</i>	0 (0%)	0 (0%)	3 (4.5%)	6 (26.1%)	<0.001
<i>Cryptosporidium spp.</i>	2 (0.5%)	1 (0.6%)	3 (4.5%)	1 (4.3%)	0.442
Microsporidia	0 (0%)	0 (0%)	0 (0%)	1 (4.3%)	0.002
<i>Cyclospora cayetanensis</i>	0 (0%)	0 (0%)	0 (0%)	1 (4.3%)	0.002
Hookworm	4 (1.0%)	1 (0.6%)	0 (0%)	0 (0%)	0.343
<i>Strongyloides stercoralis</i>	2 (0.5%)	5 (3.1%)	5 (7.5%)	1 (4.3%)	<0.001
<i>Enteriobius vermicularis</i>	1 (0.2%)	0 (0%)	1 (1.5%)	0 (0%)	0.423
<i>Ascaris lumbricoides</i>	0 (0%)	1 (0.6%)	0 (0%)	0 (0%)	0.571
<i>G. lamblia</i> and <i>E. coli</i>	1 (0.2%)	0 (0%)	0 (0%)	0 (0%)	0.503
<i>Cystoisospora belli</i> and <i>S. stercoralis</i>	0 (0%)	0 (0%)	1 (1.5%)	0 (0%)	<0.001
Microsporidia and <i>Cystoisospora belli</i>	0 (0%)	0 (0%)	0 (0%)	2 (8.7%)	<0.001
<i>G. lamblia</i> and <i>E. histolytica</i>	1 (0.2%)	2 (1.2%)	2 (3.0%)	0 (0%)	0.068

HIV⁺=HIV Positive; HIV⁻=HIV Negative; N=Number of Participants; ND=Not Determined.

Table 3: Correlation between CD4⁺ T-Lymphocyte Levels and Diarrhoea.

	CD4 ⁺ T-cell count(cells/L)				Total	p-value
	>500	200-500	50-200	<50		
Diarrhoea Stools	2 (2%)	6 (4.0%)	23 (34.3%)	18 (78.3%)	49 (14.4%)	<0.001
Non Diarrhoea Stools	98 (98%)	145 (96.0%)	44 (65.7%)	5 (21.7%)	292 (85.6%)	
Total	100 (100%)	151 (100%)	67 (100%)	23 (100%)	341 (100%)	

Table 4: Presence of Parasites in Diarrhoeic Stools of HIV Positive and Negative Participants.						
Parasite	HIV* (N=341)			HIV (N=331)		
	D	ND	p-value	D	ND	p-value
<i>Giardia lamblia</i>	10 (40.0%)	29 (9.2%)	<0.001	18 (62.1%)	21 (7.0%)	<0.001
<i>Entamoeba histolytica</i>	0 (0%)	4 (1.3%)	0.736	0 (0%)	0 (0%)	nd
<i>Cystoisospora belli</i>	8 (16.3%)	1 (0.3%)	<0.001	0 (0%)	0%	nd
<i>Cryptosporidium</i> spp.	3 (6.1%)	3 (1.0%)	0.04	0 (0%)	0 (0%)	nd
Microsporidia	1 (2.0%)	0 (0%)	0.14	0 (0%)	0 (0%)	nd
<i>Cyclospora cayetanensis</i>	1 (2.0%)	0 (0%)	0.14	0 (0%)	0 (0%)	nd
Hookworm	0 (0.0%)	3 (1.0%)	1	0 (0%)	2 (0.6%)	1
<i>Strongyloides stercoralis</i>	5 (10.2%)	8 (2.7%)	0.026	0 (0%)	0 (0%)	nd
<i>Enteriobius vermicularis</i>	1 (2.0%)	0 (0%)	0.14	0 (0%)	1 (0.3%)	1
<i>Ascaris lumbricoides</i>	0 (0%)	0 (0%)	nd	0 (0%)	1 (0.3%)	1
<i>G. lamblia</i> and <i>E. coli</i>	2 (4.1%)	2 (0.7%)	1	0 (0%)	1 (0.3%)	1
<i>Cystoisospora belli</i> and <i>S.stercoralis</i>	1 (2.0%)	0 (0%)	0.14	0 (0%)	0 (0%)	nd
Microsporidia and <i>Cystoisospora belli</i>	2 (4.1%)	0 (0%)	0.02	0 (0%)	0 (0%)	nd
<i>G. lamblia</i> and <i>E. histolytica</i>	2 (4.1%)	2 (0.7%)	0.1	0 (0%)	1 (0.3%)	1
Total diarrhoea (143) (21.3%)	36/110 (32.7%)	52/283 (18.3%)		18/33 (54.5%)	27/325 (8.3%)	

HIV*=HIV Positive; HIV=Negative; N=Number of Participants; nd=Not Determined; D=Diarrhoea Stools; ND=Non Diarrhoeal Stools.

minths and other protozoa (Table 4). The incidence of diarrhoea among participants infected with *Giardia lamblia* only, was 40% and 62.1% in HIV seropositive and seronegatives respectively. The incidence of diarrhoea among HIV seropositives infected with *C. belli*, *Cryptosporidium* and *S. stercoralis* infections were 16.3%, 6.1% and 10.2%, respectively. Microsporidia, *C. cayetanensis* and *E. vermicularis* occurred with diarrhoea in at most 2% of participants.

DISCUSSION

Intestinal parasitic infections and HIV/AIDS have been the major public health problems and remain a vital cause of morbidity and mortality in developing countries. Both problems are linked in a vicious cycle.¹ The introduction of antiretroviral therapy has lessened the prevalence of gastrointestinal infections in HIV patients. This notwithstanding, several people with HIV infection still suffer from intestinal parasite infections.^{1,6-8,25}

Intestinal parasitic infections found among HIV patients from low-income countries have been reported with prevalence 15% to over 80% in recent times.^{1,6-8,25}

In the current study, the overall prevalence of intestinal parasite among the study population was 19.3%. However, the prevalence of intestinal parasites in the HIV seropositive group was significantly higher (25%) than that observed in the HIV seronegative group which was 13.3%. The observed prevalence in this study is similar to others carried out in Zambia which reported 25% prevalence among HIV-infected cohort.²⁶ Other reports from India, Ethiopia and Tanzania were comparably higher ranging from 30% and above.^{27,28} However, much lower prevalence

of 10.6% among HIV patients have also been reported elsewhere.²⁹

It was also observed that opportunistic parasitic infections mainly the coccidian parasites occurred exclusively in HIV/AIDS patients with a corresponding depletion of CD4⁺ T cell count. This has been attributed to the modulation of immune response in the advance stages of the disease.¹⁰⁻¹⁵ The highest prevalence of parasitosis was observed among participants in the CD4⁺ T-cell level ≤ 50 cells/ μ L. This category forms 56.5% of participants in the advanced stage of the disease. The most predominant parasites recovered among this group of participants belonged to the coccidian groups (47.8%), which are well known as opportunistic parasites in HIV disease. With the exception of one participant, all participants with mixed parasitic infections had CD4⁺ T-lymphocyte level < 200 cells/ μ L. This observation has been echoed by other studies.^{6,30}

Typically, the dynamics of HIV-1 infection is known to follow a familiar pattern where there is the acute phase in which there is massive depletion of CD4⁺ T cells of the gastrointestinal tract,³¹ followed by the chronic phase, where there is a gradual reduction in CD4⁺ T cells which results in high risk of opportunistic infections, and then AIDS sets in. Recently it has been found that there is significant preferential loss of Th17 cells within the gastrointestinal tract of HIV-infected individuals³² as a result of microbial translocation after the initial structural and immunological disruption of the gut mucosa in the acute phase.³³ *Giardia lamblia* was the most common parasite among the participants. Its occurrence, among both HIV seropositive (11.4%) and seronegative (11.8%) was similar. Previous studies have demonstrated that although infection with *Giardia lamblia*

and HIV correlated with enteritis or enterocolitis, its incidence does not differ amongst HIV-positive and negative patient populations.³⁴ This underscores the non-opportunistic nature of the *G. lamblia* reviewed by Cimerman et al.³⁵

The helminths observed in this study were *A. lumbricoides*, *E. vermicularis*, hookworm and *S. stercoralis*. Helminths infections generally were low among the study groups when compared to findings of similar studies elsewhere reporting prevalence of 37.04%.³⁶ However, *S. stercoralis* was only associated with HIV seropositive individuals and hookworm infections were higher (5%). Modjarrad et al³⁶, reported relatively higher prevalence of intestinal helminths (24.9%) with *A. lumbricoides* and hookworm being prevalent among HIV-1 patients in an urban African setting.³⁷ Apart from *S. stercoralis*, other helminths had lower prevalence in our study when compared to others carried out in similar developing countries²⁰; this may be due to the widespread administration of anti-helminths and cotri-moxazole among the study participants (56% of participants were already on ART at the time of stool collection).

Studies have shown that, reconstitution of the immune system following ART administration alone resolves *Cryptosporidium* infections without specific treatment for the parasite.^{38,39} This is because ART acts against the aspartyl-protease of the parasite depriving the parasite of an essential protein.^{40,41} More than 56% of participants were already on ART at the time of stool collection. It is likely that as a patient's CD4⁺ T-cell level increases with the administration of ART, opportunistic infections are not established even if they are exposed to infection.

Diarrhoea is a life threatening complication often associated with HIV, causing severe weight loss; both conditions are independent predictors of mortality in HIV/AIDS.⁴² The incidence of diarrhoea among HIV seropositives in this study was significantly high. Among the HIV cohort diarrhoea episodes increased with declining immunity, with the highest diarrhoea prevalence (78.3%) occurring at the least CD4⁺ T-cell count <50 cells/ μ L. (Table 4). *Giardia lamblia*, *Cystoisospora belli*, *Cryptosporidium*, and *S. stercoralis* were associated with diarrhoeal stools of HIV seropositive patients (Table 4). Among the opportunistic coccidian parasites in HIV seropositives *Cystoisospora belli* (3.5%) was predominant followed by *Cryptosporidium* (2.1%). Microsporidia and *C. cayetanensis* had a prevalence of 0.9% and 0.3%, respectively, these occurred exclusively among HIV seropositives. All participants with *Cystoisospora belli* infections presented with diarrhoea. This strong association with diarrhoea may be associated with patients who were ART naïve and presenting themselves very late to the hospitals. They often present with wasting, general weakness and diarrhoea. *Cyclospora cayetanensis*, an emerging parasite, was found in only one participant with diarrhoea.

On the other hand, the presence of diarrhoea without parasites in stool can be quite intriguing. About 32.7% of HIV seropositives had diarrhoea even in the absence of parasites (Table 4). This can arise from bacteria etiology, lactose intoler-

ance or insufficient sensitivity of the diagnostic procedure.³⁹ It has been shown however, that no etiological agent is found in 15-50% of HIV patients with chronic diarrhoea.⁴¹ Munnink et al observed that unexplained diarrhoea in HIV- infected patients were not due to novel pathogens [immunodeficiency-associated stool virus (IAS virus)]⁴² or previously unknown pathogens, but may be due to HIV-1 itself having a “virotoxic” effect on the enterocytes that results in intestinal mucosal abnormalities leading to diarrhoea.⁴⁰ Again some anti-retroviral agents especially the protease inhibitors have been reported to cause diarrhoea.⁴³ This appears to be changing the etiology of diarrhoea in some parts of the world where parasite related diarrhoea appear to be falling, while the number of unexplained and drug-induced diarrhoea seems to be going up.⁴⁴ It is thus conceivable to state that the interpretation of diarrhoea associated with parasitic infections must be made cautiously.

In spite of the high prevalence (25%) of intestinal parasitosis in HIV patients, there are currently no clear guidelines that require its diagnosis. Moreover, the high burden of intestinal parasitosis results in diarrhoea and weight loss which are independent predictors of mortality in HIV patients. In order for HIV patients to obtain comprehensive healthcare, it is recommended that efforts are made towards diagnosing intestinal parasites in HIV patients especially those with CD4⁺ T cell counts less than 50 cells/ μ L.⁴⁵

COMPETING INTEREST

The authors declare no competing interest.

AUTHOR'S CONTRIBUTION

SCKT and KB conceived the study, participated in its design, supervised the field work, data analysis and drafted the manuscript; EOA conducted the field and laboratory data collection and performed the experiments. All authors read and approved the final manuscript.

ACKNOWLEDGEMENTS

The authors thank the Medical superintendent of the Atwima-Mponua District Hospital, Dr. Yusuf Al-Hassan and the nurses at the VCT Clinic of the St. Patrick's Hospital at Offinso all in the Ashanti region. Sincere gratitude also goes to all study participants for their patience and time.

We are also grateful to Dr. Jewelna Arkoli (Noguchi Memorial Institute, University of Ghana) for her kind review.

REFERENCES

- Gedle D, Kumera G, Eshete T, Ketema K, Adugna H, Feyera F. Intestinal parasitic infections and its association with under nutrition and CD4 T cell levels among HIV/AIDS patients on HAART in Butajira. *J Health Popul Nutr.* 2017; 36: 15. doi:

10.1186/s41043-017-0092-2

2. UNAIDS. Global facts and figures: The global AIDS epidemic. Web site. http://data.unaids.org/pub/globalreport/2008/20080715_fs_global_en.pdf. Accessed May 21, 2014.
3. WHO. Facts about health in the African Region of WHO. Web site. http://www.allcountries.org/health/facts_about_health_in_the_african_region_of_who.html. Accessed May 21, 2014.
4. UNAIDS Gap Report, 2014. Web site. http://files.unaids.org/en/media/unaids/contentassets/documents/unaidspublication/2014/UNAIDS_Gap_report_en.pdf. Accessed May 21, 2014.
5. The National AIDS Control Programme 2016 HIV sentinel report, May 2017.
6. Cimerman S, Cimerman B, Lewi DS. Prevalence of intestinal parasitic infections in patients with acquired immunodeficiency syndrome in Brazil. *Int J Infect Dis*. 1999; 3: 203-206. doi: [10.1016/S1201-9712\(99\)90025-5](https://doi.org/10.1016/S1201-9712(99)90025-5)
7. Sarfati C, Bourgeois A, Menotti J, et al. Prevalence of intestinal parasites including microsporidia in human immunodeficiency virus-infected adults in Cameroon: A cross-sectional study. *Am J Trop Med Hyg*. 2006; 74: 162-164. doi: [10.4269/ajtmh.2006.74.162](https://doi.org/10.4269/ajtmh.2006.74.162)
8. Nsagha DS, Njunda AL, Assob NJC, et al. Intestinal parasitic infections in relation to CD4⁺ T cell counts and diarrhea in HIV/AIDS patients with or without antiretroviral therapy in Cameroon. *BMC Infect Dis*. 2016; 16: 9. doi: [10.1186/s12879-016-1337-1](https://doi.org/10.1186/s12879-016-1337-1)
9. McGowan I, Elliott J, Fuerst M, et al. Increased HIV-1 mucosal replication is associated with generalized mucosal cytokine activation. *J Acquir Immune Defic Syndr*. 2004; 37: 1228-1236. doi: [10.1097/01.qai.0000131846.12453.29](https://doi.org/10.1097/01.qai.0000131846.12453.29)
10. MacDonald AS, Araujo MI, Pearce EJ. Immunology of parasite helminths infections. *Infect Immun*. 2002; 70: 427-433. doi: [10.1128/IAI.70.2.427-433.2002](https://doi.org/10.1128/IAI.70.2.427-433.2002)
11. Osakwe CE, Bleotu C, Chifiriuc MC, et al. TH1/TH2 cytokine levels as an indicator for disease progression in human immunodeficiency virus type 1 infection and response to antiretroviral therapy. *Roum Arch Microbiol Immunol*. 2010; 69(1): 24-34.
12. Evans TG, Fitzgerald T, Gibbons DC, Keefer MC, Soucier H. The Aids Vaccine Evaluation Group. Th1/Th2 cytokine responses following HIV-1 immunization in seronegative volunteers. *Clin Exp Immunol*. 1998; 111(2): 243-250. doi: [10.1046/j.1365-2249.1998.00486.x](https://doi.org/10.1046/j.1365-2249.1998.00486.x)
13. Clerici M, Stocks NI, Zajac RA, et al. Detection of three distinct patterns of T helper cell dysfunction in asymptomatic, human immunodeficiency virus-seropositive patients. *J Clin Invest*. 1989; 84(6): 1892-1899. doi: [10.1172/JCI114376](https://doi.org/10.1172/JCI114376)
14. Maggi E, Mazzetti M, Ravina A, et al. Ability of HIV to promote a TH1 to TH0 shift and to replicate preferentially in TH2 and TH0 cells. *Science*. 1994; 265(5169): 244-248. doi: [10.1126/science.8023142](https://doi.org/10.1126/science.8023142)
15. Clerici M, Hakim FT, Venzon DJ, et al. Changes in interleukin-2 and interleukin-4 production in asymptomatic, human immunodeficiency virus-seropositive individuals. *J Clin Invest*. 1993; 91: 759-765. doi: [10.1172/JCI116294](https://doi.org/10.1172/JCI116294)
16. Soares RLS, Camillo-Coura L, Magalhães LF, Carvalho TM, Gutstein da FA, Oliveira da FK. Isosporiase como causa freqüente de diarreia crônica em pacientes com AIDS em nosso meio [In Portuguese]. *Ann Acad Nac Med*. 1996; 156: 24-25.
17. Awole M, Gebre-Selassie S, Kassa T, Kibru G. Prevalence of intestinal parasites in HIV-infected adult Patients in South-western Ethiopia. *Ethiop J Health Dev*. 2003; 17: 71-78. doi: [10.4314/ejhd.v17i1.9783](https://doi.org/10.4314/ejhd.v17i1.9783)
18. Smith P, Lane H, Gill V, Manischewitz J, Quinnan G, Fauci A. Intestinal infections in patients with the acquired immunodeficiency syndrome (AIDS) Etiology and response to therapy. *Ann Intern Med*. 1998; 108: 328-333. doi: [10.7326/0003-4819-108-3-328](https://doi.org/10.7326/0003-4819-108-3-328)
19. Soave R, Johnson W. *Cryptosporidium* and *Isospora belli* infections. *J Infect Dis*. 1988; 2(1): 225-229. doi: [10.1093/infdis/157.2.225](https://doi.org/10.1093/infdis/157.2.225)
20. Mohandas K, Sehgal R, Sud A, Malla N. Prevalence of intestinal parasitic pathogens in HIV seropositive patients in Northern India. *Jpn J Infect Dis*. 2002; 55(Suppl 3): 83-84.
21. Munyekenye OG, Githeko AK, Zhou G, Mushinzimana E, Minakawa N, Yan G. Plasmodium falciparum spatial analysis, western Kenya highlands. *Emerg Infect Dis*. 2005; 11(10): 1571-1577. doi: [10.3201/eid1110.050106](https://doi.org/10.3201/eid1110.050106)
22. Tay SCK, Badu K, Mensah AA, Gbedema SY. The prevalence of malaria among HIV seropositive individuals and the impact of the co-infection on their hemoglobin levels. *Ann Clin Microbiol Antimicrob*. 2015; 14: 10. doi: [10.1186/s12941-015-0064-6](https://doi.org/10.1186/s12941-015-0064-6)
23. Chessbrough M. *District Laboratory Practice in Tropical Countries*. 3rd ed. part 1. United Kingdom: Cambridge University Press; 2005: 178-215.
24. Singh HR, Singh NG, Singh TB. Estimation of CD4⁺ and CD8⁺ T-lymphocytes in human immunodeficiency virus infection and acquired immunodeficiency syndrome patients in Mani-

- pur. *Indian Journal of Microbiology*. 2007; 25(2): 126-132. doi: [10.4103/0255-0857.32718](https://doi.org/10.4103/0255-0857.32718)
25. Monkemuller KE, Call SA, Lazenby AJ, Wilcox CM. Declining prevalence of opportunistic gastrointestinal disease in the era of combination antiretroviral therapy. *Am J Gastroenterol*. 2000; 95: 457-462. doi: [10.1111/j.1572-0241.2000.01768.x](https://doi.org/10.1111/j.1572-0241.2000.01768.x)
26. Kelly P, Baboo K, Wolff M, Ngwenya B, Luo N, Farthing M. The prevalence and aetiology of persistent diarrhoea in adults in urban Zambia. *Acta Trop*. 1996; 61: 183-190. doi: [10.1016/0001-706X\(95\)00142-2](https://doi.org/10.1016/0001-706X(95)00142-2)
27. Gupta S, Narang V, Singh S. Chronic diarrhoea in HIV patients. Prevalence coccidian parasites. *Indian J Med Microbiol*. 2008; 26(Suppl 2): 172-175. doi: [10.4103/0255-0857.40536](https://doi.org/10.4103/0255-0857.40536)
28. Fontanet A, Sahlou T, Rinke de Wit T, et al. Epidemiology of infections with intestinal parasites and human immunodeficiency virus (HIV) among sugar estate residents in Ethiopia. *Ann Trop Med Parasitol*. 2000; 94: 269-278. doi: [10.1080/00034983.2000.11813539](https://doi.org/10.1080/00034983.2000.11813539)
29. Tian LG, Chen JX, Wang TP, et al. Co-infection of HIV and intestinal parasites in rural area of China. *Parasit Vectors*. 2012; 5: 36. doi: [10.1186/1756-3305-5-36](https://doi.org/10.1186/1756-3305-5-36)
30. Adamu H, Petros B. Intestinal protozoan infections among HIV positive persons with and without Antiretroviral Treatment (ART) in selected ART centers in Adama and Dire-Dawa, Ethiopia. *Ethiop J Health Dev*. 2009; 23(Suppl 2): 133-139. doi: [10.4314/ejhd.v23i2.53230](https://doi.org/10.4314/ejhd.v23i2.53230)
31. Mehandru S, Poles MA, Tenner-Racz K, et al. Primary HIV-1 infection is associated with preferential depletion of CD4⁺ T lymphocytes from effector sites in the gastrointestinal tract. *J Exp Med*. 2004; 200:761-770. doi: [10.1084/jem.20041196](https://doi.org/10.1084/jem.20041196)
32. Brechley JM, Paiardini M, Knox KS, et al. Differential Th17 CD4 T-cell depletion in pathogenic and nonpathogenic lentiviral infections. *Blood*. 2008; 112(7): 2826-2835. doi: [10.1182/blood-2008-05-159301](https://doi.org/10.1182/blood-2008-05-159301)
33. Brechley JM, Price DA, Schacker TW, et al. Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nat Med*. 2006; 12: 1365-1371. doi: [10.1038/nm1511](https://doi.org/10.1038/nm1511)
34. Mendez OC, Szmulewicz G, Menghi C, Torres G, Gatta GC. Comparacion de indices de infestaciones por enteroparasitos entre poblaciones HIV- positivas y negativas [In Spanish]. *Medicina (Buenos Aires)*. 1994; 54: 307-310.
35. Cimerman S, Cimerman B, Lew DS. Enteric parasites and AIDS. *Sao Paulo Med J/Rev Paul Med*. 1999; 117(6): 266-273. doi: [10.1590/S1516-31801999000600007](https://doi.org/10.1590/S1516-31801999000600007)
36. Modjarrad K, Zulu I, Redden D, Lungowe N, Freedman D, Sten H. Prevalence and predictors of intestinal helminthes infections among human immunodeficiency virus type 1-infected adults in an urban African setting. *Am J Trop Med Hyg*. 2005; 73(Suppl 4): 777-782. doi: [10.4269/ajtmh.2005.73.777](https://doi.org/10.4269/ajtmh.2005.73.777)
37. Schmidt W, Wahnschaffe U, Schafer M. Rapid increase of mucosal CD4 T-cells followed by clearance of intestinal cryptosporidiosis in an AIDS patients receiving highly active anti-retroviral therapy. *Gastroenterology*. 2001; 120: 984-987. doi: [10.1053/gast.2001.22557](https://doi.org/10.1053/gast.2001.22557)
38. Gomez MA. Highly active anti-retroviral therapy and cryptosporidiosis. *Parassitologia*. 2004; 46(1-2): 95-99.
39. Assoumou A, Kone M, Penali LK, Coulibaly M, N'Draman AA. Cryptosporidiosis and HIV in Abidjan (Ivory Coast). *Bull Soc Pathol Exot*. 1993; 86(2): 85-86.
40. Downs JH. The gastrointestinal tract and HIV pathogenesis. *S Afr J Clin Nutr*. 2010; 23(Suppl 1): 65-68.
41. Munnink BBO, Canuti M, Deijs M, et al. Unexplained diarrhoea in HIV-1 infected individuals. *BMC Infect Dis*. 2014; 14: 22. doi: [10.1186/1471-2334-14-22](https://doi.org/10.1186/1471-2334-14-22)
42. Holtz LR, Finkbeiner SR, Kirkwood CD, Wang D. Identification of a novel picornavirus related to cosaviruses in a child with acute diarrhoea. *Virology*. 2008; 22; 5: 159. doi: [10.1186/1743-422X-5-159](https://doi.org/10.1186/1743-422X-5-159)
43. Kartalija M, Sande MA. Diarrhea and AIDS in the era of highly active antiretroviral therapy. *Clin Infect Dis*. 1999; 28: 701-77.
44. Call S, Mnkemüller K, Saag M, Wilcox CM. The changing etiology of diarrhea in patients with the acquired immune deficiency syndrome (AIDS). *Am J Gastroenterol*. 1998; 63: 1665.
45. Sharma HN, Chalise BS, Rai G, Adhikari N, Bastola A, Singh A. Prevalence of intestinal parasitic infections among people living with HIV/AIDS visiting a central hospital of Kathmandu Nepal. *Asian J Med Sc*. 2017; 8(5): doi: [10.3126/ajms.v8i5.17731](https://doi.org/10.3126/ajms.v8i5.17731)