

## Original Research

# Antihyperglycemic Mechanisms of *Allium sativum*, *Citrus sinensis* and *Persea americana* Extracts: Effects on Inhibition of Digestive Enzymes, Glucose Adsorption and Absorption on Yeast Cells and Psoas Muscles

Boris Gabin Kingue Azantsa, PhD<sup>1\*</sup>; Guy Roussel Takuissu, MSc<sup>1</sup>; Etienne Junior Tcheumeni, MSc<sup>1</sup>; Martin Fonkoua, PhD<sup>1</sup>; Edwige Ruth Kemadjou Dibacto, MSc<sup>1,2</sup>; Judith Laure Ngondi, PhD<sup>1</sup>; Julius Enyong Oben, PhD<sup>1</sup>

<sup>1</sup>Laboratory of Nutrition and Nutritional Biochemistry, Department of Biochemistry, Faculty of Sciences, University of Yaounde 1, Yaounde 812, Cameroon

<sup>2</sup>CRAN, Ministry of Scientific Research and Innovation, Yaounde, Cameroon

### \*Corresponding author

**Boris Gabin Kingue Azantsa, PhD**

Senior Lecturer, Department of Biochemistry, Faculty of Sciences, University of Yaounde 1, Yaounde 812, Cameroon; Tel: +237677920184;

E-mail: [borisazantsa@yahoo.fr](mailto:borisazantsa@yahoo.fr)

### Article information

Received: May 4<sup>th</sup>, 2019; Revised: May 16<sup>th</sup>, 2019; Accepted: May 20<sup>th</sup>, 2019; Published: June 3<sup>rd</sup>, 2019

### Cite this article

Azantsa BGK, Takuissu GR, Tcheumeni EJ, et al. Antihyperglycemic mechanisms of *Allium sativum*, *Citrus sinensis* and *Persea americana* Extracts: Effects on inhibition of digestive enzymes, glucose adsorption and absorption on yeast cells and psoas muscles. *Diabetes Res Open J.* 2019; 6(1): 1-9. doi: [10.17140/DROJ-6-143](https://doi.org/10.17140/DROJ-6-143)

## ABSTRACT

### Background

Mechanisms by which some plants with antihyperglycemic effects reduce postprandial hyperglycemia are not fully elucidated. This study was designed to investigate some action mechanisms of extracts from stem bark of *Citrus sinensis*, seeds of *Persea americana* and bulbs of *Allium sativum* including *in vitro* inhibition of  $\alpha$ -amylase and invertase; glucophagic capacity, absorption capacity on yeast cells and psoas tissues.

### Methods

Ethanollic (EE) and aqueous (AE) extracts were tested on  $\alpha$ -amylase and invertase activities. Glucose remaining in the medium was measured after direct interactions of: Glucose-extracts; Glucose-yeast-extracts and Glucose-psoas-extracts at different doses 0, 5, 7.5, 10 mg/ml.

### Results

All extracts inhibited invertase with IC<sub>50</sub> varying from 1.92 to 4.81 mg/ml for *Allium sativum* extracts.  $\alpha$ -amylase was inhibited by EE *C. sinensis* (IC<sub>50</sub> = 0.063 vs 2.73 mg/ml for arcabose) and not by EE of *A. sativum* and EE of *P. americana*. Glucophagic capacity of extracts varied significantly from 47.55 % of AE *P. americana* (5 mg/ml) to 100% with *C. sinensis* (5 mg/ml). All extracts stimulated glucose uptake ( $p < 0.05$ ) from 2.62 % AE *C. sinensis* (2.5 mg/ml) to 54.74% for EE of *Persea americana* (10 mg/ml). All extracts enhanced glucose uptake by psoas tissues increasing absorption capacity to up to 38.56 % with *A. sativum* (10.45% insulin,  $p < 0.05$ ).

### Conclusion

Cumulative actions of each plant extract on inhibition of carbohydrates' digestive enzymes, adsorption of glucose in intestine and blood, stimulation of glucose uptake and insulin action on yeast cells and psoas tissues, contribute to lower hyperglycemia and diabetes related complications. Therefore, extracts from the plants could be good candidates for diabetes therapy.

### Keywords

Mechanism; Absorption; Yeast cell; Psoas; Enzymes inhibition; Antihyperglycemic Plants.

## INTRODUCTION

Type 2 diabetes mellitus (T2DM) is a complex metabolic disorder of carbohydrates, fats and proteins resulting in an inability of insulin action and/or secretion.<sup>1</sup> T2DM is characterized by chronic hyperglycaemia with perturbations of glucose homeostasis due to absence of regulation of postprandial glycaemia. Postprandial glycaemia is at the onset of disturbances of glucose tolerance which appears earlier before high-levels of fasting blood glucose.<sup>2</sup> Several factors influence postprandial glycaemia: foods rich in carbohydrates, glucose digestion and absorption, insulin secretion as a result of rise of glucose levels, incretins actions and glucose intake by cells.<sup>3</sup> Postprandial glucose is also associated to protein glycosylation to the generation of reactive oxygen species which attack DNA and membrane lipids, to increase plasma lipid.<sup>4,5</sup> It is also a risk factor for cardiovascular diseases.<sup>6</sup>

Disturbances related to the glucose homeostasis as well as complications associated to dyslipidemia complexify management of diabetes, leading to alarming and increasing prevalence. In fact, 451 million individuals aged 18-99 years were reported to suffer from diabetes with more than 693 million of patients projected by 2045.<sup>7</sup> Such increasing prevalence is a public health concern and constitutes economic burden for governments. It is why efforts are made everyday to improve health of diabetic patients. Several oral drugs developed are used to reduce blood sugar. These include inhibitors of digestive enzymes, which blocked alpha glucosidases (amylase and invertase), inhibitors of intestinal absorption of glucose which inhibit membrane transporters: inhibitors of Dipeptidyl peptidase-4 (DPP4) which promote insulin secretion and molecules involved in the capture of peripheral glucose.<sup>8,9</sup> Thiazolidinediones, sulfonylureas, Metformin are oral antidiabetic drugs that play a prominent role in the treatment algorithm of T2DM.<sup>10,11</sup> Unfortunately, despite the efficacy of those drugs side effects like hypoglycemia, nausea, vomiting, headache and constipation are still reported.<sup>12-14</sup> Exploration of novel bioactive molecules from plants against diabetes has been intensified mainly with the availability of cell and tissue models like yeast cells and psoas muscles.<sup>14-16</sup>

Cameroonian pharmacopeia is very rich in plants with medicinal action including *Citrus sinensis*, *Persea americana* and *Allium sativum*.<sup>17</sup> They have been reported to have antihyperglycemic effects on rats in an oral glucose tolerance test.<sup>18,19</sup> However, few studies on their action mechanisms exist to justify their efficacy. This study was designed to evaluate and compare effects of ethanolic and aqueous extracts of *C. sinensis* stem bark, *P. americana* seeds and *A. sativum* bulbson digestive enzymes (invertase and alpha amylase); to determine their adsorptive/glucophagic capacity; to evaluate their capacity to enhance glucose uptake by yeast cells and psoas muscles models.

## MATERIALS AND METHODS

### Chemicals and Reagents

Krebs solution, insulin and acarbose (GLUCOR<sup>®</sup>) was purchased in local pharmacy. Baker's yeast, and *Chronolab* test kits for Glucose and amylase assays were purchased locally. Amylase Kit contained reagent R[(2-chloro-4-nitrophenyl- $\alpha$ -D-maltotriose) (CNP3;

2.25 mmol/L)+sodium Chlorure (NaCl, 350 mmol/L)+calcium acetate (CH<sub>3</sub>COO)<sub>2</sub>Ca, 6 mmol/L)+potassium Thiocyanate (KSCN, 900 mmol/L)]. Alpha amylase and invertase were purchased from Sigma Co., Louis, MO, USA.

### Plant Materials

Stem bark of *C. sinensis*, seeds of *P. americana* and bulb of *A. sativum* were used. *C. sinensis* was harvested in February 2017 in Yaounde, City capital of Cameroon. *A. sativum* and *P. americana* fruits were bought at Mokolo, a local market in Yaounde, Cameroon. *C. sinensis* L (voucher N° 25859/SRF), *P. americana* mill (voucher N° 31940 HNC), *A. sativum* (voucher N° 44810 HNC) were identified at the National Herbarium as belonging to the families of Rutaceae, Alliaceae and Lauraceae respectively. Stem bark were isolated from *C. sinensis* trees. Peels of *A. sativum* bulbs were removed and the bulbs were isolated. Seeds of *P. americana* fruits were removed. Each isolated part was cut into small pieces and dried in open air (till constant weight). The dried materials were ground finely to obtain powder, from which different extracts were prepared.

### Preparation of Ethanolic and Aqueous Extracts

One hundred grams (100 g) of each powder were placed in 800 mL of ethanol (95%) for maceration during 48 hours. The resulting supernatant was filtered using Whatman #1 filter paper (Whatman International Limited, Kent, England) through a funnel and concentrated to about 10% of the original volume by a rotavapor (BUCHI Rotavapor R-114, Switzerl) and at 45 °C, before complete drying in an oven at 50 °C. The same procedure was followed to prepare aqueous extract; ethanol being replaced by distilled water. After 24 h of maceration, the mixture was submitted to rotavapor. Extracts collected were stored in polyethylene bags to avoid moisture.

### Assessment of Antihyperglycemic Mechanisms

Three mechanisms were evaluated: inhibition of carbohydrates digestion through the  $\alpha$ -amylase and invertase assays, glucophagic effects through the glucose adsorption assay and stimulation of insulin-sensitivity through glucose uptake by yeast cells and muscles (psoas) assays.

### Invertase inhibition Assay

Invertase activity was evaluated after hydrolysis of sucrose to form glucose. Reaction mixture was made up with 290  $\mu$ L phosphate buffer, 40  $\mu$ L invertase and 250  $\mu$ L of sucrose was pre-incubated at 37 °C for 10 min. Then 40  $\mu$ L of plant extracts at different concentrations (0.5; 1; 1.5; 2; 2.5; 5 and 10 mg/ml) was added in all tubes except in the control containing 40  $\mu$ L of phosphate buffer. After incubation at 37 °C for 15 min, all the tubes were heated 100 °C in a water bath to stop the reaction. Glucose formed in each tube was measured as described by Trinder.<sup>20</sup> Absorbance (OD) was read at 505 nm. Another test tube containing no enzyme was

used as blank for calibration. Tests were done in triplicate. Results were expressed as inhibition percentage:

$$\text{Inhibition (\%)} = \frac{((\text{OD control} - \text{OD sample}))}{(\text{OD control}) \times 100}$$

#### **α-Amylase Inhibition Assay**

α-amylase activity was evaluated according to protocol described by Foo et al<sup>21</sup> with modifications. *Chronolab* test kit was used. α-amylase (AMS) hydrolysis 2-chloro-4-nitrophenyl-α-D-maltotriose (CNPG3) to yield 2-chloro-4-nitrophenol (CNP), 2-chloro-4-nitrophenyl-α-D-maltoside (CNPG2), maltotriose (G3) and glucose (G). The initial velocity measuring the formation of CNP was proportional to the enzyme catalytic concentration. Absorbance was read at 405 nm. In presence of extracts at different concentration, a reduction of absorbance was interpreted as inhibition of enzyme activity. Briefly, 75 μL of reagent Rwas diluted in 6 mL of distilled water. Then, 550 μL of reagent Rwas mixed with 100 μL of extracts at 1.25; 2.5; 5; 7.5 and 10 mg/mL. Another tube containing 100 μL of distilled water served as control. All the tubes were pre-incubated for 3 min at 37 °C, and 20 μL α-amylase at (0.31 mg/mL) was introduced in the reaction mixture. All the tubes were incubated again at 37 °C for 10 min and the reaction was stopped by placing them 5 min in a water bath at 100 °C. Another test tube containing no enzyme, but 40 μL phosphate buffer served as blank. Tests were done in triplicate. Results were expressed as inhibition percentage.

$$\text{Inhibition (\%)} = \frac{((\text{OD control} - \text{OD sample}))}{(\text{OD control}) \times 100}$$

#### **Glucose Adsorption Assay**

Glucose adsorption capacity of the extract was determined according to the method of Ou et al.<sup>22</sup> Briefly, 1 ml of extracts prepared at different concentrations (5; 10; 15 and 20 mg/ml) was added to 1 mL of glucose solution of increasing concentrations (12.5, 25, 37.5 and 50 mM), the mixture was stirred well, incubated in a shaker water bath at 37 °C for 1 hr, centrifuged at 4,000 g for 20 min. The glucose content in the supernatant was determined according to the method of Trinder.<sup>20</sup> Results were expressed as inhibition concentration. Glucose bound was calculated using the following formula:

$$\text{Glucose adsorption (\%)} = \frac{([\text{Glucose}]_{\text{initial}} - [\text{Glucose}]_{\text{final}})}{[\text{Glucose}]_{\text{initial}} \times 100}$$

#### **Glucose Uptake by Yeast Cell Assays**

Yeasts were prepared according to the method of Cirillo.<sup>23</sup> Commercial baker's yeast was washed by repeated centrifugation (4,000 g for 5 minutes) in distilled water until the supernatant became clear. Then 10% (v/v) suspension was prepared in distilled water. Extracts prepared at various concentrations (5; 10; 15 and 20 mg/ml) were added to 1 mL of glucose solution (25 mmol/L) and incubated together for 10 minutes at 37 °C. The reaction started by adding 100 μL of yeast suspension into the medium. The mixture was vortexed and further incubated at 37 °C for 60 minutes. After 60 minutes, the tubes were centrifuged (3,000 g for 5 minutes) and glucose in the supernatant was measured according to the method of Trinder.<sup>20</sup> A tube containing a mixture without extract was used as control. Glucose uptake by yeast cells was expressed in percentage and calculated using the following formula:

$$\text{Glucose uptake (\%)} = \frac{((\text{Absorbance Control} - \text{Absorbance sample}))}{(\text{Absorbance Control}) \times 100}$$

#### **Glucose Uptake in Rat Psoas Muscle Assay**

Glucose uptake in rat psoas muscle of the extract was determined according to the method of Al-Awadi et al<sup>24</sup> with modifications. Modifications included point measurements at 2h 30 min only. Psoas muscle was isolated from two anaesthetized adult rats and placed immediately in Krebs solution containing glucose (11.1 mm). Muscle tissue was cut into pieces of equal mass (0.25 g), and preincubated for 5 min in a CO<sub>2</sub> incubator. Triple sets including muscle tissue alone (control), muscle tissue with insulin (50 mU/L), muscle tissue with both insulin and extract (5, 7.5 and 10 mg/ml) were incubated for 2.5 hrs in CO<sub>2</sub> incubator under 95% O<sub>2</sub> and 5% CO<sub>2</sub> atmosphere. Changes in glucose concentrations were measured according to the method of Trinder.<sup>20</sup> The percentage of glucose uptake was calculated using the following formula:

$$\text{Glucose uptake (\%)} = \frac{((\text{Absorbance Control} - \text{Absorbance sample}))}{(\text{Absorbance Control}) \times 100}$$

#### **Statistical Analysis**


All experiments were performed in triplicate. Results obtained were expressed as mean ± SD from three distinct observations and as percentages. The software SPSS 20.0 (Chicago-Illinois Inc., IL, USA) was used for analyses. Chi square test were performed to compare percentages. One-way ANOVA followed by post-hoc Tukey's test was performed to compare variable amongst groups. Using Graph Pad Prism 6.0 (GraphPad Prism INC., CA, USA), the inhibition percentages *versus* logarithm of the extract concentrations were plotted and inhibitory concentration 50 (IC<sub>50</sub>) were

determined by regression. Analyses were done at 95% confidence interval.

## RESULTS


### Inhibition of Invertase Activity by Extracts

All the extracts inhibited invertase in a dose-dependent manner (Figure 1). Ethanolic extracts of *P. americana* and *C. sinensis* were the most efficient with values of  $IC_{50}$  of 2.69 and 2.35 mg/ml respectively, while aqueous extract of *A. sativum* was the most efficient of the aqueous extracts, with an  $IC_{50}$  of 1.92 mg/ml (Table 1).




### Inhibition of $\alpha$ -amylase activity by extracts

All extracts partially and weakly inhibited  $\alpha$ -amylase in a dose dependent manner except ethanolic extracts of *A. sativum* and aqueous extracts of *P. americana* that showed no inhibition (Figure 2A). Regression analysis revealed that aqueous extracts of *C. sinensis* was the most efficient and provided upto 91.94% inhibition of enzyme activity and an  $IC_{50}$ =4.96 mg/ml. All other extracts had inhibitions lower than 17.06%.



Under similar conditions acarbose used as reference drug gave an  $IC_{50}$  of 2.73% (Figure 2B and Table 1), higher than that of AE *C. sinensis*.



Invertase activity was reduced by extracts.  $IC_{50}$ s vary from 1.92 (AE *A. sativum*) to 4.81 (EE *A. sativum*). Inhibition of alpha-amylase activity by *C. sinensis* varies from  $IC_{50}$ =0.068 mg/ml (ethanolic extracts) to 58.09 mg/ml (aqueous extract) (Table 1). Aqueous extract of *P. americana* and ethanolic extracts of *A. sativum* showed null inhibition.


**Table 1.** Inhibitory Concentration 50 of Different Extracts ( $IC_{50}$ )

Extracts	P.americana		Citrus sinensis		Allium sativum		Acarbose
	Aqueous	Ethanolic	Aqueous	Ethanolic	Aqueous	Ethanolic	
Invertase	4.66	2.69	4.76	2.35	1.92	4.81	/
$\alpha$ -amylase	0	6.08	58.09	0.063	2.85	0	2.73

*IC50 : Inhibitory Concentration 50; Results expressed in mg/ml of extracts*


### Glucose Adsorption Capacity of Extracts

All extracts of different plants presented a minimum of 47.55% of glucose adsorption at the tested doses (Figure 3). Glucophagic activity observed was dose dependent. Aqueous extracts of *A. sativum* and *C. sinensis* were the most efficient with percentages varying from 75.87% to 82.68% ; 88.21% to 100% respectively, while ethanolic extract of *P. americana* was the most efficient with adsorption percentages of glucose from 86.71% to 96.35%. Aqueous extracts of *C. sinensis* presented the best glucophagic activity at low concentrations (<15 mg/ml).




### Glucose Uptake Capacity of Extracts on Yeast Cells Model

All extracts increased glucose capture by yeasts in a dose dependent manner (Figure 4). Ethanolic extracts were the most efficient with values varying from 19.14% to 45.46%, 18.24% to 54.74% and 27.56% to 37.19% respectively for *A. sativum*, *P. americana* and *C. sinensis*. The lowest efficacy was observed with AE of *C. sinensis* while EE of *P. americana* seeds presented the best absorption on yeast cells.



### Glucose Uptake Capacity in Rat Psoas Muscle

Using insulin as reference drug, psoas tissues were capable to uptake 10.45% of glucose. It appears that all extracts were capable to enhance glucose uptake at increasing concentrations ( $p < 0.05$  vs insulin) between 14.03% to 38.56% ; above insulin performance (Figure 5). Ethanolic extracts were the most efficient with uptake varying from 24.66% to 37.66%, 34.90% to 38.56% and 18.26% to 29.29% at increasing concentrations respectively for *P. americana*, *A. sativum* and *C. sinensis* compared to aqueous extracts. Furthermore, increment estimated as portion of glucose uptake by muscle tissue intrinsically attributable to extracts varied from 4.21% to 27.21%; 24.45% to 28.11% and 7.81% to 18.84% respectively for *P. americana*, *A. sativum* and *C. sinensis*. Bulbs of *A. sativum* presented the best lowering effects on blood glucose by mediation of glucose uptake by muscle tissues.



### DISCUSSION

Hyperglycemia is a major symptom of type 2 diabetes and a risk factor of cardiovascular diseases. However, many bioactive molecules derived from plants like polyphenols have proved their efficacy in the treatment of metabolic disorders.<sup>25</sup> Polyphenols are highly soluble in polar solvents like water and ethanol, reason why those two extracts were prepared from stem bark of *C. sinensis*, bulbs of *Allium sativum* and seeds of *Persea americana*. After consumption of foods, action of digestive enzymes  $\alpha$ -amylase and invertase in the gastrointestinal tract produces smaller molecules.<sup>15</sup> Monosaccharides resulting from digestion are absorbed into enterocytes to reach the blood stream, where they enter cells to supply energy and perform other biological functions.<sup>26</sup> Treatments administered in conditions of metabolic syndrome or diabetes target interaction with one or several of the abovementioned steps. In this study, the action of different plants have been reported on digestive enzymes. Starch and saccharose are the main polysaccharides found in meals.

An evaluation of invertase activity in the presence of both extract and sucrose revealed that all the extracts inhibited invertase (Figure 1) with  $IC_{50}$  varying from 1.92 (AE *A. sativum*) to 4.81 (EE *A. sativum*) (Table 1). Sucrase or invertase is a bifunction-

al enzyme also called sucrase-isomaltase which hydrolysis sucrose and isomaltose substrates.<sup>15</sup> If aqueous extract of *P. americana* and ethanolic extracts of *A. sativum* showed no inhibition (null) of alpha amylase at the tested concentrations (Figure 2A), EE of *Citrus sinensis* inhibited  $\alpha$ -amylase instead with an  $IC_{50}$ =0.063 mg/ml (*vs* 2.73 with acarbose) (Figure 2B and Table 1).  $\alpha$ -amylase hydrolyses the alpha bond linked polysaccharides of starch producing di- and mono-saccharides. Partial inhibition of the enzyme by *C. sinensis* extracts reduces its activity and prevents formation of monosaccharides.<sup>12-15</sup> Inhibition of amylase and invertase by extracts delays absorption of consumed carbohydrates and reduces plasma glucose.<sup>12</sup>

All the extracts strongly inhibited invertase (Figure 1), but only the AE of *P. americana* and EE extract failed to inhibit the two enzymes. Inhibitors of those enzymes surely possess in their structures functions similar to their substrates starch and sucrose although plant extracts do not possess such complex structures in their contents. Acarbose used as reference drug under the same experimental conditions provided an  $IC_{50}$  of 2.73 mg/mL, higher than *C. sinensis* extracts. Acarbose has a structure similar to oligosaccharides and therefore inhibits  $\alpha$ -glucosidases in a dose dependent manner, preventing the formation of glucose.<sup>27</sup> Flavonoids and alkaloids present in *Allium sativum* and americana extracts can justify their efficiency as well.<sup>18,19</sup> In fact, myricetin, quercetin, kaempferol are inhibitors of  $\alpha$ -amylase and  $\alpha$ -glucosidase while alkaloids like berberin are reported to be able to bind to  $\alpha$ -glucosidases and prevent sucrose fixation.<sup>28-31</sup> Their use to inhibit carbohydrates digestion is therefore beneficial in the management of diabetes because absorption of glucose is reduced and and postprandial hyperglycemia obviously.<sup>32,33</sup>

The residual glucose resulting from digestion can be trapped in the intestine limiting its absorption by adsorption, reducing its availability and diffusion into blood; reducing glucotoxicity as well. Adsorptive capacity through glucophagic activity of extracts revealed high percentages of glucose adsorption (Figure 3). Flavonoids and fibers content of extracts can justify such results. Insoluble fibers are capable of binding to glucose reducing its availability and therefore its intestinal absorption.<sup>34</sup> Also, they can reduce viscosity of secretions in intestinal lumen slowing or preventing diffusion. Absorbed glucose present in the blood can condense with hydroxyl groups of flavonoids to form complex molecules like glycosyl-flavonoids.<sup>35</sup> Decrease in glucose concentration contributes to reduce harmful effects of high plasma glucose on hemoglobin and other proteins, preventing glycation and complications such as retinopathy, neuropathy, and nephropathy observed in diabetic patients.<sup>36-38</sup>

Finally, the extracts were tested for their absorption capacity at the cellular levels. Glucose transport mechanism through cell membranes using yeast cells and psoas culture as models have been of great interest in the understanding of physiological processes of hypoglycemic activities of extracts.<sup>39,40</sup> Results revealed that all the extracts stimulated glucose entry in yeast cells (Figure 4). In fact, extracts are capable of stimulating glucose membrane transport-

ers, just like many oral antidiabetic drugs.<sup>39,40</sup> On yeast membrane, glucose is transported *via* specialised stereospecific groups of transporters called hexose transporters (HXT1 to HXT17) and specific molecules Snf3 and Rgt2, present on the membrane and involved in its capture. Binding of glucose to Snf3/Rgt2 located on yeast cell membrane permits phosphorylation of two co-receptors Mthl and Std 1 of Rgt1 (transcriptional repressor in charge of down regulation of the *HXT* gene) exposing Rgt1 to phosphorylation by a protein kinase A, causing glucose to diffusion across the membrane.<sup>41,42</sup>

Effectiveness of glucose absorption on yeast membrane mediated by extracts was confirmed by muscle cells tested *ex-vivo*, on extracts capacity to boost insulin action (Figure 5). Glucose absorption by cells is very important in glucose homeostasis. Upto 80% of glucose fixation depend on insulin action.<sup>43</sup> This study revealed that psoas in the presence of insulin produced 10.45% of glucose uptake. Insulin in the presence of each plant extract produced values higher than 10.45% in a dose dependent manner ( $p<0.05$ ). Similar observation was done with *A. aspera* extract.<sup>44</sup> Ability of extracts to boost insulin action can be attributed to polyphenols and alkaloids content of plants which act through stimulation of GLUT4 and/or AMPK activation or insulin secretory action.<sup>45</sup> Polyphenols stimulate glucose capture by muscle *via* activation of AMPK by mechanisms similar to thiazolidinediones.<sup>46,47</sup>

The enhanced action of extracts alongside with insulin on rat psoas muscles has been reported with maltitol.<sup>48</sup> Alkaloids like berberin stimulates direct glucose absorption mediated by AMPK without involvement of GLUT4.<sup>49</sup> Ability of extracts mainly ethanolic extracts to boost insulin action can be attributed to polyphenols known to stimulate binding of insulin to IRS receptors, to increase the number of insulin receptors or/and to increase insulin sensitivity.<sup>44-50</sup> Enhanced uptake of glucose by cells *via* joint action of insulin and extracts may stimulate glucose clearance from blood, through improved insulin resistance. Enhanced uptake of glucose also promote post absorptive utilisation in glycolysis for energy supply.<sup>44</sup> Absorption values above 10.45% and increments due to extracts varying from 4.21 to 28.11 % respectively for *P. americana* and *A. sativum* (Figure 5) cannot be intrinsically attributed to the stimulation of insulin action, because *in vitro*, extracts could adsorb part of glucose before insulin action, lowering its concentration in the medium. This study however demonstrated that, the cumulative actions of extracts on enzyme digestion, glucose adsorption and insulin action lead to reduced hyperglycemia, targeted in the first treatment outcome in type 2 diabetes. Reduced hyperglycemia also limits progression to complications.

## LIMITATIONS OF THE STUDY

Limitation of this study includes measurement of glucose concentration at 2 h 30 min only instead of different time intervals for better follow-up of absorption. Also, no reference drugs were used in invertase and glucose adsorption assays. Their use will have permitted better appreciate of the extracts' efficacy.

## CONCLUSION

Extracts from bulbs of *A. sativum*, seeds of *P. americana* and stem bark of *C. sinensis* exert their antihyperglycemic effects via inhibition of carbohydrate digestive enzymes, adsorption of glucose and stimulation of glucose absorption by cells. Stem bark extracts of *C. sinensis*, mainly ethanolic extracts are more effective on inhibition of digestion and glucose adsorption while bulbs of *A. sativum* stimulates glucose absorption in cells than other plant extracts. Bulbs of *A. sativum*, seeds of *P. americana* and stem bark of *C. sinensis* based on their efficiency and action mechanisms act like oral diabetic drugs. Therefore their use should be encouraged in the management of metabolic disorders.

## ACKNOWLEDGEMENTS

Not applicable.

## ETHICAL ISSUE

Not applicable.

## CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest.

## REFERENCES

- American Diabetes Association. Diagnostic and classification of diabetes mellitus. *Diabetes Care*. 2013; 36(Suppl 1): S67-S74. doi: [10.2337/dc13-S067](https://doi.org/10.2337/dc13-S067)
- Gerich J. Pathogenesis and management of postprandial hyperglycemia: role of incretin-based therapies. *Int J Gen Med*. 2013; 6: 877-895 doi: [10.2147/IJGM.S51665](https://doi.org/10.2147/IJGM.S51665)
- Agnaniet H, Mbot E, Keita O, Fehrentz J-A, et al. Antidiabetic potential of two medicinal plants used in Gabonese folk medicine. *BMC Complementary and Alternative Medicine*. 2016; 16: 71. doi: [10.1186/s12906-016-1052-x](https://doi.org/10.1186/s12906-016-1052-x)
- Shobana S, Sreerama YN, Malleshi NG. Composition and enzyme inhibitory properties of finger millet (*Eleusine coracana L.*) seed coat phenolics: Mode of inhibition of  $\alpha$ -glucosidase and pancreatic amylase. *Food Chemistry*. 2009; 115: 1268-1273. doi: [10.1016/j.foodchem.2009.01.042](https://doi.org/10.1016/j.foodchem.2009.01.042)
- Oboh G, Ademiluyi AO, Akindahunsi AA. Changes in polyphenols distribution and antioxidant activity during fermentation of some underutilized legumes. *Food Science and Technology International*. 2009; 15(1): 41-46. doi: [10.1177/1082013208101022](https://doi.org/10.1177/1082013208101022)
- Riccardi G, Rivellese AA, Giacco R. Role of glycemic index and glycemic load in the healthy state, in prediabetes, and in diabetes. *Am J Clin Nutr*. 2008; 87(1): 269S-274S. doi: [10.1093/ajcn/87.1.269S](https://doi.org/10.1093/ajcn/87.1.269S)
- Cho NH, Shaw JE, Karuranga S, et al. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res Clin Pract*. 2018; 138: 271-281. doi: [10.1016/j.diabres.2018.02.023](https://doi.org/10.1016/j.diabres.2018.02.023)
- Lordan S, Smyth TJ, Soler-Vila A, Stanton C, Ross RP. The  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibitory effects of Irish seaweed extracts. *Food Chem*. 2013; 141: 2170-2176. doi: [10.1016/j.foodchem.2013.04.123](https://doi.org/10.1016/j.foodchem.2013.04.123)
- American Diabetes Association. Standards of Medical Care in Diabetes-2017. *Diabetes Care*. 2017; 40(1): 1-80. [https://professional.diabetes.org/files/media/dc\\_40\\_s1\\_final.pdf](https://professional.diabetes.org/files/media/dc_40_s1_final.pdf)
- Inzucchi SE, Bergenstal RM, Buse JB. Management of hyperglycaemia in type 2 diabetes: A patient-centered approach: Position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care*. 2012; 55: 1577-1596.
- Song R. Mechanism of metformin: A tale of two sites. *Diabetes Care*. 2016; 39(2): 187-189. doi: [10.2337/dci15-0013](https://doi.org/10.2337/dci15-0013)
- Stein SA, Lamos EM, Davis NS. A review of the efficacy and safety of oral antidiabetic drugs. *Expert Opin Drug Saf*. 2013; 12(2): 153-175. doi: [10.1517/14740338.2013.752813](https://doi.org/10.1517/14740338.2013.752813)
- Sola D, Rossi L, Gian CS, et al. Sulfonylureas and their use in clinical practice. *Arch Med Sci*. 2015; 11(4): 840-848. doi: [10.5114/aoms.2015.53304](https://doi.org/10.5114/aoms.2015.53304)
- Chaudhury A, Chitharanjan D, Vijaya SD, et al. Clinical review of antidiabetic drugs: Implications for type 2 diabetes mellitus management. *Front Endocrinol (Lausanne)*. 2017; 8: 6. doi: [10.3389/fendo.2017.00006](https://doi.org/10.3389/fendo.2017.00006)
- Jijith US, Jayakumari S. Recent advances and methods for in-vitro evaluation of antidiabetic activity: A review. *J Res Ayurveda pharm*. 2017; 8(1): 81-87. doi: [10.7897/2277-4343.08117](https://doi.org/10.7897/2277-4343.08117)
- Maier A, Volker B, Boles E. Characterisation of glucose transport in *Saccharomyces cerevisiae* with plasma membrane vesicles (counter transport) and intact cells (initial uptake) with single Hxt1, Hxt2, Hxt3, Hxt4, Hxt6, Hxt7 or Gal2 transporters). *FEMS Yeast Res*. 2002; 2(4): 539-550. doi: [10.1111/j.1567-1364.2002.tb00121.x](https://doi.org/10.1111/j.1567-1364.2002.tb00121.x)
- Rauter A, Martins A, Lopes R, et al. Bioactivity studies and chemical profile of the antidiabetic plant *Genista tenera*. *J Ethnopharmacol*. 2009; 122(2): 384-393. doi: [10.1016/j.jep.2008.10.011](https://doi.org/10.1016/j.jep.2008.10.011)
- Azantsa KGB, Djuikoo NLI, Kuikoua W, Takuissu GN. Phytochemical screening and anti-diabetic evaluation of *C. sinensis* stem bark extracts. *International Journal of Biochemistry Research & Review*. 2017; 17(2): 1-13. doi: [10.9734/IJBCCR/2017/34051](https://doi.org/10.9734/IJBCCR/2017/34051)

19. Azantsa KGB, Oben J, Ngondi JL. Effects of polyherbal formulation of *Allium sativum* and *P. americana* seeds' extracts on postprandial hyperglycaemia and sucrose digestion in acute treatment of normoglycemic rats. *Biology and Medicine (Aligarh)*. 2018; 10: 432. doi: [10.4172/0974-8369.1000432](https://doi.org/10.4172/0974-8369.1000432)
20. Trinder P. Determination of blood glucose using 4-aminophenazone as oxygen acceptor. *J Clin Pathol*. 1969; 22(2): 246. doi: [10.1136/jcp.22.2.246-b](https://doi.org/10.1136/jcp.22.2.246-b)
21. Foo AY, Bais R. Amylase measurement with 2-chloro-4-nitrophenyl maltotriose as substrate. *Clin Chim Acta*. 1998; 27(2): 137-147.
22. Ou S, Kwok K, Li Y, Fu L. In vitro study of possible role of dietary fiber in lowering postprandial serum glucose. *J Agric Food Chem*. 2001; 49: 1026-1029. doi: [10.1021/jf000574n](https://doi.org/10.1021/jf000574n)
23. Cirillo V. Mechanism of glucose transport across the yeast cell membrane. *J Bacteriol*. 1962; 84: 485-491.
24. Al-Awadi F, Khattar M, Gumma K. On the mechanism of the hypoglycemic effect of a plant extract. *Diabetologia*. 1985; 28: 432-434. doi: [10.1007/BF00280886](https://doi.org/10.1007/BF00280886)
25. Oh Y, Jun HS. Role of bioactive food components in diabetes prevention: Effects on beta-cell function and preservation. *Nutr Metab Insights*. 2014; 7: 51-59. doi: [10.4137/NM.LS13589](https://doi.org/10.4137/NM.LS13589)
26. Meyer C, Woerle HJ, Dostou JM, Welle SL, Gerich JE. Abnormal renal, hepatic, and muscle glucose metabolism following glucose ingestion in type 2 diabetes. *Am J Physiol Endocrinol Metab*. 2004; 287(6): E1049-E1056. doi: [10.1152/ajpendo.00041.2004](https://doi.org/10.1152/ajpendo.00041.2004)
27. Scheen A. Place de l'acarbose dans le traitement du diabète sucré. *Diabetes and Metabolism*. 2018; 24(4): 1-4.
28. Mojica L, Meyer A, Berhow MA, de Mejia EG. Bean cultivars (*Phaseolus vulgaris* L.) have similar high antioxidant capacity, in vitro inhibition of  $\alpha$ -amylase and  $\alpha$ -glucosidase while diverse phenolic composition and concentration. *Food Res Inter*. 2015; 69: 38-48. doi: [10.1016/j.foodres.2014.12.007](https://doi.org/10.1016/j.foodres.2014.12.007)
29. Gaikwad S, Krishna G, Sandhya M. Phytochemicals for diabetes management. *Pharmaceutical Crops*. 2014; 5(1): 11-28.
30. Lanzotti V. The analysis of onion and garlic. *J Chromatogr A*. 2006; 1112: 3-22. doi: [10.1016/j.chroma.2005.12.016](https://doi.org/10.1016/j.chroma.2005.12.016)
31. Melgara B, Diasa M, Ciricc A, et al. Bioactive characterization of *P. americana* Mill. by-products: A rich source of inherent antioxidants. *Industrial Crops & Products*. 2018; 111: 212-218. doi: [10.1016/j.indcrop.2017.10.024](https://doi.org/10.1016/j.indcrop.2017.10.024)
32. Koyasu M, Ishii H, Watarai M, et al. Impact of acarbose on carotid intima-media thickness in patients with newly diagnosed impaired glucose tolerance or mild type 2 diabetes mellitus: A one-year, prospective, randomized, open-label, parallel-group study in Japanese adults with established coronary artery disease. *Clin Ther*. 2010; 32: 1610-1617. doi: [10.1016/j.clinthera.2010.07.015](https://doi.org/10.1016/j.clinthera.2010.07.015)
33. Kwon YI, Apostolidis E, Kim YC, Shetty K. Health benefits of traditional corn, beans and pumpkin: In vitro studies for hyperglycemia and hypertension management. *J Med Food*. 2007; 10: 266-275. doi: [10.1089/jmf.2006.234](https://doi.org/10.1089/jmf.2006.234)
34. Bhingre SD, Bhutkar MA, Randive DS, Wadkar GH, Hasabe TS. In-vitro hypoglycemic effects of unripe and ripe fruits of *Musa sapientum*. *The Brazilian Journal of Pharmaceutical Sciences*. 2017; 53(4): 1-4. doi: [10.1590/s2175-97902017000400159](https://doi.org/10.1590/s2175-97902017000400159)
35. Li S, Wu J, Chen L, Du H, et al. Biogenesis of C-glycosyl flavones and profiling of flavonoid glycosides in lotus (*Nelumbo nucifera*). *PLoS One*. 2014; 9(10): e108860. doi: [10.1371/journal.pone.0108860](https://doi.org/10.1371/journal.pone.0108860)
36. Ahmed N. Advanced glycation end products role in pathology of diabetic complications. *Diabetes Res Clin Pract*. 2005; 67(1): 3-21. doi: [10.1016/j.diabres.2004.09.004](https://doi.org/10.1016/j.diabres.2004.09.004)
37. Cade WT. Diabetes-related microvascular and macrovascular diseases in the physical therapy setting. *Phys Ther Phys Ther*. 2008; 88(11): 1322-1335. doi: [10.2522/ptj.20080008](https://doi.org/10.2522/ptj.20080008)
38. Haddadinezhad S, Ghazaleh N. Relation of fasting and postprandial and plasma glucose with hemoglobin A1c in diabetics. *Int J Diabetes Dev Ctries*. 2010; 30(1): 8-10. doi: [10.4103/0973-3930.60002](https://doi.org/10.4103/0973-3930.60002)
39. Boles E, Hollenberg CP. The molecular genetics of hexose transport in yeasts. *FEMS Microbiol Rev*. 1997; 21: 85-111. doi: [10.1111/j.1574-6976.1997.tb00346.x](https://doi.org/10.1111/j.1574-6976.1997.tb00346.x)
40. Roy A, Dement AD, Cho KH, Kim JH. Assessing glucose uptake through the yeast hexose transporter 1 (Hxt1). *PLoS One*. 2015; 10(3): e0121985. doi: [10.1371/journal.pone.0121985](https://doi.org/10.1371/journal.pone.0121985)
41. Gancedo J. The early steps of glucose signalling in yeast. *FEMS Microbiol Rev*. 2008; 32: 673-704. doi: [10.1111/j.1574-6976.2008.00117.x](https://doi.org/10.1111/j.1574-6976.2008.00117.x)
42. Revathi P, Thirumala P. In-vitro antidiabetic activity of ethanolic leaf extract of *Bruguiera cylindrica* L. - glucose uptake by yeast cells method. *IBBJ*. 2016; 2(4): 171-175.
43. Tripathy D, Chavez AO. Defects in insulin secretion and action in the pathogenesis of type 2 diabetes mellitus. *Curr Diab Rep*. 2010; 10(3): 184-191. doi: [10.1007/s11892-010-0115-5](https://doi.org/10.1007/s11892-010-0115-5)
44. Srinivasulu NP, Mallaiah G, Sudhakar B, Sasi BR, Sarala KD. Alpha amylase inhibitory activity and in vitro glucose uptake in psoas muscle and adipose tissue of male wistar rats of leaf



methanolic extract of *Achyranthes aspera*. *Journal of Pharmacognosy and Phytochemistry*. 2016; 5(3): 176-180.

45. Zhao FQ, Keating AF. Functional properties and genomics of glucose transporters. *Curr Genomics*. 2007; 8(2): 113-128. doi: [10.2174/138920207780368187](https://doi.org/10.2174/138920207780368187)

46. Breen DM, Sanli T, Giacca A, Tsiani E. Stimulation of muscle cell glucose uptake by resveratrol through sirtuins and AMPK. *Biochem Biophys Res Commun*. 2008; 374(1): 117-122. doi: [10.1016/j.bbrc.2008.06.104](https://doi.org/10.1016/j.bbrc.2008.06.104)

47. Zygumnt K, Faubert B, MacNeil J, Tsiani E. Naringenin, a citrus flavonoid, increases muscle cell glucose uptake via AMPK. *Biochem Biophys Res*. 2010; 398(2): 178-183. doi: [10.1016/j.bbrc.2010.06.104](https://doi.org/10.1016/j.bbrc.2010.06.104)

[bbrc.2010.06.048](https://doi.org/10.1016/j.bbrc.2010.06.048)

48. Chukwuma CI, Mohammed AI, Shahidul I. Maltitol increases-insulin mediated muscle glucose uptake *ex vivo* but not in normal and type 2 diabetic rats. *International Journal of Food Sciences and Nutrition*, 2016 ; 73-81. doi: [10.1080/09637486.2016.1216527](https://doi.org/10.1080/09637486.2016.1216527)

49. Yin J, Gao Z, Liu D, Liu Z, Ye J. Berberine improves glucose metabolism through induction of glycolysis. *Am J Physiol Endocrinol Metab*. 2008; 294(1): E148-E156. doi: [10.1152/ajpendo.00211.2007](https://doi.org/10.1152/ajpendo.00211.2007)

50. Gupta R, Kesari A, Watal G, et al. Hypoglycaemic and antidiabetic effects of aqueous extracts of leaves of *Annona squamosa* in experimental animal. *Current Science*. 2005; 88(8): 1244-1254.