

Research

*Corresponding author Srimathi Kannan, PhD

Address (where this research was funded and completed) Department of Nutrition School of Public Health and Health Sciences University of Massachusetts Amherst, USA Tel. 4132306847 Fax: 4132306847 E-mail: srimathik@gmail.com

Volume 4 : Issue 1 Article Ref. #: 10000R0J4128

Article History

Received: October 20th, 2016 Accepted: November 21st, 2016 Published: November 22nd, 2016

Citation

Kannan S, Visintainer P, Ganguri HB, Conroy R, Gudala M, Wittcopp C. Body mass index is a strong predictor of vitamin D deficiency in multiethnic obese children. *Obes Res Open J.* 2016; 4(1): 11-18. doi: 10.17140/OROJ-4-128

Copyright

©2016 Kannan S. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

= Open Journal 👌 =

Body Mass Index is a Strong Predictor of Vitamin D Deficiency in Multiethnic Obese Children

Srimathi Kannan, PhD¹; Paul Visintainer, PhD²; Harish B. Ganguri, MS³; Rushika Conroy, MD⁴; Meghana Gudala, MBBS⁵; Chrystal Wittcopp, MD⁵

¹Department of Nutrition, School of Public Health and Health Sciences, University of Massachusetts, Amherst, USA (where this work was initiated and completed) ²Department of Medicine, UMMS-Baystate, Springfield, Massachusetts, USA ³Department of Computer Science, Southern Illinois University, Carbondale, Illinois, USA; Kronsys Inc, Raleigh, North Carolina, USA

⁴Department of Pediatrics, Tufts University School of Medicine, Boston, Massachusetts, USA ⁵Department of Pediatrics, Baystate Children's Hospital, Chestnut St., Springfield, MA, USA

ABSTRACT

Background: Vitamin D deficiency is highly prevalent among obese children in the United States.

Objective: The main objective of this study is to determine predictors of vitamin D deficiency in obese children.

Methods: Children aged 5-14 years with body mass index z-score (BMIZ)>95% were enrolled. Data included height, weight, ethnicity, season, dietary intake, serum 25-hydroxy vitamin D test: (25(OH)D), parathyroid hormone, calcium, insulin, glucose, high sensitivity C-reactive protein (CRP), lipids, aspartate aminotransferase (AST), alanine transaminase (ALT) and waist circumference. BMIZ calculation in children was categorized into \geq 2.5 (high) and <2.5 (low). Serum 25(OH)D levels were dichotomized as <20 ng/mL (deficient) or \geq 20 ng/mL (sufficient). **Results:** Ninety-one children completed the study. Mean BMIZ was 31.3±6.3 kg/m². Mean 25(OH)D level was 21.2±7.6 ng/mL. A base logistic regression model showed odds of a deficient 25(OH)D was more than 3-times higher in children with a high BMIZ (OR: 3.35, 95% CI: 1.07-10.43). African-American children were more likely than Caucasian children to have low 25(OH)D (OR: 6.68, 95% CI: 1.31-36.13). Children enrolled in seasons other than winter were less likely to have a low 25(OH)D. Adjustment for nutrients and serum cardiovascular disease risk markers did not significantly alter the relationship between low 25(OH)D and BMIZ.

Conclusions: The risk for vitamin D deficiency increases in children with severe obesity (BMIZ≥2.5 SD) and does not significantly change by season, ethnicity, nutrient intake, or cardiovascular disease risk parameters.

KEYWORDS: Vitamin D deficiency; Obesity; Multiethnic children; BMIZ; Parathyroid hormone (PTH).

ABBREVIATIONS: BMIZ: Body Mass Index Z-Score ; PTH: Parathyroid hormone; AST: Aspartate aminotransferase; ALT: Alanine aminotransferase; VDBP: Vitamin D Binding Protein; CDC: Centers for Disease Control and Prevention; FFQ: Food Frequency Questionnaire.

INTRODUCTION

Vitamin D deficiency is highly prevalent in the pediatric population.¹ Evidence suggests that obese children are more prone to vitamin D deficiency than non-obese children,² with the degree of deficiency directly related to the degree of adiposity. To our knowledge, however, there are no recent data available on the predictors of low 25(OH)D levels among multiethnic obese

Open Journal 👌

ISSN 2377-8385

children. In adults, adiposity has long been known to be inversely related to circulating 25(OH)D,^{2,3} so it is not surprising that this association is being recognized in obese children, with studies showing that as many as 75% of obese children are vitamin D deficient.¹

Vitamin D deficiency is defined by the Institute of Medicine (IOM) as a serum 25-hydroxy vitamin D <20 ng/mL (<50 nmol/L).⁴ The etiology of vitamin D deficiency is multi-factorial.⁵ Proposed risk factors include race/ethnicity,⁶ lack of sufficient sunlight exposure,⁷ insufficient activity,⁸ decreased consumption of vitamin D-containing foods such as fish and for-tified dairy products.⁹ malabsorptive states,¹⁰ increased vitamin D degradation and increased body weight.⁶ Other predictors of low vitamin D status in healthy children include age,¹¹ gender¹¹ and socio-economic status.¹

In children, vitamin D is essential for skeletal health.¹² Deficiency in children has also been linked to increased risk of allergies and asthma,¹³ and cardiovascular risk markers.¹⁴ In obese children, low levels of vitamin D may play a role in the pathophysiology of impaired glucose tolerance.¹⁵ It is therefore essential to know what factors predict vitamin D deficiency in children.

The objective of the study is to examine the association between vitamin D deficiency and BMI z-score (BMIZ) in obese children. We hypothesize that predictors of deficient serum 25(OH)D in obese children will be low dairy intake, ethnicity, season and higher BMIZ.

MATERIALS AND METHODS

Participants

Children aged 5-14 years with a BMIZ≥95% for age and gender followed at a pediatric weight management clinic were enrolled in the study from March 2011 to July 2012. Exclusion criteria were: use of vitamin D supplementation; history of hyperparathyroidism, rickets, type 1 or type 2 diabetes (T2D), renal failure, nephrolithiasis or new-on set hypothyroidism; and, use of prednisone, phenytoin, thiazides, statins or orlistat. Informed consent and/or assent were obtained for all participants and their caregivers prior to participation and study protocols were an Institutional Review Board (IRB) approved.

Data Collection

Fasting blood was drawn for serum 25(OH)D, parathyroid hormone (PTH), calcium, insulin, glucose, HbA1c, high sensitivity C-reactive protein (hs-CRP), lipids, aspartate aminotransferase (AST) and alanine aminotransferase (ALT). Anthropometric data were collected: height to the nearest 0.1 cm on a standard wall stadiometer, weight to the nearest 0.1 kg on an electronic scale (Scale-tronic model 6002; Wheaton, IL, USA) with children in light clothing, waist circumference (WC) measured at the level of the iliac crest, and blood pressure. BMIZ was calculated as weight in kg divided by the square of the height in meters. BMIZ percentiles were determined by using age- and gender-specific Centers for Disease Control and Prevention (CDC) growth charts.¹⁶ Child age, gender, ethnicity, past medical history and family history of vitamin D deficiency, T2D and early chemical vapor deposition (CVD) were reported by parents. Season of serum collection was recorded as summer, fall, winter and spring.

Biomarker Assays

Serum 25(OH)D concentrations were determined after extraction with acetonitrile, using a standard radioimmunoassay (RIA) developed by DiaSorin, Inc. (Stillwater, MN, USA). Serum 25(OH)D was assayed using an equilibrium RIA with 125I. Lower limit of detection was 1.5 ng/mL (DiaSorin, Inc.). Serum lipids and blood glucose were measured by enzymatic methods with Cobas 8000 (Roche Diagnostics, Basel, Switzerland). hs-CRP was measured with a particle enhanced immunoturbidimetric method (Roche Diagnostics, Basel, Switzerland). PTH and fasting insulin were measured using Electro chemiluminescence immunoassay (Roche Diagnostics, Basel, Switzerland). Calcium levels were measured using Cobas method according to Schwarzenbach¹⁷ Roche Diagnostics is a diagnostic division of Hoffmann-La Roche which manufactures equipment and reagents for research and medical diagnostic (Roche Diagnostics, Basel, Switzerland).¹⁷ Hemoglobin A1c Test Normal and high chart and ranges (HbA1C) was measured using Cobas Turbimetric (Roche Diagnostics, Basel, Switzerland). AST and ALT were measured using NADH oxidation (Roche Diagnostics, Basel, Switzerland).

Dietary Assessment

Dietary intakes were assessed using the 8-page Block Kids 2004 FFQ (Nutrition Quest, Berkeley, CA, USA), incorporating approximately seventy food and beverage items. The Food Frequency Questionnaire (FFQ) has been validated for multiethnic school-age children⁹ residing in the same geographic region as the current study. It has also been specifically validated for estimating beverage, Ca and vitamin D intakes in children when compared with 3-day food diaries.¹⁸ Portion size pictures were used to increase the accuracy of their estimates.

The FFQs were administered by trained staff. Proxy respondents reported for 5-year-old children and proxy-assisted interviews were conducted with children 6-14 years of age. FFQs were reviewed for completion and processed through the Block Dietary Data Systems (Berkeley, CA, USA),¹⁹ and quantified by Nutrition Quest. Nutrient values were calculated for each item and further summarized into daily intakes of energy and nutrients using an algorithm from Nutrition Quest. Additional vitamin D contribution from foods and beverages was requested separately from Nutrition Quest. Nutrient values were determined based on updated values for fortified foods from the USDA 1998 nutrient database²⁰ and adjusted for bioavailability through nutrient-equivalent estimates when relevant.²¹

Open Journal 👌

ISSN 2377-8385

Statistical Analysis

All analysis were conducted using Stata (version 14.0, Stata-Corp, College Station, TX, USA). Descriptive statistics, univariate analysis, and Fisher's exact test were employed. If continuous variables were not normally distributed, the Wilcoxon rank-sum test was employed.

Serum 25(OH)D levels were dichotomized into <20 ng/ mL (deficient) or \geq 20 ng/mL (sufficient). BMIZ was dichotomized into \geq 2.5 (high) vs. <2.5 (low). Nutrient intakes were energy-adjusted using the residual method.²² Daily milk intake was determined with "number of milk-cup equivalents" (servings), and dairy, using "daily servings of milk, yogurt and cheese" and total dairy intake was assessed as total dairy intake in milk equivalent servings. The dietary variables were correlated with the cardiovascular markers and waist circumference and then compared by BMIZz category, using *t*-tests and rank-sum tests.

The month of enrollment was used as a coarse indicator of average sun exposure. Months were combined into seasons for analysis: winter (December, January, February), spring (March, April, May), summer (June, July, August), and fall (September, October, November).

We computed an age- and gender- standardized WC score based on Sabo, 2012.²³ For purposes of analysis, we categorized the sample into two groups above and below the median for the sample. So, the sample was split at the sample median standardized WC values, which were 3.4.

A base regression model evaluated BMIZ, adjusting for age, gender, ethnicity and season. Significance testing of their overall contribution to the model was assessed using a likelihood ratio test. Significance testing relative to the reference group, as http://dx.doi.org/10.17140/OROJ-4-128

for all other variables in the model, was conducted using a standard Wald test. In light of the restricted sample size, to account for nutritional factors, energy-adjusted nutrient variables were added singly to the base regression model. Similarly, cardiovascular disease (CVD) risk markers were evaluated as confounders by introducing each marker individually into the base model. Adjusted risk estimates are reported as odds ratios with 95% confidence intervals (CI). Significance testing was conducted at a critical level of 5%.

RESULTS AND DISCUSSION

Demographics

Of 115 children enrolled, 24 were excluded for incomplete dietary data, implausible energy intake <500 or >5000 kcal/d,²⁴ or for missing serum markers. Ninety-one children [mean age 10.9±2.6 (SD) years] had complete dietary data. The majority were female (52.7%). Of the total, 26.4% were Caucasian, 51.7% were Hispanic, and 22% were African American. The majority of participants were enrolled in the spring or summer (71.5%) (Table 1). Clinical characteristics are shown in Table 2. The mean BMIZ was 31.6±6.5 kg/m², BMIZ was 2.40±0.37. The mean standardized waist circumference was 3.7 ± 1.5 (n=90). For subsequent analysis, BMIZ was dichotomized at 2.5 (about the 99.5%): 37.4% (n=34)≥2.5, and 62.6% (n=57)<2.5.

25(OH)D and PTH Levels

The mean 25(OH)D was 21.2 \pm 7.6 ng/mL, with 47.3% (n=43) having levels <20 ng/mL. Significant associations with 25(OH) D levels were seen with age, season and ethnicity (p<0.01) (Table 2). The prevalence of vitamin D deficiency was higher among ethnic minority and older children.

Sample Characteristic	Overall (n=91)	VDL≥20 (n=48)	VDL<20 (n=43)	p
	mean (SD)	mean (SD)	mean (SD)	
Age	10.9 (2.6)	10.1 (2.5)	11.8 (2.5)	<0.001
	n (%)	n (%)	n (%)	
Male	47.3 (43)	54.2 (26)	39.5 (17)	0.21
Ethnicity				0.12
Caucasian	26.4 (24)	33.3 (16)	18.6 (8)	
African-American	22.0 (20)	14.6 (7)	30.2 (13)	
Hispanic	51.7 (47)	52.1 (25)	51.2 (22)	
Season				0.009
Winter	15.4 (14)	4.2 (2)	27.9 (12)	
Spring	44.0 (40)	45.8 (22)	41.9 (18)	
Summer	27.5 (25)	31.3(15)	23.3 (10)	
Fall	13.2 (12)	18.8 (9)	7.0 (3)	

*among children with total calories ≥500

Table 1: Characteristics of study sample (n=91), overall and by vitamin D Level.

Open Journal 👌

ISSN 2377-8385

Sample	Overall (n=91)	25(OH)D≥20 (n=48)	25(OHD<20 (n=43)	Р
Characteristic	mean (SD)	mean (SD)	mean (SD)	
SBP	114.8 (10.1)	111.9 (8.8)	118.1 (10.5)	0.003
DBP	66.2 (7.0)	64.7 (6.2)	67.9 (7.5)	0.03
	n (%)	n (%)	n (%)	
BMIZ≥2.5	37.4 (34)	31.3 (15)	44.2 (19)	0.28
WC≥3.4ª	51.1 (46)	47.9 (23)	54.8 (23)	0.53

a90 children total

SBP=Systolic Blood Pressure; DBP=Diastolic blood pressure *Among children with total calories≥500; BMIZ=Body mass index by age percentile z-score; WC=waist circumference

Table 2: Clinical Characteristics of study sample (n=91), overall and by vitamin D level.

Vitamin D levels were lower in the winter season of serum collection and in children with higher BMIZ. The effect of BMIZ on serum 25(OH)D may be explained by the fact that persons with high BMIZ may have excess body fat, potentially resulting in increased sequestration of vitamin D and low bioavailability and thus, low serum 25(OH)D levels.³ Other evidence suggests that excess body fat may disrupt hormonal pathways important for skeletal health. For example, leptin that binds to osteoblasts appears to activate a pathway that inhibits renal synthesis of the active form of vitamin D.²⁵ Leptin levels are markedly elevated in obese individuals thereby decreasing 25(OH) D levels. vitamin D deficiency is associated with insulin resistance,²⁶ which, in turn, is inversely correlated with vitamin D binding protein (VDBP).²⁶ Protein bound vitamin D is inactive, whereas non-protein bound vitamin D is the active form. We speculate that the total 25(OH)D we observed may have been counter balanced by a parallel insulin resistance-driven reduction of VDBP.

This is corroborated by potentially higher HOMA values in children with 25(OH)D < 20 ng/mL vs. those with non-deficient levels.

There were no significant differences in parathyroid hormone (PTH) levels between those with 25(OH)D<20 ng/mL vs. those ≥ 20 ng/mL (Table 3). Our sub-analysis of the relationship between serum 25(OH)D and PTH revealed no association. Maximal suppression of serum parathyroid hormone: PTH in adults has been reported over a wide range of serum 25(OH) D from 20-110 nmol/L.26 Giudice27 found comparable levels of PTH despite different levels of total 25(OH)D in obese and nonobese children. In a recent study, however, Amini²⁸ showed that the level of serum 25(OH)D below which PTH began to rise was lower in overweight/obese compared to normal weight children (12.4 vs. 17.0 ng/mL). This might be a consequence of similar ionized calcium levels despite lower 25(OH)D.²⁹ We speculate that the PTH is responding to total body 25(OH)D, not just circulating serum levels. Other factors that can influence PTH include the assay utilized, diurnal variation, age, and dietary intake of calcium and phosphorus.³⁰ Our study participants were all adolescents and children, and normal ranges for PTH are not dissimilar in these age groups. Our assay was specific enough to consider only intact PTH molecules minimizing falsely elevated levels. We factored in dietary intake of calcium in our analysis

but did not consider specifically phosphorous intake.

Dietary Intakes

Children in the vitamin D deficient group had lower mean daily intakes of vitamin D, calcium, n-3 fatty acids (134.8 vs.173.8 IU/d, p<0.03; 657.9 vs. 763.8 mg/d, p<0.005; 0.8 vs. 1.0 mg/d, p<0.005), and total sugars (80.5+/-36.4 g/d vs. 110.0+/-58.7 g/d, p<0.005, Table 3). The median number of servings from the meat group was lower in children with 25(OH)D<20 ng/mL compared to those with 25(OH)D≥20 ng/mL (median 1.1 vs. 1.4 servings; inter quartile range 1.0-1.7, p<0.05).

Other Clinical Measures

As shown in Table 1, children with low 25(OH)D had higher fasting insulin (13.1 vs. 19.1 IU/ml, p<0.01), systolic (111.9 vs. 118.1 mmHg, p<0.04) and diastolic blood pressure (64.7 vs. 67.9 mmHg, p<0.03) and lower serum calcium (9.8 vs. 10.0 mg/dL, p<0.005). AST was higher in those with 25(OH)D≥20 ng/mL when compared to those with <20 ng/mL (24.6+/- 5.0 vs. 21.1+/- 5.1 U/L; p<0.002).

Multiple factors impacting risk for vitamin d deficiency

The base logistic regression model examining vitamin D deficiency included BMIZ (<2.5 vs. ≥2.5), gender, ethnicity, age, and season is shown in Table 4. Odds of vitamin D deficiency were more than three times higher in children with BMIZ >2.5 compared to those <2.5 when adjusting for the other covariates (OR: 3.35, 95% CI: 1.07-10.43; p= 0.04). The odds of Vitamin D deficiency increased by 47% for each year of age (OR: 1.47, 95% CI: 1.17-1.83; p=0.001); African-American children were more than six times as likely as Caucasian children to have low 25(OH)D (OR: 6.68, 95% CI: 1.31-36.13; p=0.02); and, relative to winter, children entering the study during the other seasons were less likely to have lower 25(OH)D levels. Although males were less likely to have low 25(OH)D than females, this difference did not reach statistical significance (or: 0.44, 95% CI: 0.15-1.26; p=0.13). When the base model was further adjusted for nutrient residuals, risk of Vitamin D Deficiency for those with a BMIZ 2.5 is unchanged and remained stable regardless of the nutrient added to the base model (Figure 1). The adjust-



http://dx.doi.org/10.17140/OROJ-4-128

Open Journal 👌

ISSN 2377-8385

Cardiovascular	Overall (n=91)	VDL≥20 (n=48)	VDL<20 (n=43) P	
Weasure	mean (SD)	mean (SD)	mean (SD)	
Fasting glucose ^a	89.7 (7.9)	89.7 (8.9)	89.8 (6.6)	0.94
Fasting insulin	15.9 (13.3)	13.1 (9.6)	19.1 (15.9)	0.04
PTH⁴	31.3 (11.2)	29.4 (8.9)	33.6 (13.2)	0.09
Ca⁵	9.9 (0.4)	10.0 (0.3)	9.8 (0.4)	0.005
ASTª	23.0 (5.3)	24.6 (5.0)	21.1 (5.1)	0.002
ALT ^a	20.9 (8.7)	21.2 (7.3)	20.6 (10.1)	0.77
sCRP⁰	3.7 (4.0)	4.1 (4.3)	3.3 (3.6)	0.34
HOMA ^a	3.5 (3.1)	3.0 (2.5)	4.2 (3.7)	0.08
Nutritional	Overall (n=91)	VDL≥20 (n=48)	VDL<20 (n=43)	Р
weasure	mean (SD)	mean (SD)	mean (SD)	
Nutrient ¹				
Vitamin D	155.4 (85.6)	173.8 (90.2)	134.8 (75.9)	0.03
Calcium	713.8 (306.5)	763.8 (320.1)	657.9 (284.0)	0.10
Magnesium	167.9 (76.6)	183.9 (82.0)	150.0 (66.6)	0.03
Phosphorus	884.7 (360.4)	951.8 (381.2)	809.7 (323.9)	0.06
Omega-3 FA	0.9 (0.4)	1.0 (0.5)	0.8 (0.4)	0.05
MUFAs	17.4 (7.6)	18.5 (7.5)	16.2 (7.6)	0.14
PUFAs	9.5 (4.5)	10.2 (4.5)	8.6 (4.5)	0.10
Saturated fats	16.7 (6.8)	17.8 (6.9)	15.4 (6.6)	0.10
Trans-fats	47.4 (20.0)	50.6 (19.9)	43.7 (19.8)	0.10
Cholesterol	165.0 (87.6)	175.7 (74.9)	153.0 (99.4)	0.23
Total sugars	96.0 (51.3)	110.0 (58.7)	80.5 (36.4)	0.005
Servings ²	Median (IQR)	Median (IQR)	Median (IQR)	
Meat	1.2 (0.8, 1.7)	1.4 (1.0, 1.7)	1.1 (0.7, 1.7)	0.05
Eggs	0.2 (0.1, 0.4)	0.2 (0.1, 0.4)	0.1 (0, 0.4)	0.06
Dairy	1.7 (0.9, 2.3)	1.8 (1.0, 2.4)	1.5 (0.9, 2.0)	0.12
Vegetables	0.8 (0.4, 1.8)	0.8 (0.5, 1.8)	0.8 (0.4, 1.8)	0.64
Grains	3.2 (2.3, 4.6)	3.8 (2.3, 4.9)	2.8 (2.1, 4.2)	0.08
Milk	1.2 (0.6, 1.9)	1.3 (0.6, 2.1)	1.0 (0.6, 1.5)	0.08
Total Dairy	1.6 (0.9, 2.3)	1.7 (1.0, 2.4)	1.4 (0.9, 2.0)	0.11
Total Grains	3.5 (2.5, 4.8)	3.8 (2.6, 4.9)	3.2 (2.5, 4.1)	0.12
Whole Grain	0.3 (0.2, 0.5)	0.3 (0.2, 0.5)	0.3 (0.1, 0.6)	0.56
Fruit Juice	0.3 (0.1, 0.8)	0.3 (0.1, 1.0)	0.3 (0, 0.5)	0.05
Fish (High)	0.0 (0.0, 0.1)	0.0 (0, 0.1)	0.0 (0, 0.1)	0.55
Fish (Low)	0.0 (0.0, 0.2)	0.0 (0.0, 0.1)	0.0 (0, 0.2)	0.81

Note: test; analysis using t-tests, with unequal variances; ²analysis using the Wilcoxon rank-sum

test, *n=90; *n=89; *n=87; *n=86 TC= total cholesterol; TG =triglycerides; HDL= high density lipoprotein; LDL=low density lipopro-tein; A1C=glycosylated hemoglobin; PTH=parathyroid hormone; Ca=calcium; AST= aspartate aminotransferase; ALT=alanine aminotransferase; hsCRP=high sensitivity C-reactive protein; HOMA=homeostatic model assessment; SBP=systolic blood pressure; DBP=diastolic blood pressure. *n-3 = comega 3 fatty acids **MUFA=monounsaturated fatty acids *PUFA=polyunsaturated fatty acids

Fish (High) Fish (Low)

Table 3: Laboratory cardiovascular measures, overall and by vitamin D level (n=91).

http://dx.doi.org/10.17140/OROJ-4-128

penventio

PUBLISHERS

Characteristic	Odds Ratio	95% CI	<i>p</i> -value
BMI z-score			
< 2.5 (ref)	1.0		
≥2.5	3.35	(1.07-10.43)	0.04
Gender			
Female (ref)	1.0		
Male	0.44	(0.15-1.26)	0.13
Ethnicity			0.05ª
Caucasian (ref)	1.0		
African-American	6.68	(1.31-36.13)	0.02
Hispanic	2.91	(0.77-10.96)	0.11
Age (years)	1.47	(1.17-1.83)	0.001
Season			0.02ª
Winter (ref)	1.0		
Spring	0.12	(0.02-0.75)	0.02
Summer	0.08	(0.01-0.55)	0.01
Fall	0.07	(0.01-0.57)	0.01

Note: «likelihood ratio test of variable's overall significance in model

Table 4: Logistic regression model of adjusted risk of low vitamin D level (<20 ug/mL) (n=91).

Model	ORs (95% CI)	р
Base models		
Unadjusted	1.74 (0.74, 4.10)	0.20
Model w/o nutrients	3.35 (1.07, 10.43)	0.03
Adjusting for nutrients		
Vit D	3.50 (1.10, 11.17)	0.03
Calc	3.35 (1.07, 10.45)	0.03
Magn	3.46 (1.10, 10.93)	0.03
Phos	3.35 (1.07, 10.47)	0.03
Omega 3	3.71 (1.14, 12.01)	0.02
Mono Fat	3.40 (1.08, 10.69)	0.03
Poly Fat	3.42 (1.09, 10.74)	0.03
Sat Fat	3.27 (1.05, 10.21)	0.04
Total Fat	3.29 (1.05, 10.24)	0.04
Chol	3.34 (1.07, 10.42)	0.03
Total Sugar	3.01 (0.96, 9.48)	0.06
Adjusting for servings		
Meat	3.33 (1.07, 10.37)	0.03
Just eggs	3.35 (1.07, 10.48)	0.03
Dairy	3.44 (1.09, 10.83)	0.03
Vegetables	3.48 (1.10, 11.03)	0.03
Grains	3.50 (1.10, 11.14)	0.03
Milk	3.38 (1.07, 10.61)	0.03
Total dairy	3.42 (1.09, 10.75)	0.03
Total grains	3.60 (1.12, 11.58)	0.03
Whole grains	3.44 (1.10, 10.72)	0.03
Fruit juice	2.96 (0.92, 9.58)	0.07
Fish (Hi)	3.31 (1.06, 10.35)	0.04
Fish (Lo)	3.28 (1.05, 10.29)	0.04

Figure 1: Logistic regression analysis of association between Vitamin D and BMI z-score category, controlling for children's Intakes of nutrients and food groups (n=91).

Open Journal 👌

ISSN 2377-8385

Model	ORs (95% CI)	p
Base models			
Unadjusted	= 1.74 (i	0.74, 4.10)	0.20
Model w/o cardio factors	- 3.35 (1.07, 10.43)	0.03
LR models w/ cardiovascular factors			
Tchol	3.30 (1.06, 10.31)	0.04
Trig	3.59 (1.08, 11.92)	0.03
HDL	3.85 (1.17, 12.62)	0.02
LDL	3.20 (1.02, 10.03)	0.04
A1c	3.35 (1.06, 10.60)	0.04
f-Glucose	3.27 (1.04, 10.28)	0.04
f-Insulin	3.07 (0.97, 9.74)	0.05
PTH	2.53 (0.77, 8.32)	0.12
CA	3.30 (0.99, 11.00)	0.05
AST	3.65 (1.14, 11.67)	0.02
ALT	3.50 (1.08, 11.35)	0.03
CRP	5.52 (1.51, 20.17)	0.01
HOMA	3.11 (0.98, 9.84)	0.05
SBP	2.55 (0.77, 8.45)	0.12
	3.13 (0.95, 10.34)	0.06

Figure 2: Logistic regression analysis of association between vitamin D and BMI z-score category, controlling for cardiovascular risk factors in children (n=91).

ed odds ratios for BMIZ ranged from 2.96 when adjusting for whole grains to 3.71 when adjusting for n-3 FAs. As shown in Figure 2, most cardiovascular risk markers did not confound the adjusted odds ratio for BMIZ in the base model. Adjustment for hs-CRP increased the estimated odds ratio for BMIZ (OR=5.52; 95% CI: 1.51- 20.17; p=0.01).

CONCLUSION

Our data suggests that those children with high BMIZ > 2.5 had lower serum 25(OH)D levels as compared to those with low BMIZ<2.5. This study documented 47.3% of children studied having 25(OH)D levels <20 ng/mL. Significant associations with 25(OH)D levels were seen with older age, winter season and African American ethnicity and with higher BMIZ.

Furthermore, we found that the risk for vitamin D deficiency increases in children with severe obesity (BMIZ ≥ 2.5 SD) and is not significantly altered by ethnicity, nutrient intake or season. Such results have not been found in other studies assessing factors affecting vitamin D status in obese children. Factors associated with a higher prevalence of Vitamin D Deficiency include poverty, winter/spring season of serum collection, lack of vitamin D supplement use, low milk intake, greater TV/computer use, and low physical activity.¹ Our study suggests that although these factors are important, they do not necessarily influence the change in risk for vitamin D deficiency in this sample of severely obese children. Sample size might be one of the reasons for these differences. Our study included 91 children whereas the National Health and Nutrition Examination Survey (NHANES) study had 12,292 children in the sample. Another difference is the geographic location, as our population was mostly from Western Massachusetts.

Our study has several limitations. First, these results

arise from a cross-sectional study design. As such, the associations should be interpreted as co-relational. It would be premature to assume based on these results that reducing BMIZ would necessarily increase 25(OH)D. Such evidence would need to be derived from prospective studies of change in BMIZ. Second, the limited sample size led to rather imprecise estimates, particularly for the multivariate analyses. Small sample sizes can exaggerate estimates of the odds ratios derived from logistic regression, regardless of adjustment.³¹ We used only a crude measure of sun exposure as a covariate in our multivariable models. Although limited, the use of season was significant in our base model and the odds ratios associated with each season appeared reasonable.

The study's design was strengthened by the use of a validated FFQ^{23,24} for estimating vitamin D intake in this highrisk pediatric population and was pilot-tested in the clinical community of interest. The inclusion of a comprehensive list of multiple risk factors, as well as the use of a study population comprised of a racially and ethnically diverse sample of children from an urban location geographically at high risk for limited sunlight exposure, were other strengths of our study.

Given the numerous behavioral and environmental factors that influence sun exposure and its effect on 25(OH)D level, it is not clear that a self-reported survey would contribute more meaningfully to the model than the adjustment for season. This relationship needs further study, particularly in children. The question also arises as to whether better markers of 25(OH) D levels exist for obese children and adults, and this should be investigated in future research. Recently, Power,³² investigating both African American and White Americans, suggested that measurement of the bioavailable, 25(OH)D may provide a better marker of sufficiency compared with total 25(OH)D. It has also been proposed that serum free 25(OH)D may better reflect

Open Journal 👌

ISSN 2377-8385

the vitamin D action than total 25(OH)D, for which an Enzymelinked immunosorbent assay (ELISA) for serum free 25(OH)D has recently become available.³³ Given the normal PTH levels in the setting of low 25(OH)D levels, and the observed increase in 25(OH)D following weight loss in many studies, it appears the treatment of low 25(OH)D levels in these adolescents and children could be weight reduction, and not supplementation. Future randomized controlled trials (RCTs) addressing this research question should be conducted. Our findings for the inverse association between BMIZ classification and vitamin D deficiency warrant more research to further elucidate the specific mechanisms by which severe obesity, and potential change in obesity status may influence serum 25(OH)D levels.

ACKNOWLEDGEMENTS

SK and CW conceived the study, received grant support, and conducted/supervised the study. PV conducted the primary analysis of the data, and facilitated data modeling and interpretations. HG managed the literature review with updated references, and incorporated revisions of the paper for critical comments from coauthors. RC contributed critical edits, with updated references from the scientific literature. MG provided scientific reference updates. All authors contributed to the review and interpretation of data, and in the editing and approval of the final paper. SK and CW had primary responsibility for the final content presented. All authors critically revised the paper for intellectual content and read and approved the final draft.

This work was funded by UMass-Baystate Medical Center Collaborative Biomedical Research (CBR) Grant. (Co-PIs: Srimathi Kannan, PhD and Chrystal Wittcopp, MD) We would like to thank our participants for their support and commitment to this study.

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest.

REFERENCES

1. Turer CB, Lin H, Flores G. Prevalence of vitamin D deficiency among overweight and Obese US Children. Pediatrics. 2013; 131(1): e152-e161. doi: 10.1542/peds.2012-1711

2. Forsythe LK, Livingstone MB, Barnes MS. Effect of adiposity on vitamin D status and the 25-hydroxycholecalciferol response to supplementation in healthy young and older Irish adults. Br J Nutr. 2012; 107: 126-134. doi: 10.1017/ S0007114511002662

3. Wortsman J, Matsuoka LY, Chen TC, Lu Z, Holick MF. Decreased bioavailability of vitamin D in obesity. Am J Clin Nutr 2000; 72: 690-693. Web site. http://ajcn.nutrition.org/ content/72/3/690.short. Accessed October 18, 2016

4. Ross AC, Manson JE, Abrams SA, Aloia JF, Brannon PM,



Clinton SK. Clarification of DRIs for calcium and vitamin D across age groups. J Am Diet Assoc. 2011; 111(10): 1467. doi: 10.1016/j.jada.2011.08.022

5. Zittermann A, Frisch S, Berthold HK, Gotting C, Kuhn J, Kleesiek K. Vitamin D supplementation enhances the beneficial effects of weight loss on cardiovascular disease risk markers. Am J Clin Nutr. 2009; 89(5): 1321-1327. doi: 10.3945/ajcn.2008.27004

6. Clemens TL, Adams JS, Henderson SL, Holick MF. Increased skin pigment reduces the capacity of skin to synthesis vitamin D3. Lancet. 1982; 1(8263): 74-76. Web site. https:// www.ncbi.nlm.nih.gov/pubmed/6119494. Accessed October 18, 2016

7. Webb AR, Kline L, Holick MF. Influence of season and latitude on the cutaneous synthesis of vitamin D3: Exposure to winter sunlight in Boston and Edmonton will not promote vitamin D3 synthesis in human skin. J Clin Endocrinol Metab. 1988; 67(2): 373-378. doi: 10.1210/jcem-67-2-373

8. Dong Y, Pollock N, Stallmann-Jorgensen IS. Low 25-hydroxyvitamin D levels in adolescents: race, season, adiposity, physical activity, and fitness. Pediatrics. 2010; 125(6): 1104-1111. Web site. http://pediatrics.aappublications.org/content/125/6/1104.short. Accessed October 18th, 2016

9. Au LE, Economos CD, Goodman E, Must A, Chomitz VR, Sacheck JM. Vitamin D intake and serum vitamin D in ethnically diverse urban schoolchildren. Public Health Nutr. 2012; 15(11): 2047-2053. doi: 10.1017/S1368980012003217

10. Kumar J, Muntner P, Kaskel FJ, Hailpern SM, Melamed ML. Prevalence and associations of 25-hydroxyvitamin D deficiency in US children: NHANES 2001-2004. Pediatrics. 2009. 124(3): e362-e370. Web site. http://pediatrics.aappublications.org/content/124/3/e362. Accessed October 18, 2016

11. Holick MF. High prevalence of vitamin D inadequacy and implications for health. Mayo Clin Proc. 2006; 81(3): 353-373. doi: 10.4065/81.3.353

12. Kiess W, Bae YJ, Penke M, Geserick M, Kratzsch J. Vitamin D in health and disease: the global threat of vitamin D deficient rickets. J Pediatr Endocrinol Metab. 2016. doi: 10.1515/jpem-2016-0054

13. Litonjua AA. Childhood asthma may be a consequence of vitamin D deficiency. Curr Opin Allergy Clin Immunol. 2009; 9(3): 202-207. doi: 10.1097/ACI.0b013e32832b36cd

14. Atabek ME, Eklioglu BS, Akyürek N, Alp H. Association between vitamin D level and cardiovascular risk in obese children and adolescents. J Pediatr Endocrinol Metab. 2014; 27(7-8): 661-666. doi: 10.1515/jpem-2013-0379

Open Journal 👌

Openventio PUBLISHERS

15. Olson ML, Maalouf NM, Oden JD, White PC, Michele R. Vitamin d deficiency in obese children and its relationship to glucose homeostasis. *J Clin Endocrinol Metab.* 2012; 97(1): 279-285. doi: 10.1210/jc.2011-1507

16. Krebs NF, Himes JH, Jacobson D, Nicklas TA, Guilday P, Styne D. Assessment of child and adolescent overweight and obesity. *Pediatrics*. 2007; 120(4): S193-228. doi: 10.1542/peds.2007-2329D

17. Schwarzenbach G. The Complexones and their analytical application. *The Analyst.* 1955; 80(955): 713-729. doi: 10.1039/AN9558000713

18. Marshall TA, Eichenberger GJM, Broffitt B. Relative validity of the Iowa Fluoride Study targeted Nutrient Semi-Quantitative Questionnaire and the Block Kids' Food Questionnaire for estimating beverage, calcium, and vitamin D intakes by children. *J Am Diet Assoc*. 2008; 108: 465-472. doi: 10.1016/j.jada.2007.12.002

19. Block G. Invited commentary: Another perspective on food frequency questionnaires. *Am J Epid* 2001; 154(12), 1103-1104. doi: 10.1093/aje/154.12.1103

20. US Department of Agriculture: *USDA National Nutrient Database for Standard Reference*. Beltsville, Maryland, US: Nutrient Data Laboratory; 1998.

21. Institute of Medicine. *Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids*. Washington, USA: National Academics Press; 2000.

22. Willett WC, Howe GR, Kushi LH. Adjustment for total energy intake in epidemiologic studies. *Am J Clin Nutr.* 1997; 65(4): 1220S-1228S. Web site. http://ajcn.nutrition.org/ content/65/4/1220S.short. Accessed October 18, 2016

23. Sabo RT, Ren C, Sun SS. Comparing Height-Adjusted Waist Circumference Indices: The Fels Longitudinal Study. *Open J Endocr Metab Dis.* 2012; 2(3): 40-48. doi: 10.4236/ojemd.2012.23006

24. Field AE, Peterson KE, Gortmaker SL. Reproducibility and validity of a food frequency questionnaire among fourth to seventh grade inner-city school children: Implications of age and day-to-day variation in dietary intake. *Public Health Nutr.* 1999; 2: 293-300. Web site. https://www.ncbi.nlm.nih. gov/pubmed/10512564. Accessed October 18, 2016

25. Tsuji K, Maeda T, Kawane T, Matsunuma A, Horiuchi N. Leptin stimulates fibroblast growth factor 23 expression in bone and suppresses renal 1alpha,25-dihydroxyvitamin D3 synthesis in leptin-deficient mice. *J Bone Miner Res.* 2010; 25(8): 1711-1723. doi: 10.1002/jbmr.65

26. Sung C-C, Liao MT, Lu KC. Role of vitamin D in insulin

resistance. J Biomed Biotechnol 2012; 234195.

27. Giudice MDE, Grandone A, Cirillo G. Bioavailable vitamin D in obese children: The role of insulin resistance. *J Clin Endocrinol Metab.* 2015; 100(10): 3949-3955. doi: 10.1210/ jc.2015-2973

28. Touvier M, Deschasaux M, Montourcy M, Sutton A. Interpretation of plasma PTH concentrations according to 25OHD status, gender, age, weight status, and calcium intake: Importance of the reference values. *J Clin Endocrinol Metab.* 2014; 99(4): 1196-1203. doi: 10.1210/jc.2013-3349

29. Amini Z, Bryant S, Smith C, Singh R, Kumar S. Is the serum vitamin D-parathyroid hormone relationship influenced by obesity in children? *Horm Res Pediatr*. 2013; 80(4): 252-256. doi: 10.1159/000354645

30. Hansen KE. High-Dose Vitamin D: helpful or harmful? *Curr Rheumatol Rep.* 2011; 13(3): 257-264. doi: 10.1007/ s11926-011-0175-9

31. Nemes S, Jonasson JM, Genell A, Steineck G. Bias in odds ratios by logistic regression modelling and sample size. *BMC*. 2009; 9: 56. doi: 10.1186/1471-2288-9-56

32. Powe CE, Karumanchi SA, Thadhani R. Vitamin Dbinding protein and vitamin D in blacks and whites. *N Engl J Med* 2014; 27: 370(9): 880-881. doi: 10.1056/NEJMc131 5850#SA4

33. Aloia J, Dhaliwal R, Mikhail M. Free 25(OH)D and calcium absorption, PTH, and markers of bone turnover. *J Clin Endocrinol Metab.* 2015; 100(11): 4140-4145. doi: 10.1210/ jc.2015-2548