

Research

Corresponding author*C. V. Singh, BSc, MSc, PhD**

Professor

Department of Animal Genetics and Breeding

College of Veterinary and Animal Science

Govind Ballabh Pant University of

Agriculture and Technology

Pantnagar, US Nagar

Uttarakhand 263145, India

E-mail: cvsingh2010@gmail.com

Volume 1 : Issue 2

Article Ref. #: 1000VMOJ1108

Article HistoryReceived: September 2nd, 2016Accepted: October 22nd, 2016Published: October 25th, 2016**Citation**Singh VK, Singh CV. Comparative study of different sire evaluation methods using first lactation traits in crossbred cattle. *Vet Med Open J.* 2016; 1(2): 43-49. doi: [10.17140/VMOJ-1-108](https://doi.org/10.17140/VMOJ-1-108)**Copyright**

©2016 Singh CV. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Comparative Study of Different Sire Evaluation Methods Using First Lactation Traits in Crossbred Cattle

V. K. Singh¹; C.V. Singh, BSc, MSc, PhD^{2*}¹Narendra Dev University of Agriculture & Technology, Faizabad, UP, India²Department of Animal Genetics and Breeding, College of Veterinary and Animal Science, Govind Ballabh Pant University of Agriculture & Technology, Pantnagar, Uttarakhand 263145, India**ABSTRACT**

The records of 1085 crossbred cattle daughters of 57 sires maintained during 1974-2004 at Instructional Dairy Farm of Govind Ballabh Pant University of Agriculture & Technology, Pantnagar, Uttarakhand, India were used to evaluate sires for first lactation performance traits. The data were analyzed to estimate the breeding values of sires using least squares methods (LSM), best linear unbiased prediction (BLUP) and derivative free restricted maximum likelihood (DFREML) method. The average breeding values of sires estimated by all 3 methods were similar and estimated for age at first calving (AFC), first lactation period (FLP), first dry period (FDP) and first calving interval (FCI) as 1359.95 days, 356.56 days, 177.58 days and 534.15 days respectively. The estimated breeding values (EBV'S) showed large genetic variation between sires for first lactation. The EBV'S of sires had very high and significant product moment correlations and rank correlations among all first lactation traits. These results revealed that estimated breeding values of sires estimated for first lactation traits by LSM method showed small genetic variation in comparison to BLUP and DFREML methods. The LSM method was found to be more efficient, accurate, and stable with lowest genetic variation.

KEYWORDS: Best linear unbiased prediction (BLUP), Derivative free restricted maximum likelihood (DFREML), Breeding value; Genetic Variation; First lactation traits.

ABBREVIATIONS: LSM: Least Squares Methods; BLUP: Best Linear Unbiased Prediction; DFREML: Derivative free restricted maximum likelihood; EBV'S: Estimated Breeding Values; FCI: First Calving Interval; AFC: Age at First Calving; FDP: First Dry Period; REML: Restricted Maximum Likelihood.

INTRODUCTION

In India, most of the progeny testing is limited to organized farms with small herd size and this is a limiting factor in assessing the sires breeding values. Today, the need for sire evaluation for the additive genetic value (breeding value or transmitting ability) to its progeny has been recognized. Therefore, accurate, efficient and early evaluation of breeding value of sires is of prime importance. With advancement in computing power, many procedures have been proposed for evaluating sires based on the records of their progeny. In recent past the best linear unbiased prediction (BLUP) procedure has been widely used as standard method of sire evaluation. However, Henderson¹ opined that analysis of variance and covariance may give biased components of variance from selected population by BLUP method; whereas restricted maximum likelihood (REML) can give biased free estimate. Hence, the sire evaluation using advanced statistical techniques based on derivative free REML methods were also used to judge the effectiveness of different methods.

In the present study, the breeding values of sires were estimated, evaluated and ranked on the basis of DFREML as described by Meyer,² would be expected to enhance the accuracy of selection of bulls. However, an animal model in Indian breeds has been used preciously.^{3,4} Hence, in this paper an attempt is made to compare effectiveness of different methods of sire evaluation on the basis of first lactation traits in crossbred cattle.

MATERIALS AND METHODS

The data for the present study were obtained from the performance records of 1085 crossbred daughters of 57 sires maintained at Instructional Dairy Farm of Gobind Ballabh Pant University of Agriculture and Technology, Pantnagar, Uttarakhand, India during 1974-2004. They were used to estimate breeding value of sires for first lactation and life time traits. The exotic breeds used were Holstein Frisian (HF), Jersey (J) and Red Dane (RD) and the indigenous breeds were Sahiwal (S) and Rathi (R). The crossbreds were HFxS, JxS, RDxS, HFxSxRD, RDxSxJ, HFxRxJ and HFxSxRDxJ. Cows with abnormal and incomplete records were excluded from the study. The data on sires having at least 3 daughters per sire were included for sire evaluation of sires. Each year was divided into 3 seasons i.e. summer (March to June), rainy (July to October) and winter (November to February) based on climatic conditions. Daughters were grouped into 7 genetic groups on the basis of their breeding history. Period of study was divided into 7 periods, each period of 5 years in which their daughters were born.

STATISTICAL ANALYSIS

As the data in the present study were non-orthogonal in nature with unequal subclass numbers, they were subjected to least squares analysis of variance without interactions using different models to examine the effect of genetic as well as non-genetic factors on various first lactation traits as per standard procedures of Harvey.⁵ The model was based on the assumption that different components fitting in the model were linear, independent and additive. While sire was treated as random effect, the other genetic and non-genetic factors (genetic group, season and period) were taken as fixed effects in the model. The main objective of sire evaluation is to obtain an accurate and unbiased estimate of breeding value of bulls and ranking them on the basis of performance of their daughters so as to enable the breeders to choose the best bull for subsequent improvement of the herd. Many sire evaluation methods based on performance of adequate number of progeny recorded have been proposed. The following four methods of sire evolution were used to estimate the breeding values of sires.

Least Squares Method (LSM)

The least squares method (LSM)⁵ was used to estimate the breeding value of sires, using the following statistical model.

$$Y_{ij} = \mu + S_i + e_{ij}$$

Where,

Y_{ij} = jth dependent single trait of the daughter of ith sire

μ = population mean

S_i = effect of ith sire

e_{ij} = random error assumed to be normally and independently distributed with mean zero and variance σ^2 i.e. NID $(0, \sigma_e^2)$.

Best Linear Unbiased Prediction (BLUP F90-Dairy Pack, 2004)

BLUP F90 and related programs were developed in the lab of Misztal et al⁶ with the purpose of providing comprehensive computing capabilities to problems related to mixed models in animal breeding.

Dairy pack provides basic animal model which allows data of animals in the pedigree to be included in the analysis so that all known relationships can be taken into account. Other effects, fixed and random, can be including for comprehensive use of mixed model. Fixed effects such as herd-year-season can be performed during the analysis; therefore, no added step of data preparation is required. Normally, all traits used in dairy pack should be continuous rather than ordinal scale for proper use in the analysis of linear mixed model. Model with single record per animal can be analyzed by BLUP univariate (single trait) animal model as follows:

$$y = X\beta + Za + \varepsilon, \text{ and } V \begin{bmatrix} a \\ \varepsilon \end{bmatrix} = \begin{pmatrix} A\sigma_a^2 & 0 \\ 0 & I\sigma_e^2 \end{pmatrix}$$

Where, y is a vector of response variable, β is a vector of fixed effects, a is vector of random additive genetic effects, ε is a vectors of random residual, X and Z are incident matrices related to fixed and random effects, A is the numerator relationship matrix, σ_a^2 is the additive genetic variance, and $I\sigma_e^2$ is the residual variance.

To perform BLUP, Henderson's¹ MME can be written as:

$$\begin{pmatrix} X'X & X'Z \\ Z'X & Z'Z + \alpha A^{-1} \end{pmatrix} \begin{pmatrix} \beta \\ a \end{pmatrix} = \begin{pmatrix} X'Y \\ Z'Y \end{pmatrix},$$

Where $\alpha = \frac{\sigma_e^2}{\sigma_a^2}$

Multivariate Model

$$\begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{pmatrix} X_1 & 0 \\ 0 & X_2 \end{pmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} + \begin{pmatrix} Z_1 & 0 \\ 0 & Z_2 \end{pmatrix} \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} + \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \end{bmatrix}, V \begin{bmatrix} a_1 \\ a_2 \\ \varepsilon_1 \\ \varepsilon_2 \end{bmatrix} = \begin{pmatrix} G \otimes A & 0 \\ 0 & R \otimes I \end{pmatrix},$$

Where y_1, y_2 is a vector of response variable for trait 1

and $2, \beta_1, \beta_2$ is vectors of fixed effects, a_1, a_2 is vector of random additive genetic effects, ϵ, ϵ_2 is vector of random residual effect, X, W and Z are incident matrices related to fixed and random effects, A is a numerator relationship matrix, G is a matrix of direct genetic variance-covariance for trait 1 and 2. R is matrix of residual variance-covariance for trait 1 and 2.

Derivative Free Restricted Maximum Likelihood Method (DFREML)

In DFREML method the density function of the multivariate distribution is maximized after correcting all observations first for the fixed effects without taking consideration of the first or second derivatives in the analysis. Single and multiple trait models were used for estimation of different variance and covariance components. In multiple traits two or three traits were considered simultaneously.

For DFREML estimation following animal model was considered:

$$Y_{ijk} = X b_i + Z u_j + e_{ijk}$$

For three traits the model can be represented as:

$$\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix} = \begin{bmatrix} X_1 & 0 & 0 & 0 \\ 0 & X_2 & 0 & 0 \\ 0 & 0 & X_3 & 0 \\ 0 & 0 & 0 & X_4 \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix} + \begin{bmatrix} Z_1 & 0 & 0 & 0 \\ 0 & Z_2 & 0 & 0 \\ 0 & 0 & Z_3 & 0 \\ 0 & 0 & 0 & Z_4 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{bmatrix} + \begin{bmatrix} e_1 \\ e_2 \\ e_3 \\ e_4 \end{bmatrix}$$

Where,

- Y_{ijk} = k^{th} observation of j^{th} random effect of i^{th} fixed effect
- b_i = Vector of observation of fixed effect i.e. season and period of calving
- u_j = Vector of additive genetic effect (random effect/sire effect)
- X = Design matrix/incident matrix of fixed effect
- Z = Design matrix/incident matrix of random effect
- e_{ijk} = Vector of residual errors.
- With $E(y) = Xb$ and variance V .

The variance covariance structure given by

$$\text{Var} \begin{bmatrix} u_1 \\ u_2 \\ u_3 \\ e_1 \\ e_2 \\ e_3 \end{bmatrix} = \begin{bmatrix} A\sigma_{a1}^2 & A\sigma_{a12} & A\sigma_{a13} & 0 & 0 & 0 \\ A\sigma_{a21} & A\sigma_{a2}^2 & A\sigma_{a23} & 0 & 0 & 0 \\ A\sigma_{a31} & A\sigma_{a32} & A\sigma_{a3}^2 & 0 & 0 & 0 \\ 0 & 0 & 0 & I\sigma_{e1}^2 & I\sigma_{e12} & I\sigma_{e13} \\ 0 & 0 & 0 & I\sigma_{e21} & I\sigma_{e2}^2 & I\sigma_{e23} \\ 0 & 0 & 0 & I\sigma_{e31} & I\sigma_{e32} & I\sigma_{e3}^2 \end{bmatrix}$$

The following are the assumptions of the model:

$$\text{Var} (u) = G$$

- $\text{Var} (e) = R$
- and $\text{Cov} (u, e) = 0$
- So that, $V(y) = ZGZ' + R$

The normal distribution of a variable y with mean μ and standard deviation σ , the normal distribution of this variable represented as: $y \sim N(\mu, \sigma^2)$.

The mathematical distribution of the density function for this normally distributed variable is:

$$f(y) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}(y-\mu)^2/\sigma^2}$$

This is known as probability density function (PDF). In case of multivariate, multidimensional normal distribution PDF follows, $y \sim N(Xb, V)$. Where, N is the length of y and V is the variance of the parameter. This function $f(y)$ gives the probability of finding a certain 'y' such that the parameters are given. The parameters are the means in Xb "location parameters" and the variances in V "dispersion parameters". Observed data gives probability of having such data for certain parameter value. So the probability density function can be used as a likelihood function.

When the data y is known; the $f(y)$ is a likelihood function which can be maximized in parameters; in other words, find the parameters for which $f(y)$ has highest value is to be found.

The effectiveness of different sire evaluation methods was judged by the estimated breeding value of sires taken as twice the sire genetic group solution plus sire solution within sire genetic group for that trait. After estimation of breeding value of sire, the sires were given ranks as per their genetic merit.

After estimation of breeding value of sires, the sires were ranked as by their genetic merit. The Spearman's rank correlation between breeding values of sires derived by various methods was used to judge the effectiveness of different methods. The rank correlation was estimated as per Steel and Torrie⁷:

$$r_s = 1 - \frac{6 \sum d_i^2}{n(n^2 - 1)}$$

Where,

- r = rank correlation coefficient
- n = number of sires under evaluation
- d_i = difference of rank between paired items under 2 methods

The significance of rank correlation was tested by t -test as given below:

$$t = r \sqrt{\frac{(n-2)}{(1-r^2)}}$$

It was compared with *t*-table value with (n-2) degree of freedom.

RESULTS

In the present study, the breeding values of sires were estimated and sires ranked on the basis of breeding value of their progeny by 3 different sires evaluation methods and presented in Table 1. The estimated overall average breeding values of sires by LSM, DFREML and BLUP methods for age at first calving (AFC), first lactation period (FLP), first dry period (FDP) and first calving interval (FCI) were found to be 1359.95 days, 356.56 days, 177.58 days and 534.15 days respectively. The EBV'S for AFC, ranged from 1296.20 to 1414.70 days, for FLP 330.70 to 399.80 days for FDP 171.40 to 188.90 days and for FCI 509.50 to 562.00 days respectively. By DFREML method the estimated breeding value for AFC ranged from 1162.00 to 1522.20 days, for FLP 305.90 to 449.70 days, for FDP 137.80 to 254.60 days for FCI 470.20 to 616.20 days respectively. By BLUP method the estimated breeding value for AFC ranged from 1204.40 to 1512.70 days, for FLP 302.80 to 447.70 days, for first dry period 151.00 to 226.20 days and for FCI 469.30 to 610.60 days respectively.

The estimated breeding values of sires estimated for first lactation traits by LSM method showed small genetic variation in comparison to BLUP and DFREML methods. The BLUP method was considered to be second more appropriate method than the DFREML method. The product moment correlations between the breeding value of sires and the rank correlations between the rankings of sires by various methods are presented in Table 2. Accuracy of the methods of sire evaluation was de-

termined by comparing the simple (product moment correlations and spearman's rank correlation) correlation of breeding value of sires estimated by various sire evaluation methods for first lactation traits. The simple correlations and rank correlations among all the three methods of sire evaluation were very high and significant. The product moment correlations among LSM and DFREML and BLUP methods of sire evaluation ranged from 0.974 to 1.00, where as the rank correlations among breeding value of sires ranged from 0.994 to 0.998 for first lactation traits.

Top 10 sires ranked on the basis of estimated breeding values of sires for first lactation traits by all 3 methods are presented in Table 3. The top 10 sires ranked on the basis of first lactation traits revealed that all sires would not rank same for all the methods. However, the rank of sires for different sire evaluation methods revealed that 4-5% top sires almost had similar rank for all the methods.

In comparison of these methods of sire evaluation LSM method was found to be more efficient, accurate, and stable with lowest genetic variation. The rank correlation coefficients amongst breeding value of sires by different methods of sire evaluation showed reasonably high degree of agreement (79%) in the ranking of sires.

DISCUSSION

The estimated overall average breeding values of sires by LSM, DFREML and BLUP methods for AFC, FLP, FDP and FCI were 1359.95 days, 356.56 days, 177.58 days and 534.15 days respectively. The EBV'S for AFC, ranged from 1296.20 to 1414.70 days, for FLP 330.70 to 399.80 days for FDP 171.40 to 188.90 days and for FCI 509.50 to 562.00 days respectively. By DFREML method the estimated breeding value for AFC ranged

Traits	Sire evaluation method	Average breeding value	Minimum B.V. (% below average)	Maximum B.V. (% below average)	Number of sires over average BV (% of sires)	Number of sires below average BV (% of sires)
AFC	LSM	1359.95	1296.20 (4.69)	1414.70 (4.03)	26.00 (45.61)	31.00 (54.39)
	DFREML	1359.95	1162.0 (14.56)	1522.20 (11.93)	27.00 (47.37)	30.00 (52.63)
	BLUP	1359.95	1204.70 (11.42)	1512.70 (11.23)	27.00 (47.37)	30.00 (52.63)
FSP	LSM	271.51	270.50 (0.37)	272.30 (0.29)	21.00 (36.84)	36.00 (63.16)
	DFREML	271.50	268.80 (1.00)	273.50 (0.74)	25.00 (43.86)	32.00 (56.14)
	BLUP	271.50	269.80 (0.63)	272.80 (0.48)	25.00 (43.86)	32.00 (56.14)
FDP	LSM	177.58	171.40 (3.48)	188.90 (6.37)	32.00 (56.14)	25.00 (43.86)
	DFREML	177.58	137.80 (22.40)	254.60 (43.37)	32.00 (56.14)	25.00 (43.86)
	BLUP	177.57	151.00 (14.96)	226.20 (27.38)	32.00 (56.14)	25.00 (43.86)
FCI	LSM	534.15	509.50 (4.61)	562.00 (5.21)	33.00 (57.89)	24.00 (42.11)
	DFREML	534.15	470.20 (11.97)	616.20 (15.36)	31.00 (54.39)	26.00 (45.61)
FLP	LSM	356.56	330.70 (7.25)	399.80 (12.13)	27.00 (47.37)	30.00 (52.63)
	DFREML	356.56	305.90 (14.21)	449.70 (26.12)	27.00 (47.37)	30.00 (52.63)
	BLUP	356.56	302.80 (15.08)	447.70 (25.56)	27.00 (47.37)	30.00 (52.63)

Table 1: Average breeding value estimates for first lactation traits by different methods of estimation.

from 1162.00 to 1522.20 days, for FLP 305.90 to 449.70 days, for FDP 137.80 to 254.60 days for FCI 470.20 to 616.20 days respectively. By BLUP method the estimated breeding value for AFC ranged from 1204.40 to 1512.70 days, for FLP 302.80 to 447.70 days, for first dry period 151.00 to 226.20 days and for FCI 469.30 to 610.60 days respectively (Table 1). These estimates showed large genetic differences between the ranges of breeding values of sires for first lactation traits. Similar results were also reported by Dalal et al,⁸ Bajetha,⁹ Moges et al,¹⁰ Singh and Singh,¹¹ Bajetha and Singh¹² and Bajetha et al.¹³

The estimated breeding values of sires estimated for first lactation traits by LSM method showed small genetic variation in comparison to BLUP and DFREML methods. The BLUP method was considered to be second more appropriate than the DFREML method. The results obtained in the present study are in conformation to the reports of Tajane and Rai,¹⁴ Sahana and Gurnani,¹⁵ Deulkar and Kothekar,¹⁶ Kumar¹⁷ and Mukharjee.¹⁸ However, Dhaka and Dahia,¹⁹ Dahia,²⁰ Bajetha,⁹ Dubey et al²¹ and Bajetha and Singh¹² reported BLUP as best procedure in comparison to other procedures of sire evaluation.

Accuracy of the methods of sire evaluation was determined by comparing the simple (product moment correlations and spearman's rank correlation) correlation of breeding value

of sires estimated by various sire evaluation methods for first lactation performance traits. The simple correlations and rank correlations among all the three methods of sire evaluation were very high and significant. The product moment correlations among LSM and DFREML and BLUP methods of sire evaluation ranged from 0.974 to 1.00, where as the rank correlations among breeding value of sires ranged from 0.994 to 0.998 for first lactation traits (Table 2).

In the present study, all the estimates of simple and rank correlations were highly significant. These findings agreed with the reports of Dalal et al,²² Dubey et al,²¹ Banik and Gandhi,²³ Bajetha,⁹ Mukharjee et al,⁴ Kumar et al³ and Moges et al.¹⁰

The top 10 sires ranked on the basis of first lactation traits revealed that all sires would not rank same for all the methods. However, the rank of sires for different sire evaluation methods revealed that 4-5% top sires almost had similar rank for all the methods (Table 3). Similar results were also reported by Dalal et al,²² Bajetha,⁹ Dubey et al,²¹ Moges et al,¹⁰ Singh and Singh,¹¹ Singh et al²⁵ and Bajetha and Singh,¹² Lodhi et al,²⁴ Lodhi et al²⁶ and Singh and Singh.²⁷

In comparison of these methods of sire evaluation LSM method was found to be more efficient, accurate, and stable

	Traits under study														
	AFC			FLP			FDP			FCI			FSP		
	LSM	DFREML	BLUP	LSM	DFREML	BLUP	LSM	DFREML	BLUP	LSM	DFREML	BLUP	LSM	DFREML	BLUP
LSM	-	0.99	0.99	-	0.99	0.99	-	0.99	0.99	-	0.99	1.00	-	0.99	0.99
DFREML	0.97	-	0.99	0.99	-	0.99	0.98	-	0.99	0.98	-	0.99	0.99	-	0.98
BLUP	0.99	0.98	-	0.99	0.99	-	0.99	0.98	-	0.99	0.99	-	1.00	0.99	-

Table 2: Spearman's Rank and Product moment correlation among different sire evaluation methods.

Rank	AFC			FLP			FDP			FCI		
	LSM	DFREML	BLUP	LSM	DFREML	BLUP	LSM	DFREML	BLUP	LSM	DFREML	BLUP
1.	25	25	25	20	20	20	43	43	43	20	20	20
2.	38	46	38	21	64	21	23	35	23	25	25	25
3.	32	32	58	64	21	64	50	23	50	47	47	47
4.	58	58	32	42	42	42	19	50	19	42	53	42
5.	24	38	24	26	26	26	12	12	12	53	42	53
6.	40	24	46	44	44	44	26	26	35	21	56	56
7.	46	40	40	23	29	22	15	66	15	56	61	21
8.	26	41	53	22	50	23	66	19	66	45	45	45
9.	47	52	47	50	32	50	35	59	26	61	21	61
10.	34	34	26	29	22	29	33	28	59	29	29	29

Table 3: Top 10 sires ranks on the basis of estimated breeding values of sires for first lactation traits by different methods.

with lowest genetic variation. The rank correlation coefficients amongst breeding value of sires by different methods of sire evaluation showed reasonably high degree of agreement (79%) in the ranking of sires.

CONCLUSION

These results indicated that the estimated breeding values (EBV'S) of sires had very high and significant product moment correlations and rank correlations among all lifetime traits estimated by all the methods. These estimates indicating that there was higher degree of similarity (about 85-90%) in ranking of sires by different methods. However, the ranks of sires for different traits revealed that 4-5% top sires had similar rank for all life time traits. In comparison of these methods of sire evaluation LSM method was found to be more efficient, accurate and stable with lowest genetic variation. The rank correlation coefficients amongst breeding value of sires by different methods of sire evaluation showed reasonably high degree of agreement (79%) in the ranking of sires.

CONFLICTS OF INTERESTS

The authors declare that they have no conflicts of interest.

REFERENCES

- Henderson CR. Use of relationship among sires to increase accuracy of sire evaluation. *J Dairy Sci.* 1975; 58(11): 1731-1738. doi: [10.3168/jds.S0022-0302\(75\)84777-1](https://doi.org/10.3168/jds.S0022-0302(75)84777-1)
- Meyer K. *Derivative free restricted maximum likelihood (DFREML) programme version 3.0β users note.* Armidale, NSW 235, Australia: University of New England; 1998.
- Kumar A, Gandhi RS, Singh A, Haile A. Comparison of Animal model with other conventional methods of sire evaluation for milk production in karanfries cattle. *Indian J Anim Sci.* 2008; 78(12): 1393-1396. Web site. <https://cgspace.cgiar.org/handle/10568/1752>. Accessed September 1, 2016.
- Mukharjee S, Joshi BK, Gaur GK. Sire evaluation methods in frieswal cattle. *Indian J Anim Sci.* 2007; 77(8): 123-126.
- Harvey WR. *User guide for LSMLMW and MIXMDL package. Mix Model Least Squares And Maximum Likelihood Computer Programme.* Columbia, Chio, USA: PC-2 version mimeograph; 1990.
- Misztal I, Duangjinda M, Tsuruta S. *BLUPF 90, Dairy Pack Version 2; Genetic Evaluation Program for Dairy Cattle.* Athens, GA 30602, USA: Department of Animal and Dairy Science, The University of Georgia; 2004. Web site. <http://nce.ads.uga.edu/ignacy>. Accessed September 1, 2016.
- Steel RGD, Torrie JH. *Principles and Procedures of Statistics.* 2nd ed. New York, USA: Mcgraw Hill Book Co.; 1960.
- Dalal DS, Rathi SS, Raheja KL. Relationship between sires estimated breeding values for first lactation and lifetime traits in Haryana cattle. *Indian J Anim Sci.* 1999; 69(8): 592-595. Web site. <http://epubs.icar.org.in/ejournal/index.php/IJAnS/article/view/21174>. Accessed September 1, 2016.
- Bajetha G. *Selection of sires by using different sire evaluation methods in crossbred cattle,* [PhD Thesis]. Pantnagar, U. S. Nagar, Uttarakhand, India: G. B. Pant University of Agriculture & Technology; 2006.
- Moges TG, Singh CV, Barwal RS, Kumar D, Singh CB. Evaluation of sires using different multitrait sire evaluation methods in crossbred cattle. *Indian J Anim Sci.* 2009; 62(3): 1-4. Web site. <http://cat.inist.fr/?aModele=afficheN&cpsid=22178856>. Accessed September 1, 2016.
- Singh VK, Singh CV. Sire evaluation using animal model and conventional methods for milk production in crossbred cattle. *Indian J Anim Sci.* 2011; 81(1): 77-79. Web site. <http://epubs.icar.org.in/ejournal/index.php/IJAnS/article/view/2964>. Accessed September 1, 2016.
- Bajetha G, Singh CV. Efficiency of different sire evaluation methods to improve life time production traits in crossbred. *Cattle Livest Res Intern.* 2015; 3(3): 64-70. Web site. http://www.jakraya.com/journal/pdf/9-lriArticle_3.pdf. Accessed September 1, 2016.
- Bajetha G, Singh CV, Barwal RS. Sire evaluation by different methods in crossbred cattle. *Ameri Adv J of Natu Sci.* 2015; 1(1): 22-26. Web site. <http://casrp.co.uk/journals/index.php/AA-JNS/article/view/93>. Accessed September 1, 2016.
- Tajane KR, Rai AV. Efficiency of sire evaluation methods to improve milk yield of Sahiwal X Holstein Frisian cattle. *Indian J Anim Sci.* 1990; 60(2): 183-191. Web site. <https://www.cabdirect.org/cabdirect/abstract/19900178588>. Accessed September 1, 2016.
- Sahana G, Gurnani M. Effectiveness of sire evaluation methods for milk production along with auxiliary traits vis a vis other methods in crossbred cattle. *Indian J Dairy Sci.* 1996; 51(6): 439-442.
- Duelkar PB, Kothekar MD. Sire evaluation considering FLY for improvement of lifetime production in Sahiwal. *Indian J Anim Sci.* 1999; 69(4): 240-242.
- Kumar A. *Genetic analysis of economic traits of Sahiwal cattle using animal models,* [PhD Thesis]. Pantnagar, US Nagar, Uttarakhand, India: G. B. Pant University of Agriculture & Technology; 2003.

18. Mukharjee S. *Genetic evaluation of frieswal cattle*, [PhD Thesis, NDRI]. India: Deemed University Karnal; 2005.
19. Dhaka SS, Raheja KL. A comparison of sire evaluation methods. *Indian J Anim Sci*. 2000; 70(6): 643-644.
20. Dahia DS, Singh RP, Khanna AS. Genetic group differences and the effect of non-genetic factors in crossbred cattle for reproduction traits. *Indian J Anim Res*. 2003; 37(1): 61-64. Web site. <http://www.indianjournals.com/ijor.aspx?target=ijor:ijar1&volume=37&issue=1&article=016>. Accessed September 1, 2016.
21. Dubey PP, Singh CV, Prasad RB. Relationship between sires estimated breeding values for first lactation and lifetime traits and ranking of sires in Sahiwal and its cross. *Indian J Anim Sci*. 2006; 76(10): 824-828. Web site. <http://epubs.icar.org.in/ejournal/index.php/IJAnS/article/view/4424>. Accessed September 1, 2016.
22. Dalal DS, Rathi SS, Raheja KL. Relationship between sires estimated breeding values for first lactation and lifetime traits in Hariana cattle. *Indian J Anim Sci*. 1999; 69(8): 592-595. Web site. <http://epubs.icar.org.in/ejournal/index.php/IJAnS/article/view/21174>. Accessed September 1, 2016.
23. Banik S, Gandhi RS. Animal model versus conventional models of sire evaluation in Sahiwal cattle. *Asian Austr J Anim Sci*. 2006; 19(9): 1225-1228. doi: [10.5713/ajas.2006.1225](https://doi.org/10.5713/ajas.2006.1225)
24. Lodhi G, Singh CV, Barbal RS, Shahi BN, Dalal DS. Estimation of breeding values by different sire evaluation methods for selection of sires in crossbred cattle. *J Vet Sci Med Diagn*. 2015; 4: 5. doi: [10.4172/2325-9590.1000175](https://doi.org/10.4172/2325-9590.1000175)
25. Singh VK, Singh CV, Barwal RS, Shahi BN. Estimation of breeding values by different animal models for selection of sires in crossbred cattle. Published in Proceedings '10th World Congress on Genetics Applied to Livestock Production (WCGALP)' August 17-22, 2014: Vancouver, British Columbia, Canada.
26. Geeta L, Singh CV, Barwal RS, Shahi BN. Estimation of sires breeding value by animal model and conventional methods for milk production in crossbred cattle. *Journal of Agriculture and Crops*. 2016; 2(3): 31-34. Web site. http://econpapers.repec.org/article/arpjacarp/2016_3ap_3a31-34.htm. Accessed September 1, 2016.
27. Jaswant S, Singh CV. Evaluation of sires using different sire evaluation methods on the basis of first lactation traits in Sahiwal Cattle. *J Veterinar Sci Technol*. 2016; 7(2): 296(P 1-P6).