

Case Report

Corresponding author:*Maksym L. Golovakha, MD**

Chief

Department of Orthopedics &
Traumatology

Zaporozhye State Medical University

Zaporozhye Oblast 69035, Ukraine

Tel. +380-67-563-0117; 061-7663396

Fax: 061 7698193

E-mail: golovaha@ukr.net

Volume 1 : Issue 4

Article Ref. #: 100SEMOJ1118

Article History:Received: April 14th, 2015Accepted: September 10th, 2015Published: September 10th, 2015**Citation:**

Golovakha ML, Aghayev E, Orljanski W. Early loading after closed wedge high tibial osteotomy for knee arthritis. *Sport Exerc Med Open J.* 2015; 1(4): 114-117. doi: [10.17140/SEMOJ-1-118](https://doi.org/10.17140/SEMOJ-1-118)

Copyright:

© 2015 Golovakha ML. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Early Loading after Closed Wedge High Tibial Osteotomy for Knee Arthritis

Maksym L. Golovakha^{1*}, Emin Aghayev² and Wenjamin Orljanski³¹Chief, Department of Orthopedics & Traumatology, Zaporozhye State Medical University, Zaporozhye Oblast 69035, Ukraine²Institute for Surgical Technology and Biomechanics, University of Bern, Stauffacherstrasse 78, Bern 3014, Switzerland³Department of Orthopedic Surgery, Vienna Private Clinic, Pelikangasse 15, Vienna 1090, Austria**ABSTRACT**

Operative treatment approaches for the arthritis in the medial tibiofemoral compartment combined with a varus deformation of the knee are closed wedge high tibial osteotomy or knee arthroplasty. In contrast to the knee arthroplasty, tibial osteotomy maintains primary the function of natural knee joint. However, the limitations of the method are the necessity of postoperative immobilization, late term of loading and long period of functional recovery. The study on early loading after closed wedge high tibial osteotomy as the treatment of knee arthrosis was initiated. Two group of different postoperative rehabilitation approach were allocated. The first group included 22 patients in whom the loading was applied after 6th postoperative week, and the second group included 15 patients with early weight bearing at 3rd-5th postoperative day. Estimation of complications and comparison of the results of the treatment in two groups was carried out. The short- and long-term results of the treatment in the both groups showed similar distribution of good and poor result without significant differences. Studying rehabilitation dynamic, early loading allowed for faster postoperative functional recovery. No statistically significant differences regarding postoperative complications between the groups were observed. Our results indicate that early weight bearing after high tibial osteotomy does not worsen process of rehabilitation and result of the treatment. The absence of statistical differences regarding postoperative complications indicates the safety of the approach. In conclusion, we assumed that after closed wedge high tibial osteotomy the early postoperative loading is an advantageous rehabilitation approach in comparison with the late loading due to a faster functional recovery and a shortening of the patient hospitalisation. Nevertheless, further multi-patient studies on this approach with similar long-term follow ups are necessary.

KEYWORDS: Knee; Arthritis; High tibial osteotomy; Loading; Closed wedge.**INTRODUCTION**

Treatment of gonarthritis by closed wedge high tibial osteotomy is a widely applied method in practice of modern orthopaedics.¹⁻³ Numerous scientific works, which recently appeared in the periodic literature, confirm both high interest to the given problem and efficiency of this method.^{4,7} The discussion deals basically with the choice of correct indications for the osteotomy, and also with selection of a method for its carrying out.

Currently, high tibial osteotomy is recognised as a confident alternative to total knee prosthesis. At first sight, total knee prosthesis is more advantageous. It has a short term of postoperative rehabilitation, and provides more precise positive outcome of the treatment. In contrast, the basic advantage of closed wedge high tibial osteotomy within the treatment of knee arthritis is the opportunity to maintain the function of natural knee joint. It is well documented

that the choice of closed wedge high tibial osteotomy by correct management of its indications gives high density of successful results of treatment.⁴⁻⁷

By the combination of arthritis in the medial tibiofemoral compartment and varus deformations of the knee, the traditionally applied treatment is the closed wedge high tibial osteotomy through external access with osteosynthesis by staples.^{2,3} The limitations of the method are the necessity of postoperative immobilization, late term of loading and long period of functional recovery. Furthermore, it is necessary to note a small range of correction for the varus deformation. From different medical sources, the maximum of correction makes between 15 and 25°.^{2,4,7}

The studies performed in our department on early loading after arthroscopy and total knee arthroplasty have forced us to reconsider the approach for postoperative rehabilitation after closed wedge high tibial osteotomy. The following study describes the effects of early loading after closed wedge high tibial osteotomy as the treatment of gonarthritis.

MATERIAL AND METHODS

The study included 37 patients, which were treated at our department by closed wedge high tibial osteotomy during the period between 1997 and 2004. All the patients were surveyed in dynamic. Thirty three were female and four male. As the indication for the operation was knee arthritis with primary lesion of the medial tibiofemoral compartment. A retrospective analysis of the treatment results of the patients was performed.

For the estimation of the joint condition a knee x-rays with functional loading (standing) were carried out prior to operative treatment. The roentgenograms formed a basis for planning and calculation of axis correction. Subsequently, operations were carried out under general anesthesia.

At the first stage, an arthroscopy checking joint surfaces, patella position, synovial capsule, presence of intraarticular bodies, the condition of meniscuses and ligaments was performed and reasons for flex contracture were assessed. Removal of the intraarticular bodies, partial resection of the meniscus, local synovectomy and fossa plastic were carried out for restoration of the full extension.

The operation using the external curved approach to the proximal tibia made the second stage. The osteotomy was performed by a chisel or oscillation saw according to the preoperative plan. The osteosynthesis was carried out using staples. For prevention of postoperative complications antibacterial prophylactic and Fraxiparin-forte for 14 days were applied.

According to the postoperative period two different

groups of patients were allocated. The first group included 22 patients (two male and 20 female) in the age between 38 and 78 years who were operated during the period from 1997 till 2000. In this group isometric gymnastics began on the second postoperative day and immobilization was used within four weeks. The splint removal was followed by rehabilitation course. Partial weight bearing was allowed after 6th-8th and full bearing after 10th-12th postoperative week.

The second group included 15 patients (two male and 13 female) in the age between 42 and 68 years which were treated during the period from 2002 till 2004. The rehabilitation course in this group began also with isometric exercises on the second postoperative day and immobilization was used within two or four weeks. Partial weight bearing was allowed already after 3rd-5th postoperative day. Walking with a stick began after 4th-6th postoperative week depending on physical condition of the patient. As immobilisation, in two patients a brace was applied from 7th postoperative day. For all other patients a plaster cast was used during all period of fixation.

For the knee assessment at follow ups a standard "Knee Society" scale was applied.⁸ For the estimation of the effect of early loading after the osteotomy the analysis of the postoperative period and, importantly, of the early complications were performed.

The short-term follow up for both groups was performed on the 4th postoperative month and one year after operation. The long-term follow-ups for the first group lay between 5th and 8th and for the second group between 1st and 3rd postoperative year. Statistical differences of the treatment results in the groups were assessed using Student's t-test (two tailed, $\alpha=0.05$).

RESULTS

In the first group of patients the estimation of the short-term results using the "Knee Society" scale showed (on the average) following points combinations: preoperative - 98 points; 4th months follow up - 145 points and one year follow up - 167 points. In the second group of patients comparable points combinations were documented: preoperative - 102 points; 4th months follow up - 167 points and one year follow up - 172 (Diagram 1) (Figure 1).




Diagram 1: The results of the treatment of patients using the "Knee Society" score at the short-term follow ups.



Figure 1: Case report: a 51-year-old male was treated using closed wedge high tibial osteotomy. Immobilization was 2 weeks. Partial weight bearing was allowed from the 3rd postoperative day. The images shows a) the preoperative x-ray; b) the x-ray performed at the second postoperative day; c) the x-ray carried out at the follow up of 8 years; d) the functional result of the patient at the 10 years follow up.

The long-term results of the treatment in the first group (late loading) of the patients were excellent in four, good in 12, satisfactory in four and unsatisfactory in two patients. The long-term results in the second group (early loading) of the patients were excellent in four, good in eight, satisfactory in two and unsatisfactory in one patient (Diagram 2) (Figure 1).




Diagram 2: The results of the treatment of patients using the "Knee Society" score at the long-term follow ups.

The comparative analysis of functional recovery in the first and second group of patients showed that early loading allowed for faster postoperative rehabilitation. This was confirmed by the computer system for gait analysis. This is also indicated by significantly different ($p < 0.05$) results in the two groups at the 4 months follow up (Diagram 1).

The number of complications after operations in researched group of patients was not too large (Table 1). Significant differences in complications between the first and second group of patients were not seen (Table 1). Thus, application of early weight bearing after closed wedge high tibial osteotomy has not led to increase in the number of postoperative complications.

No.	Complication	Late loading	Early loading	Total
1	Deep venous thrombosis	1	1	2
2	Delayed union	1	0	1
3	Recurrence of the varus deformations	3	2	5
4	Total	5	3	8

Table 1: The distribution of postoperative complications after closed wedge high tibial osteotomy between the two studied groups.

DISCUSSION

Currently, development of techniques and approaches for operative interventions in orthopaedics goes on the way of achievement of the shortest possible rehabilitation period, which is mainly due to the high hospital costs for patients stay.

The idea for this study was born after introduction of early full loading after total hip and knee arthroplastic in our clinic. This technique has been borrowed from our German colleagues, and it has repeatedly passed check, both on efficiency and on safety.⁹ Observing patients after total hip arthroplasty, we concluded that early loading does not slow down but accelerates process of functional recovery of the joint. Similarly, considering our experience with knee arthroscopy, which was initially, before 10 years, followed by late weight bearing on the 4th or even 10th postoperative day and currently is the subject of one day hospitalisation, an early loading is advantageous.

Additionally to these ideas, the other reason for this study was the fact that three of our patients have began with early loading without our advice and in contrary to our recommendations after they were discharged from the hospital (within the 2nd postoperative week). At the follow up of the 4th postoperative week each of these patients showed a good range of motions and a good radiologically confirmed consolidation of the osteotomy.

The study was performed on two groups of patients with comparable age and sex distribution. The indication for the treatment of knee arthritis by closed wedge high tibial osteotomy was carefully examined in all patients.

Our short- and long-term results of the treatment in the both studied groups showed similar distribution of good and poor result without significant differences. This indicates that early weight bearing after high tibial osteotomy does not worsen process of rehabilitation and result of the treatment. In contrary, early loading allowed for faster postoperative functional

recovery. The absence of statistical differences regarding postoperative complications indicates the safety of the approach.

In the presented study, the second group of patients, in which the early postoperative loading was applied, included 15 patients. Furthermore, long term follow ups for the groups were different. Although, in comparison with the group of late loading, good results of the treatment and even faster functional recovery of the patients with early loading was documented, further studies using the described rehabilitation approach are necessary.

In conclusion, based on our results we assumed that after closed wedge high tibial osteotomy the early postoperative loading is an advantageous rehabilitation approach in comparison with the late loading due to a faster functional recovery and a shortening of the patient hospitalisation. Nevertheless, further multi-patient studies on this approach with similar long-term follow ups are necessary.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare.

CONSENT

The authors have obtained written informed consent from the patient for submission of this manuscript for publication.

REFERENCES

1. Levenets VN, Pljatsko VV. Diagnostic arthroscopy at gonarthrosis. *Orthopedics, traumatology, prosthesis*. 1985; 15: 60-65.
2. Loskutov AE, Golovakha ML. Arthroscopy and wedge osteotomy at treatment of gonarthrosis. *Bulletin of Orthopedics, traumatology, prosthesis*. 2002; 2: 5-7.
3. Popov VA. High tibial osteotomy at knee arthrosis. *Orthopedics, traumatology and prosthesis*. 1984; 2: 54-55.
4. Billings A, Scott DF, Camargo MP, Hofmann AA. High Tibial Osteotomy with a Calibrated Osteotomy Guide, Rigid Internal Fixation and Early Motion. Long-Term Follow-up. *The Journal of Bone and Joint Surgery*. 2000; 82: 70-79.
5. Koshino T, Yoshida T, Ara Y, Saito I, Saito T. Fifteen to twenty-eight years' follow-up results of high tibial valgus osteotomy for osteoarthritic knee. *The Knee*. 2004. 11(6): 439-444.
6. Rozkydal Z, Kura V, Ondrusek S. The arthroscopic debridement in the management of osteoarthritis of the knee joint by high tibial osteotomy. *Bratisl Lek Listy*. 2003; 104(11): 362-366.
7. Tang WC, Henderson IJP. High tibial osteotomy: Long term survival analysis and patients' perspective. *The Knee*. 2005; 12(6): 410-413.
8. Insall JN, Dorr LD, Scott R, Scott WN. Rationale of The knee society clinical rating. *Clin Orthop*. 1989; 248: 13-14.
9. Bosch HC, Springorum HW. Immediate loading after cement Wi Huftgelenkersatz. *Mod.Orthop.technik*. 1992; 112(5): 252-256.