

Research

*Corresponding author

Fabio Apone, PhD

Vitalab srl / Arterra Bioscience srl

via B. Brin 69

80142 Naples, Italy

Tel. +39 0816584411/6584386

Fax: +39 0812144864

E-mail: fapone@arterrabio.it

Volume 1 : Issue 1

Article Ref. #: 1000TCOJ1103

Article History

Received: April 18th, 2016

Accepted: May 12th, 2016

Published: May 17th, 2016

Citation

Bimonte M, De Lucia A, Carola A, et al. *Galdieria sulphuraria* relieves oily and seborrheic skin by inhibiting the 5- α Reductase expression in skin cells and reducing sebum production in vivo. *Trichol Cosmetol Open J.* 2016; 1(1): 11-18.

doi: [10.17140/TCOJ-1-103](https://doi.org/10.17140/TCOJ-1-103)

Copyright

©2016 Apone F. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Galdieria sulphuraria Relieves Oily and Seborrheic Skin By Inhibiting the 5- α Reductase Expression in Skin Cells and Reducing Sebum Production *In Vivo*

M. Bimonte, PhD¹; A. De Lucia, PhD¹; A. Carola, PhD¹; A. Tito, PhD¹; S. Buono, PhD²; A. L. Langellotti, PhD²; V. Fogliano, PhD³; G. Colucci, PhD⁴; Fabio Apone, PhD⁴

¹Arterra Bioscience, srl via Benedetto Brin, 69 80142, Napoli, Italy

²CAISIAL, University of Naples Federico II, Parco Gussone - Edificio H Via Università 100, 80055 Portici, Naples, Italy

³Food Quality and Design Group, Wageningen University, Bornse Weilanden 9, 6708 WG Wageningen, Netherlands

⁴Vitalab srl / Arterra Bioscience srl, via B. Brin 69, 80142 Napoli, Italy

ABSTRACT

The sebum has important protective functions for the skin, although its overproduction may lead to undesired aesthetic problems and even more serious skin diseases. Sebum overproduction is caused by the androgen hormone testosterone, which is regulated by the enzyme 5- α Reductase, and can be associated with an over-proliferation of parasitic microbes that cause serious infections linked to skin inflammatory reactions. From the extremophile microalga *Galdieria sulphuraria*, we obtained a water-soluble extract and we found that it was capable of inhibiting the enzyme 5- α Reductase, inducing the expression of the β -defensins, which represents the first defense response mechanism triggered by the skin cells to fight against undesired microbes, and stimulating the wound healing process in skin cell cultures. The microalga extract was finally employed in clinical tests on human volunteers and it was confirmed its capacity to regulate sebum production. The results suggested that the *Galdieria* extract had interesting potentialities as active ingredient in cosmetic and dermatological formulas, in particular in those addressed to oily and seborrheic skins.

KEY WORDS: Skin cell cultures; Microalga *Galdieria*; Sebum regulation; 5- α Reductase; β -Defensins.

INTRODUCTION

The skin is the first barrier of our body with important protective functions. Its protective role is dependent on the sebum, a complex mixture of lipids, triglycerides, wax esters, cholesterol esters, sterols, squalene and free fatty acids, all essential components for the formation of the hydro-lipid film which coats the skin and the scalp. The function of the hydrolipidic film is to lubricate and moisturize the skin surface, allowing the skin to maintain an appropriate level of hydration and protecting it against microbial invasions.

Sebum is produced by small specialized organs, the sebaceous glands, which are located over the entire body, with the exception of the hand palms, feet soles and some genital areas: they are more abundant and productive on the face and scalp, and usually associated with hair follicles. Although the sebum has got an important protective role for the skin, its overproduction by the sebaceous glands may lead to aesthetic problems and even more serious skin disfunctions.

Seborrhea is one of the most common cosmetic problems affecting both men and

women, typically highest in the age range of 15-35 years and declining throughout the adult age range,¹ and affecting more Asian than Caucasian people. Seborrhea gives an oily, shiny and greasy aspect to the skin and it is frequently accompanied by large pores. Many individuals feel embarrassed and annoyed with the appearance of their oily skin, often associated with a feeling of dirtiness.

Sebum production is influenced by several factors, among which the most important is the level of the androgen hormone testosterone: indeed the massive activation of the sebaceous glands occurs during puberty, when the production of testosterone is more intense. The testosterone concentration influences the production of sebum as it is converted, at level of sebaceous gland, into dihydro-testosterone by the enzyme 5- α Reductase. There are two isoenzymes of 5- α Reductase, the 5- α Reductases 1 (5 α R1) and the 5- α Reductases 2 (5 α R2), both of them have the capacity to convert testosterone into dihydro-testosterone, but they have different tissue distribution and functions.² The 5 α R1 is mainly expressed by sebaceous cells, whereby 5 α R2 is preferentially produced in the hair follicles and prostate; moreover, the activity of 5 α R1 is significantly higher in sebaceous glands from face and scalp compared to non-acne prone areas.³ Thus, a down regulation of 5 α R1 in the skin represents one of the solutions to contrast excessive sebum production, and some other strictly related diseases, such as seborrheic dermatitis and acne.

In seborrheic skins the overproduction of sebum is usually associated to a massive bacterial proliferation, more commonly the species of *Staphylococcus* and *Propionibacterium acnes*, because the sebum contains nutrients and thus represents a substrate for bacteria growth.^{4,5} The β -defensins, small cationic peptides, are the major components of innate immunity, a non-specific immune defense response against microorganism attack and hyper-proliferation. They have a strong antimicrobial, antiviral and antifungal activity,⁶ since their particular cationic nature allows to destroy the microbial membrane, negatively charged, by binding it, and leave intact human cell membranes that have a neutral charge. Thus defensins represent the initial defense response against microbial invasion. To date, three human β -defensins (hBD) have been shown to play a role in the defense of the skin, namely human β -defensin1, 2 and 3 (hBD1, hBD2, hBD3). They are expressed in all human epithelial tissues, either constitutively or up regulated in response to microbial or pro-inflammatory stimuli.⁷ In patients with acne the expression of the hBD2 is increased in the sebaceous glands,⁸ where it is likely to exert an important role in protecting from microbial invasion.

Among the unicellular organisms, the red microalga *Galdieria sulphuraria* has attracted the attention of applicative research, because it has well adapted to the extreme environment of the volcanic calderas:⁹ it is able to withstand high temperatures and grow in high-sulfur growth media, at pH-1. Its pres-

ence on the walls behind the Bocca Grande of volcanic crater Solfatara at Pozzuoli, near Naples, has been documented for several years.¹⁰ Recent studies have shown that this microalga is capable of synthesizing a wide range of antioxidant compounds, including the sulfur containing glutathione, making it an excellent candidate for skin care applications.¹¹ Indeed, the use of sulfur in dermatology has demonstrated efficacy in the treatment of many dermatological disorders including acne vulgaris, rosacea, seborrheic dermatitis.^{12,13} Moreover, for its high content of vitamins, essential fatty acids, proteins, carotenoids and phycocyanins, *Galdieria* has even been proposed as dietary supplement and food ingredient.¹⁴

On the basis of these properties, we investigated the potential benefits of a hydrosoluble extract obtained from *Galdieria sulphuraria* on the skin, in particular as cosmetic ingredient aimed at balancing sebum production. We demonstrated that *G. sulphuraria* extract was able to reduce the 5 α R1 expression in skin cells, to reinforce the skin's natural defense system by up regulating hBD2 expression and to promote the skin integrity accelerating the wound healing process. A clinical study performed on human volunteers confirmed the efficacy of *G. sulphuraria* extract as potential active ingredient for skin care applications, particularly directed to patients with oily and seborrheic skin.

MATERIAL AND METHODS

Galdieria sulphuraria cultivation and extract preparation

G. sulphuraria (strain n. 017) was obtained from ACUF (Algal Collection of the University of Naples Federico II, Italy), originally collected from a thermal spring site in Pozzuoli, Napoli (Italy). Stock cultures of axenic *G. sulphuraria* strain were maintained routinely on both liquid and agar slants of Allen medium¹⁵ in heterotrophic conditions, being the glucose (30 g L⁻¹) the main source of organic carbon (Figure 1). The biomass production was carried out in batches, and the culture flasks were incubated at a temperature of 26 \pm 2 °C with continuous air insufflation. After 2 weeks of growth, the cultures were harvested by a continuous centrifuge (2,500 g); the collected biomass was rinsed with 10 L of distilled water and frozen in sterilized containers. For the preparation of *G. sulphuraria* extracts, the frozen *G. sulphuraria* biomass was suspended in two volumes of phosphate-buffered saline (PBS) pH 7.4, and homogenized in an industrial homogenizer. The resulting lysate was centrifuged at 4,000 rpm to precipitate the particulate fraction and isolate the soluble components. The supernatant was then collected, adjusted to pH 6.5 using NaOH, and lyophilized until obtaining a dry powder, that represented the *Galdieria sulphuraria* Hydrosoluble Extract (GsHE). The powder was then dissolved in water or cell culture medium at the required concentrations for the different analyses. Three independently grown batches were tested in the different experiments.




Figure 1: Microscopic picture of the microalga *Galdieria sulphuraria* taken by an Inverted Microscope Motic BA. Magnification=1000 X; the black bar corresponds to 20 μm .

Cell Viability Assay (MTT Assay)

1.3×10^4 cells (keratinocytes, HaCat) were seeded in each well of a 96-well plate, grown for 8 h and treated for 48 h with different concentrations of GsHE. After treatments, cells were washed with PBS and incubated with 100 μl per well of “reaction buffer” containing: 10 mM Hepes, 1.3 mM CaCl_2 , 1 mM MgSO_4 , 5 mM glucose and 0.5 mg ml^{-1} of colorimetric substrate MTT [3-(4,5-dimethylthiazolyl)-2,5-diphenyltetrazoliumbromide] in PBS buffer at pH 7.4, according to the method described by Mosmann et al.¹⁶ After 3 h at 37 °C and 5% CO_2 , cells were solubilized by the addition of 100 microliters of solubilization solution (10% TritonX-100, 0.1 N HCl in isopropanol), and the plate incubated for 4 h at room temperature. The number of healthy cells was directly proportional to the level of the formazan product created. The developed color was then quantified at 595 nm by the multi well-plate reader Victor3 (Perkin Elmer, Waltham, MA, USA).

5 α -Reductase type 1 and human β -Defensin 2 gene expression analysis

For the gene expression analyses of the 5- α Reductase type 1 (5 α R1) and the human β -Defensin 2 (hBD2), Human Dermal Fibroblasts (HDF) and keratinocytes (HaCat) were used respectively. The cells were incubated with GsHE for 6 or 8 h, and then processed for RNA extraction using a RNA Purification Kit according to the manufacturer’s instructions (GenElute Mammalian Total RNA Miniprep Kit, Sigma-Aldrich). Reverse transcription polymerase chain reaction (RT-PCR) was performed using gene-specific primers and an oligo-nucleotide internal standard Kit (Quantum RNA™ 18S Ambion, Austin, Texas, USA). The PCR products obtained were visualized and quantified with an Imaging system (Geliance 200, Perkin Elmer, Waltham, MA, USA). The amplification band corresponding to the analyzed gene was normalized to the amplification band corresponding to the internal standard 18S. The values obtained were finally converted into percentage values by considering as 100% the measure of the samples of untreated cells.

Scratch Assay

For the scratch assay, HDF were seeded in 6-well plates. 12 h later, same areas of each well were displaced by scratching a line through the cell layer by a pipet tip and floating cells were removed by PBS washing. Culture medium containing 0.5% FBS with or without GsHE was added and the cells incubated for 7 hours. Mitomycin C (10 $\mu\text{g } \mu\text{l}^{-1}$) was always included in the medium to prevent cell proliferation and evaluate the cell migration capacity. To estimate the wound healing capacity of the cells, the unclosed cell-free area at time 0 and the one at 7 hours after the treatments were compared for each condition by using the software Image J.

Clinical Test

The study was performed on 20 male and female volunteers, aged from 18 to 50, who had an acne-prone face skin, although not suffering from any disease to the skin areas to treat. Each subject applied a cream containing 0.002% of GsHE, twice a day for a period of 30 days on the right side of the face massaging until absorption; the left side was treated with a placebo and the forehead remained untreated for the entire duration of the study. The amount of skin sebum excretion was measured on the lateral creases of the nose and on the forehead, before the treatments (T0), and after 15 days (T15) and 30 days (T30) of a bi-daily application of the product, by using the Sebumeter® SM815 (Courage-Khazaka GmbH, Germany), a digital probe for instrumental measure of the cutaneous sebum amount.¹⁷ The study was carried out under standard environmental conditions for each reading time, maintaining constant room temperature and monitoring humidity by a Thermo-hygrometer (Taylor Precision, NM).

RESULTS AND DISCUSSION

Calculation of the usage doses for GsHE

To evaluate the potential cell toxicity and determine the correct

doses of GsHE to use in the bioassays, we performed a cytotoxicity test (MTT assay) by testing scalar concentrations of GsHE, ranging from 0.5% to 0.0006%, on HaCaT keratinocytes. After 48 h, the vitality rate was determined. The IC₅₀ calculated was 0.198 %, and the highest non-toxic dose (which gave no mortality) was 0.01% (data not shown). Thus, the concentrations of GsHE chosen for the studies on skin cells were 0.002 and 0.01 %.

GsHE on 5αR1 Expression

In order to study whether *G. sulphuraria* had the capacity to affect the activity of 5αR1, the main enzyme regulating sebum production in the skin, we used GsHE at the 2 established concentrations to treat HDF for 6 h. Fibroblasts express 5αR1 in normal conditions and therefore are an appropriate model to investigate 5αR1 activity and regulation.¹⁸ After treatment, the cells were collected, RNA extracted and the expression of 5αR1 gene analyzed by RT-PCR. As shown in Figure 2, the treatments with both the concentrations of GsHE reduced the expression of 5αR1 (- 45% for 0.01% and -35% for 0.002%), suggesting a potential role of the extract in reducing the sebum production in the skin by acting through an inhibition of 5αR1 expression.




Figure 2: RT-PCR analysis of gene expression in fibroblasts treated with GsHE at the indicated concentrations. After RNA extraction, the expression level of the 5αR1 gene was analyzed by RT-PCR. The values obtained from the PCR band quantification were normalized to the standard amplification bands of 18S, reported in the column graph and expressed as percentages to the control, arbitrarily set as 100%. The error bars represent standard deviations, and the asterisks indicate statistically significant measures (* $p < 0.01$; ** $p < 0.001$).

GsHE on hBD2 Expression

As explained in the introduction, an overproduction of sebum can be often associated to a hyper proliferation of microbes in the skin, leading to skin injuries and to the triggering of inflammatory processes. Therefore, β-defensins represent an efficient defense mechanism developed by the skin cells to fight microbe proliferation. In order to test the ability of GsHE to induce the skin's natural defense system, gene expression analysis of hBD2 was performed in human keratinocytes treated with the two different concentrations of GsHE. The keratinocytes are the cells mostly exposed to microbial attacks, thus more keen to respond by activating a rapid production of inducible β-defensins. As shown in Figure 3, both the doses of GsHE increased hBD2

expression, but the lowest concentration of 0.002% was even better than the higher one to induce hBD-2 (170 % increase *versus* 60%), suggesting a good potential stimulatory effect on the innate skin defense system. Three independent extract preparations gave all the same result, indicating that this inverted dose response relationship was reproducible. The higher efficacy of the lower dose compared to higher one could be explained by considering that the extracts are mixtures of several types of molecules that may act synergically or counteract each other whether negative regulators become predominant. In this case, the choice of the right concentration of the extract would be of primary importance to produce a final positive effect on the cells both *in vitro* and *in vivo*.




Figure 3: RT-PCR analysis of gene expression in keratinocytes treated with GsHE at the indicated concentrations. After RNA extraction, the expression level of the BD2 gene was analyzed by RT-PCR. The values obtained from the PCR band quantification were normalized to the standard amplification bands of 18S, reported in the column graph and expressed as percentages to the control, arbitrarily set as 100%. The error bars represent standard deviations, and the asterisks indicate statistically significant measures (* $p < 0.01$; ** $p < 0.001$).

GsHE In Wound Repair

Wound healing is a natural restorative response to tissue injury, regulated by several proteins and secreted signals, which orchestrate the interactions among all the cells forming the skin. Recent studies have shown that hBD2, besides their roles as antimicrobial peptides and chemo attractants, play an active role in several growth-dependent processes as well, like wound healing and cell migration or proliferation.¹⁹⁻²¹ Furthermore, the increased expression of hBD2 can positively affect the wound healing process, which could be an additional advantage for the treatment of skin microlesions and scars linked to seborrhea.

To test the ability of GsHE to promote the wound healing process, a scratch assay was performed on fibroblasts treated with the extract. In a typical scratch assay, a “wound gap” in a cell monolayer is created by scratching, and the “healing” of this gap by cell migration towards the center of the gap is monitored and quantified. As showed in Figure 4(A,B), GsHE increased the capacity of the cells to repair the wound: it accelerated the cell migration and the healing process by 17% at the concentration of 0.002%, compared to the untreated control, similarly to the Transforming Growth Factor-β (TGF-β), known for its ability to accelerate the wound repair.²² Also in this case, there was

a slight difference between the two doses, confirming that the lower dose was the one giving a better positive effect on the cells.

the best results in most of the cell based assays. As control, the forehead remained untreated for the entire duration of the study and the formula used in the tests is reported in Table 1.




Figure 4: Scratch assay. Confluent fibroblast mono-layer cultures were scratched by a pipet tip and then treated with GsHE or the growth factor TGFβ. (A) Photographs of the cultures just after the scratching (T0), and after 7 h from the scratching (T7). The relative treatment is indicated below each picture. (B) The cell-free areas at T0 and T7 for each treatment were measured and compared, and the values expressed as percentages to the control, arbitrarily set as 100%. The error bars represent standard deviations, and the asterisks indicate statistically significant measures (* $p < 0.01$; ** $p < 0.001$).

GsHE Regulates Sebum Production *in Vivo*

On the basis of the results obtained from the *in vitro* studies, we employed the extract of *G. sulphuraria* in a clinical sebum normalizing efficacy test, performed on 20 male and female volunteers, aged from 18 to 50, having a greasy face skin. Twice a day, for 30 consecutive days, the volunteers applied a cream containing 0.002% of GsHE on the left side of the face and an emulsion without GsHE on the right side of the face. The concentration of 0.002% was chosen because it was the one giving

As showed in Figure 5, the mean cutaneous sebum amount was significantly decreased by GsHE treatment: the sebum excretion decreased by an average value of 58.5% after 15 days and 75.71% after 30 days of application with respect to the mean value measured at T0; the variations were extremely statistically significant ($p < 0.0001$). A significant decrease in the sebum level was observed also in the placebo treatment after 30 days, but it was more limited and corresponded to approximately half of that observed with GsHE application. The mean sebum value did not change in the forehead treatments (control).

This results confirm that GsHE can be an effective cosmetic ingredient, particularly suitable to relieve the problems associated with sebum overproduction in oily skins.

CONCLUSION

Research on microalgae has highlighted the huge applicative potentiality of these groups of microorganisms in industry: they are extremely diversified and phylogenetically distant one from the other, which makes them a natural inexhaustible source of unexplored compounds with the most various and divergent industrial applications, ranging from food, cosmetics, detergent to bio-fuel. Among the microalgae, the extremeophiles are even more attracting in terms of the applicative outputs of the molecules they synthesize: they have adapted to survive in extremely harsh environments, thus they represent highly effective biofactories of novel chemicals with unknown biological functions.

CAS	INCI Name	%INCI
7732-18-5	AQUA (WATER)	83.552
73398-61-5	CAPRYLIC/CAPRIC TRIGLYCERIDE	4.600
57-11-4	STEARIC ACID	3.300
31566-31-1	GLYCERYL STEARATE	2.400
9004-99-3	PEG-100 STEARATE	1.700
67762-27-0	CETEARYL ALCOHOL	1.050
36653-82-4	CETYL ALCOHOL	1.000
122-99-6	PHENOXYETHANOL	0.800
56-81-5	GLYCERIN	0.490
9004-99-3	PEG-20 STEARATE	0.350
70445-33-9	ETHYLHEXYGLYCERIN	0.300
76050-42-5	CARBOMER	0.260
139-33-3	DISODIUM EDTA	0.100
1310-73-2	SODIUM HYDROXIDE	0.096
68917-51-1	GsHE (ABSENT IN PLACEBO)	0.002
		100.000

Table 1: Cosmetic formula used in the clinical tests on sebum normalizing activity.




Figure 5: Clinical test. 20 panelists applied a cream without the active (placebo) and a cream containing GsHE (GsHE 0.002%), respectively on the right and left half face, twice a day for 28 consecutive days. After 15 days (T15) and 30 days (T30) of application, the sebum level on the skin was measured by the instrument Sebumeter® SM815, and compared with that measured in untreated skins (Control). The values were reported as micrograms of sebum per cm² of skin at the 3 different times of measure. Asterisks indicate significant differences (* $p < 0.01$; ** $p < 0.001$).

Thanks to their peculiarities, extremeophile microalgae have drawn the attention of researchers in the fields of Dermatology and Cosmetics, and some ingredients, derived from these species, have already found applications in the skin care market²³ and used by many top cosmetic brands. The species *Dunaliella salina*, living in high salinity environments, is the most used source of β -carotene, that is one of the leading antioxidant compounds employed in skin care formulas.²⁴ In addition, several types of extracts from *Arthrospira maxima*, commonly known as Spirulina, a cyanophyte living in alkaline waters, are claimed to repair the signs of early skin aging, exert a tightening effect and have wound healing properties, thanks to the presence of the protein C-phycoerythrin.^{25,26} Recently, an extract deriving from the snow micro alga *Chlamydocapsa* was also developed as product to fight extrinsic and intrinsic skin aging.²⁷

The extremeophilic red alga *Galdieria sulphuraria* lives in hot, sulfur-rich, acidic environments and shows an enormous metabolic flexibility, growing either photo autotrophically or heterotrophically on more than 50 carbon sources. It is also able to change its metabolism and its storage products according to the different growth conditions.²⁸ As extremophile, *Galdieria* is remarkably flexible because it has acquired a large number of genes from bacteria, and integrated them into a functional network.²⁹ Thanks to this evolutionary process, the species *Galdieria* has developed unique properties and adaptations due to the production of specific secondary metabolites. Some of these metabolites have been identified by a recent study which demonstrated the synthetic capacity of *Galdieria* to produce a class of compounds, the galactosylglycerols, specifically involved in the acclimation to various abiotic stresses, especially high salinity. Among them, the main photosynthetic products floridside and

isoforidoside are known also to contribute to the osmotic stress resistance of cells.³⁰

In the present study, we investigated the potential use of the microalga *G. sulphuraria* in skin care and, by performing both cellular assays *in vitro* and clinical tests, we demonstrated that the extract obtained from *G. sulphuraria* was able to reduce 5 α R1 expression in fibroblasts and enhance the skin's natural defense response by upregulating hBD2 expression in keratinocytes, resulting in a significant reduction of sebum production in the skin. Thanks to these activities, GsHE represents an attractive active ingredient for the skin care market, in particular as adjuvant in products designed to fight greasy skins, or even those affected by seborrheic dermatitis or acne. The extract was also capable of enhancing the healing process, as demonstrated by the cell culture tests, reinforcing the role of the extract as active ingredient in formulas addressed to alleviate the outbreak of lesions typical of acneic skins. Interestingly, these activities can be easily exploited in the hair care market as well, which is always looking for new and safe ingredients having a sebum reducing effect on the scalp, formulated either in conditioners or shampoos.^{31,32}

ACKNOWLEDGEMENT

We want to thank the professors A. Pollio and G. Pinto, University of Naples Federico II, Italy, for providing the microalga strain for the studies. The research was supported by the grants "Campus Bioframe" - POR CAMPANIA FESR2007 2013, and "Campania Bioscience" -PON03PE_00060_3.

CONFLICTS OF INTEREST: None.

REFERENCES

1. Jacobsen E, Billings JK, Frantz RA, Kinney CK, Stewart ME, Downing DT. Age-related changes in sebaceous wax ester secretion rates in men and women. *J Invest Dermatol.* 1985; 85(5): 483-485. doi: [10.1111/1523-1747.ep12277224](https://doi.org/10.1111/1523-1747.ep12277224)
2. Jenkins EP, Andersson S, Imperato-McGinley J, Wilson JD, Russell DW. Genetic and pharmacological evidence for more than one human steroid 5 alpha-reductase. *J Clin Invest.* 1992; 89(1): 293-300. Web site: <http://www.jci.org/articles/view/115574>. Accessed April 17, 2016
3. Thiboutot D, Harris G, Iles V, Cimis G, Gilliland K, Hagari S. Activity of the type 1 5-alpha-reductase exhibits regional differences in isolated sebaceous glands and whole skin. *J Invest Dermatol.* 1995; 105(2): 209-214. doi: [10.1111/1523-1747.ep12317162](https://doi.org/10.1111/1523-1747.ep12317162)
4. Burkhart CN, Gottwald L. Assessment of etiologic agents in acne pathogenesis. *Skin med.* 2003; 2(4): 222-228. doi: [10.1111/j.1540-9740.2003.02134.x](https://doi.org/10.1111/j.1540-9740.2003.02134.x)
5. Ro BI, Dawson TL. The role of sebaceous gland activity and scalp microfloral metabolism in the etiology of seborrheic dermatitis and dandruff. *J Invest Dermatol Symp Proc.* 2005; 10(3): 194-197. doi: [10.1111/j.1087-0024.2005.10104.x](https://doi.org/10.1111/j.1087-0024.2005.10104.x)
6. Yamasaki K, Gallo RL. Antimicrobial peptides in human skin disease. *Eur J Dermatol.* 2008; 18(1): 11-21. doi: [10.1684/ejd.2008.0304](https://doi.org/10.1684/ejd.2008.0304)
7. Raj PA, Dentino AR. Current status of defensins and their role in innate and adaptive immunity. *FEMS Microbiol Lett.* 2002; 206: 9-18. doi: [10.1111/j.1574-6968.2002.tb10979.x](https://doi.org/10.1111/j.1574-6968.2002.tb10979.x)
8. Chronnell CM, Ghali LR, Ali RS, et al. Human beta defensin-1 and -2 expression in human pilosebaceous units: upregulation in acne vulgaris lesions. *J Invest Dermatol.* 2001; 117(5): 1120-1125. doi: [10.1046/j.0022-202x.2001.01569.x](https://doi.org/10.1046/j.0022-202x.2001.01569.x)
9. Ott FD, Seckbach J. New classification for the genus Cyanidium Geitler 1933. In: Seckbach J. Ed., *Evolutionary Pathways and Enigmatic Algae: Cyanidium caldarium Rhodophyta and Related Cells*. Kluwer, Dordrecht: 1994; 145-152.
10. Merola A, Castaldo R, DeLuca P, Gambardella R, Musachio A, Taddei R. Revision of Cyanidium caldarium: three species of acidophilic algae. *G Bot Ital.* 1981; 115(4-5): 189-195. doi: [10.1080/11263508109428026](https://doi.org/10.1080/11263508109428026)
11. Carfagna S, Napolitano G, Barone D, Pinto G, Pollio A, Venditti P. Dietary supplementation with the microalga Galdieria sulphuraria (Rhodophyta) reduces prolonged exercise-induced oxidative stress in rat tissues. *Oxid Med Cell Longev.* 2015; Article ID 732090.
12. Gupta AK, Nicol KJ. The use of sulfur in dermatology. *Drugs Dermatol.* 2004; 3(4): 427-431. Web site: <http://europepmc.org/abstract/med/15303787>. Accessed April 17, 2016
13. Del Rosso JQ. The use of sodium sulfacetamide 10%-sulfur 5% emollient foam in the treatment of acne vulgaris. *J Clin Aesthet Dermatol.* 2009; 2(8): 26-29.
14. Graziani G, Schiavo S, Nicolai MA, et al. Microalgae as human food: chemical and nutritional characteristics of the thermo-acidophilic microalga Galdieria sulphuraria. *Food Funct.* 2013; 4(1): 144-152. doi: [10.1039/c2fo30198a](https://doi.org/10.1039/c2fo30198a)
15. Allen M. Studies with cyanidium caldarium, an anomalously pigmented chlorophyte. *Arch Mikrobiol.* 1959; 32(3): 270-277. doi: [10.1007/BF00409348](https://doi.org/10.1007/BF00409348)
16. Mosmann TJ. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Immunol Methods.* 1983; 65(1-2): 55-63. doi: [10.1016/0022-1759\(83\)90303-4](https://doi.org/10.1016/0022-1759(83)90303-4)
17. Crowther JM. Method for quantification of oils and sebum levels on skin using the Sebumeter®. *Int J Cosm Sci.* 2016; 38(2): 210-216. doi: [10.1111/ics.12258](https://doi.org/10.1111/ics.12258)
18. Habib FK, Ross M, Bayne CW, et al. The localisation and expression of 5 alpha-reductase types I and II mRNAs in human hyperplastic prostate and in prostate primary cultures. *J Endocrinol.* 1988; 156(3): 509-517. doi: [10.1677/joe.0.1560509](https://doi.org/10.1677/joe.0.1560509)
19. Niyonsaba F, Ushio H, Nakano N, et al. Antimicrobial peptides human beta-defensins stimulate epidermal keratinocyte migration, proliferation and production of proinflammatory cytokines and chemokines. *J Invest Dermatol.* 2007; 127(3): 594-604. doi: [10.1038/sj.jid.5700599](https://doi.org/10.1038/sj.jid.5700599)
20. Hirsch T, Spielmann M, Zuhaili B, et al. Human beta-defensin-3 promotes wound healing in infected diabetic wounds. *J Gene Med.* 2009; 11(3): 220-228. doi: [10.1002/jgm.1287](https://doi.org/10.1002/jgm.1287)
21. Baroni A, Donnarumma G, Paoletti I, et al. Antimicrobial human beta-defensin-2 stimulates migration, proliferation and tube formation of human umbilical vein endothelial cells. *Peptides.* 2009; 30(2): 267-272. doi: [10.1016/j.peptides.2008.11.001](https://doi.org/10.1016/j.peptides.2008.11.001)
22. Nall AV, Brownlee RE, Colvin CP, et al. Transforming growth factor beta 1 improves wound healing and random flap survival in normal and irradiated rats. *Arch Otolaryngol Head Neck Surg.* 1996; 122(2): 171-177. doi: [10.1001/archotol.1996.01890140057011](https://doi.org/10.1001/archotol.1996.01890140057011)
23. Wang HM, Chen CC, Huynh P, Chang JS. Exploring the potential of using algae in cosmetics. *Bioresour Technol.* 2015; 184: 355-362. doi: [10.1016/j.biortech.2014.12.001](https://doi.org/10.1016/j.biortech.2014.12.001)

24. Ye ZW, Jiang JG, Wu GH. Biosynthesis and regulation of carotenoids in *Dunaliella*: progresses and prospects. *Biotechnol Adv.* 2008; 26(4): 352-360. doi: [10.1016/j.biotechadv.2008.03.004](https://doi.org/10.1016/j.biotechadv.2008.03.004)

25. Stolz P, Obermayer B. Manufacturing microalgae for skin care. *Cosmetics & Toiletries.* 2005; 120(3): 99-106.

26. Madhyastha H, Madhyastha R, Nakajima Y, Omura S, Maruyama M. Regulation of growth factors-associated cell migration by C-phycoyanin scaffold in dermal wound healing. *Clin Exp Pharmacol Physiol.* 2012; 39(1): 13-19. doi: [10.1111/j.1440-1681.2011.05627.x](https://doi.org/10.1111/j.1440-1681.2011.05627.x)

27. SchürchStutz C, Schmid D, Züllli F. US Patent 20100316720 A1. Use of an extract from snow algae in cosmetic or dermatological formulations. 2010.

28. Sakurai T, Aoki M, Ju X, et al. Profiling of lipid and glycogen accumulations under different growth conditions in the sulfotermophilic red alga *Galdieria sulphuraria*. *Bioresour Technol.* 2016; 200: 861-866. doi: [10.1016/j.biortech.2015.11.014](https://doi.org/10.1016/j.biortech.2015.11.014)

29. Schönknecht G, Chen WH, Ternes CM, et al. Gene transfer from bacteria and archaea facilitate devolution of an extremophilic eukaryote. *Science.* 2013; 339(6124): 1207-1210. doi: [10.1126/science.1231707](https://doi.org/10.1126/science.1231707)

30. Pade N, Linka N, Ruth W, Weber AP, Hagemann M. Floridoside and isofloridoside are synthesized by trehalose 6-phosphate synthase-like enzymes in the red alga *Galdieria sulphuraria*. *New Phytol.* 2015; 205(3): 1227-1238. doi: [10.1111/nph.13108](https://doi.org/10.1111/nph.13108)

31. Kim YR, Kim JH, Shin HJ, Choe YB, Ahn KJ, Lee YW. Clinical evaluation of a new-formula shampoo for scalp seborrheic dermatitis containing extract of *Rosa centifolia* petals and epigallo catechingallate: a randomized, double-blind, controlled study. *Ann Dermatol.* 2014; 26(6): 733-738. doi: [10.5021/ad.2014.26.6.733](https://doi.org/10.5021/ad.2014.26.6.733)

32. Draelos ZD. Essentials of hair care often neglected: hair cleansing. *Int J Trichology.* 2010; 2(1): 24-29. doi: [10.4103/0974-7753.66909](https://doi.org/10.4103/0974-7753.66909)