

Observational Study

*Corresponding author

Huseyin Sari, MD
Department of Otolaryngology
Okmeydani Training and Research
Hospital, Istanbul, Turkey
E-mail: drhusyinsari@gmail.com

Volume 3 : Issue 3

Article Ref. #: 1000OTLOJ3143

Article History

Received: September 14th, 2017

Accepted: October 31st, 2017

Published: November 1st, 2017

Citation

Sari H, Yıldırım Y, Deniz S, et al. Importance of keros classification in surgery. *Otolaryngol Open J.* 2017; 3(3): 54-58. doi: [10.17140/OTLOJ-3-143](https://doi.org/10.17140/OTLOJ-3-143)

Copyright

©2017 Sari H. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Importance of Keros Classification in Surgery

Hüseyin Sari, MD^{1*}; Güven Yıldırım, MD²; Suzan Deniz Önoel, MD³; Erdi Özdemir, MD¹; Ahmet Arslanoglu, MD¹; Berk Gurpınar, MD¹; Tolgar Lütü Kumral, MD¹; Yavuz Uyar, MD¹; Ziya Salturk, MD¹

¹Department of Otolaryngology, Okmeydani Training and Research Hospital ENT Clinic, Istanbul, Turkey

²Department of Otolaryngology, Giresun University ENT Clinic, Giresun, Turkey

³Department of Otolaryngology, Okmeydani Training and Research Hospital Radiology Clinic, Istanbul, Turkey

ABSTRACT

Objective: The relationship between the cribriform plate and the uncinat process may be elaborated with the help of the Keros classification. The observations were analysed using high-resolution computed tomography (HRCT). Additionally, the relationship between the superior attachment of the uncinat process, the existence of concha bullosa, and the different types of ethmoid roofs were examined.

Methods: Five-hundred and sixteen subjects complaining of sinonasal disorders between 2015 and 2016 were enrolled retrospectively at the Okmeydani Training and Research Hospital, Department of Otolaryngology. HRCT scans of 1-mm sections were obtained. Keros classification was used for the measurement of the depth of the olfactory fossa as follows: Keros I: 1-3 mm, Keros II: 3-7 mm, and Keros III: 7-16 mm.

Results: Fifty-one point nine percentage of cases were of Keros type II, 27.7% of type III, and 20.3% of type I. In 82.4% of the reported cases, the uncinat process was attached to the lamina papyracea, in 11% of the cases it was attached to the middle nasal turbinate, and in 6.4% cases to the skull base. A concha bullosa was observed in the right nasal cavity in 13.8% of the cases, in the left nasal cavity in 11% of the cases, and bilaterally in 16.3% of the cases. Variances observed in the Keros types were not statistically significant with respect to sex ($p>0.05$). Concha bullosa was significantly more frequently seen in females (53.1%) relative to males ($p=0.001$).

Conclusion: Endoscopic sinus surgery (ESS) is the primary mode of treatment for chronic sinus diseases. In this procedure, a proper assessment of the vital structures is very important to avoid further complications. In the present study, it has been suggested that determining the depth of the ethmoid roof is necessary to avoid injuring the bony lamella, which can lead to cerebrospinal fluid rhinorrhea.

KEY WORDS: Keros classification; Ethmoid roof; Paranasal Sinus; Cribriform plate.

ABBREVIATIONS: HRCT: High-resolution computed tomography; ESS: Endoscopic Sinus Surgery.

INTRODUCTION

Chronic sinonasal diseases are among the most common diseases demanding surgical attention in the field of otolaryngology.¹ Although, some diseases can be treated with medicines; surgical intervention is more often needed. Endoscopic sinus surgery (ESS) is the most commonly implemented approach for treatment at present.^{2,3} Thorough anatomical knowledge is considered mandatory, especially with respect to the skull base, to prevent serious surgical complications. Therefore, variations in anatomy should be diagnosed before surgery and examined carefully for every patient. Keros classification is a widely used procedure for the evaluation of the depth

of the nasal roof and should be assessed pre-operatively; which may otherwise lead to many life-threatening complications.

The uncinata process and the attachment of the middle turbinate can be associated with several anatomical regions. Rarely, those structures may attach directly to the nasal roof. Interfering with this anatomical location can cause serious clinical complications including cerebrospinal rhinorrhea, meningitis, etc. High-resolution computed tomography (HRCT) is the gold standard for diagnosis in the paranasal sinuses and neighboring structures and is a vital tool for pre-operative assessment to prevent related medical complications.⁴

In this study, the relationship between the cribriform plate and the uncinata process was investigated using the Keros classification. The relationship between the superior attachment of the uncinata process, the existence of concha bullosa, and the different types of ethmoid roofs were also examined carefully.

MATERIALS AND METHODS

Institutional review board approval was obtained from the Okmeydani Training and Research Hospital Ethical Committee. HRCT of the paranasal sinuses of 516 subjects who had sinonasal disorders in 2015-2016 were included in the study. Exclusion criteria were a previous history of nasal surgery, maxillofacial trauma, benign or malignant tumors of the sinonasal tract, and cases with asymmetrical morphology of the olfactory fossa. The pediatric population was not included in the study.

HRCT (Philipps 128 slice, Holland) scans of 1-mm sections were obtained. The Keros classification is used for measuring the depth of the olfactory fossa and defined as follows: Keros I: 1-3 mm, Keros II: 3-7 mm, and Keros III: 7-16 mm (Figures 1 and 2).⁵ In all the scans, the attachment of the uncinata process was also observed and grouped as: type 1 attached to the lamina papyracea, type 2 to the skull base, and type 3 to the middle turbinate. Concha bullosa cases were also identified. Concordance between the Keros types and the other groups was observed.




Figure 1: A Schematic Representation of Keros Types.

Statistical analysis was performed using SPSS software version 22 (IBM, USA). Continuous data has been reported as mean±standard deviation. Statistical significance was accepted when $p < 0.05$. The chi-squared test was used to determine the statistically significant variances.

RESULTS

The average age of the participants in the study was 34.37 ± 12.03 years (Table 1). Fifty-one point nine percentage of cases were associated with Keros type II, 27.7% with type III, and 20.3% with type I. In 82.4% of cases, the uncinata process was attached to the lamina papyracea, in 11% cases to the middle nasal turbinate, and in 6.4% cases to the skull base. Concha bullosa was in the right nasal cavity in 13.8% of cases, in the left nasal cavity in 11% of the cases, and bilateral in 16.3% of cases (Table 2, Figure 3). On the basis of the Chi-square test, differences in Keros types were not statistically significant as was observed with respect to sex ($p > 0.05$) and Concha bullosa (53.1%) was significantly higher in females relative to males ($p = 0.001$).




Figure 2: Computerized Tomography Images of Keros Types.

Table 1: Patient Demographics.

		n	%
Age	14-29	213	41.3
	30-39	130	25.2
	40 and above	173	33.5
Sex	Male	356	69
	Female	160	31
Keros type	Keros 1	105	20.3
	Keros 2	268	51.9
	Keros 3	143	27.7
Unsinat process attachment	L.Papyracea	425	82.4
	Middle Turbinate	57	11
	Skull Base	33	6.4
Concha Bullosa	None	304	58.9
	Right	71	13.8
	Left	57	11.0
	Bilateral	84	16.3

Table 2: Keros Types and Concha Bullosa According to Sex.

		Sex		p
		Male	Female	
		n (%)	n (%)	
Keros type	Keros 1	74 (%20.8)	31 (%19.4)	0.759
	Keros 2	181 (%50.8)	87 (%54.4)	
	Keros 3	101 (%28.4)	42 (%26.3)	
Concha Bullosa	None	229 (%64.3)	75 (%46.9)	0.001*
	Right	39 (%11.0)	32 (%20.0)	
	Left	32 (%9.0)	25 (%15.6)	
	Bilateral	56 (%15.7)	28 (%17.5)	

Chi-Square test *p<0.05




Table 3: Keros Types, Uncinate Process Attachments and Concha Bullosa.

		Keros Type			P
		Keros 1	Keros 2	Keros 3	
		n (%)	n (%)	n (%)	
Unsinate process attachment	L. Papyraceae (Type 1)	83 (%79)	230 (%85.8)	112 (%78.3)	0.100
	Middle Turb. (Type 2)	14 (%13.3)	32 (%11.9)	11 (%7.7)	0.299
	Skull base (Type 3)	8 (%7.6)	7 (%2.6)	18 (%12.6)	0.001*
Concha bullosa	None	57 (%54.3)	150 (%56)	97 (%67.8)	0.297
	Right	17 (%16.2)	39 (%14.6)	15 (%10.5)	
	Left	14 (%13.3)	32 (%11.9)	11 (%7.7)	
	Bilateral	17 (%16.2)	47 (%17.5)	20 (%14)	

Among the Keros types, having performed chi-square analysis, there was no significant relationship between the attachment of the uncinate process to the concha bullosa ($p>0.05$). However, among the Keros types with the uncinate process attached to the skull base, statistical significance was observed ($p=0.001$). There were significant differences in the ratio of the skull base attachment between Keros types 1 and 2 ($p=0.038$), and between types 2 and 3 ($p=0.001$) (Table 3) The ratio of the skull base attachment in Keros type 2 (2.6%) was significantly lower than that of types 1 (7.6%) and 3 (12.6%). There was no significant difference between the Keros types and the prevalence of concha bullosa ($p>0.05$). The analysis of the recorded data was performed using Chi-square test.

DISCUSSION

Our data revealed that Keros type 2 was the most commonly observed type in medical conditions and lamina papyrycea was the most common site of insertion of uncinate process but the variations were not rare. Especially, Keros type 3 with skull base attachment was a potential dangerous clinical variation. The sinonasal tract can be visualized with the naked eye or by using nasal endoscopy. If clinically suspected, diseases or variations can be examined using the HRCT evaluation, which showed the anatomical structures and their relationships to each other, anatomical variations, and pathologies.⁶⁻⁸

Although, anatomical knowledge and experience reduces the chances of arising complications, ESS has been associated with some problematic symptoms; minor complications including bleeding, infection, crusting, synechiae, ostial stenosis, and recurrence of the disease;⁹ and major complications including extraocular muscle damage, cerebrospinal fluid rhinorrhea, intracranial damage, and orbital emphysema.¹⁰

Pre-operative assessment may help to prevent cerebrospinal fluid rhinorrhea. If the olfactory fossa is deep, as in Keros III, the surgeon must take care not to approach medially during the surgery, as a thin bony lamella constitutes most of the roof and the frontal support is missing. These patients are prone to complications associated with the ethmoid roof. However, in Keros I, surgery may be safer to perform.⁵⁻¹¹

Uncinectomy is the primary step of ESS. The attachment of the uncinate process can vary and should be analyzed before surgery. Stamberger et al¹² reported three different types of attachments. The uncinate process was attached to the middle turbinate in the first type, to the frontal roof in the second, and to the lamina papyraceae in the third. These variations of the uncinate process alter surgery in terms of finding the frontal recess. If not examined properly, damage to the skull base and cerebrospinal fluid rhinorrhea may possibly result.¹³

Pneumatized middle concha is another frequently observed variation in the sinonasal region, with a rate of incidence of 13-73%,¹⁴⁻¹⁵ subsequently exerting pressure on the middle concha and ethmoid labyrinth.^{16,17} An uncinate process may also come in contact with a pneumatized middle concha, which may require surgical intervention.

Our data revealed that Keros type 2 was the most common type, followed by type 1, which was consistent with the results of Guldner et al.¹⁸ However, Elwany et al¹⁹ reported that Keros type 3 was the most common one to be observed. Our study indicated that gender was not a significant factor for influencing variations, which was consistent with the conclusions of Elwany et al.¹⁹

Studies examining the point of attachment of the uncinate process were consistent with the recorded results.²⁰⁻²² The only exception was that Keros type 3 were observed to be more common than type 2, which may have been so, on account of the limited number of participants in the study.

ESS is a delicate surgery that is associated with major complications. The only approach to avoid those complications is to conduct a meticulous assessment of each patient before surgery. This evaluation is only possible with HRCT, and there are some key points to differentiate the danger areas of the sinonasal tract prior to ESS.

CONCLUSION

Our study suggested the determination of the depth of the ethmoid roof to avoid injuring the bony lamella and thus reduced

the risks of developing cerebrospinal fluid rhinorrhea.

ACKNOWLEDGMENT

The authors report nothing to declare.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

REFERENCES

1. Zacharek MA, Han JK, Allen R, et al. Sagittal and coronal dimensions of the ethmoid roof: A radioanatomic study. *Am J Rhinol.* 2005; 19: 348-352.
2. Luong A, Marple BF. Sinus surgery: Indications and techniques. *Clin Rev Allergy Immunol.* 2006; 30: 217-222.
3. Ohnishi T, Tachibana T, Kaneko Y, Esaki S. High-risk areas in endoscopic sinus surgery and prevention of complications. *Laryngoscope.* 1993; 103: 1181-1185. doi: [10.1288/00005537-199310000-00020](https://doi.org/10.1288/00005537-199310000-00020)
4. Leunig A, Betz CS, Sommer B, Sommer F. Anatomic variations of the sinuses; multiplanar CT-analysis in 641 patients. *Laryngorhinootologie.* 2008; 87: 482-489. doi: [10.1055/s-2007-995572](https://doi.org/10.1055/s-2007-995572)
5. Keros P. On the practical value of differences in the level of the lamina cribrosa of the ethmoid. *Z Laryngol Rhinol Otol.* 1962; 41: 809-813.
6. Zeinreich SJ. Paranasal sinus imaging. *Otolaryngol Head Neck Surg.* 1990; 103: 868-869. doi: [10.1177/01945998901030S505](https://doi.org/10.1177/01945998901030S505)
7. Messerklinger W. On the drainage of the frontal sinus of man. *Acta Otolaryngol.* 1967; 63: 176-181. doi: [10.3109/00016486709128748](https://doi.org/10.3109/00016486709128748)
8. Stammberger H. Endoscopic endonasal surgery-concepts in treatment of recurring rhinosinusitis. Part 1. Anatomik and pathophysiologic considerations. *Otolaryngol Head Neck Surg.* 1986; 94(2): 143-147. doi: [10.1177/019459988609400202](https://doi.org/10.1177/019459988609400202)
9. McMains KC. Safety in endoscopic sinus surgery. *Curr Opin Otolaryngol Head Neck Surg.* 2008; 16: 247-251. doi: [10.1097/MOO.0b013e3282fdccad](https://doi.org/10.1097/MOO.0b013e3282fdccad)
10. Ulualp SO. Complications of endoscopic sinus surgery: Appropriate management of complications. *Curr Opin Otolaryngol Head Neck Surg.* 2008; 16: 252-259. doi: [10.1097/MOO.0b013e3282fdc3b2](https://doi.org/10.1097/MOO.0b013e3282fdc3b2)
11. Erdogan S, Keskin G, Topdag M, Ozturk M, Sari F, Mutlu F. Ethmoid roof radiology; analysis of lateral lamel-

- la of cribriform plate. *Otolaryngol Pol.* 2015; 69: 53-57. doi: [10.5604/00306657.1184543](https://doi.org/10.5604/00306657.1184543)
12. Stammberger H, Posawetz W. Functional endoscopic sinus surgery. Concept, indications and results of Messerklinger technique. *Eur Arch Otorhinolaryngol.* 1990; 247(2): 63-76.
13. Sheng YC, Chih JY, Chiao HL, et al. The association of superior attachment of uncinate process with pneumatization of middle turbinate: A computed tomographic analysis. *Eur Arch Otorhinolaryngol.* 2017; 274: 1905-1910. doi: [10.1007/s00405-016-4441-3](https://doi.org/10.1007/s00405-016-4441-3)
14. Bolger WE, Butzin CA, Parsons DS. Paranasal sinus bony anatomic variations and mucosal abnormalities: CT analysis for endoscopic sinus surgery. *Laryngoscope.* 1991; 101: 56-64. doi: [10.1288/00005537-199101000-00010](https://doi.org/10.1288/00005537-199101000-00010)
15. Perez-Piñas I, Sabaté J, Carmona A, Catalina-Herrera CJ, Jiménez-Castellanos J. Anatomical variations in the human paranasal sinus region studied by CT. *J Anat.* 2000; 197: 221-227. doi: [10.1046/j.1469-7580.2000.19720221.x](https://doi.org/10.1046/j.1469-7580.2000.19720221.x)
16. Unlu HH, Akyar S, Caylan R, Nalça Y. Concha bullosa. *J Otolaryngol.* 1994; 23: 23-27.
17. Maru YK, Gupta Y. Concha bullosa: Frequency and appearances on sinonasal CT. *Indian J Otolaryngol Head Neck Surg.* 1999; 52: 40-44. doi: [10.1007/BF02996431](https://doi.org/10.1007/BF02996431)
18. Guldner C, Zimmermann AP, Diogo J, Werner JA, Teymoortash A. Age-dependent differences of the anterior skull base. *Int J Pediatr Otorhinolaryngol.* 2012; 76(6): 822-828. doi: [10.1016/j.ijporl.2012.02.050](https://doi.org/10.1016/j.ijporl.2012.02.050)
19. Elwany S, Medanni A, Eid M, Aly A, El-Daly A, Ammar SR. Radiological observations on the olfactory fossa and ethmoid roof. *J Laryngol Otol.* 2010; 124: 1251-1256. doi: [10.1017/S0022215110001313](https://doi.org/10.1017/S0022215110001313)
20. Ercan I, Cakir BO, Sayin I, Başak M, Turgut S. Relationship between the superior attachment type of uncinate process and presence of agger nasi cell: A computer-assisted anatomic study. *Otolaryngol Head Neck Surg.* 2006; 134: 1010-1014. doi: [10.1016/j.otohns.2006.01.021](https://doi.org/10.1016/j.otohns.2006.01.021)
21. Srivastava M, Tyag M. Role of Anatomic variations of uncinate process in frontal sinusitis. *Indian J Otolaryngol Head Neck Surg.* 2016; 68(4): 441-444. doi: [10.1007/s12070-015-0932-6](https://doi.org/10.1007/s12070-015-0932-6)
22. Tuli IP, Sengupta S, Munjal S, Kesari SP, Chakraborty S. Anatomical variations of uncinate process observed in chronic sinusitis. *Indian J Otolaryngol Head Neck Surg.* 2013; 65(2): 157-161. doi: [10.1007/s12070-012-0612-8](https://doi.org/10.1007/s12070-012-0612-8)