

Research

*Corresponding author

Cristina Palacios, PhD
Nutrition Program
Graduate School of Public Health
PO Box 365067, San Juan
00936-5067, Puerto Rico
Tel. (787) 758-2525
Fax: (787) 759-6719
E-mail: cristina.palacios@upr.edu

Volume 1 : Issue 1
Article Ref. #: 1000DROJ1102

Article History

Received: October 25th, 2014
Accepted: January 20th, 2015
Published: January 21st, 2015

Citation

Ramos-Trautmann G, González L, Díaz-Luquis G, M. Pérez C, Palacios C. Inverse association between vitamin D status and diabetes in a clinic based sample of Hispanic adults in Puerto Rico. *Diabetes Res Open J*. 2015; 1(1): 5-11. doi: [10.17140/DROJ-1-102](https://doi.org/10.17140/DROJ-1-102)

Copyright

©2014 Palacios C. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Inverse Association between Vitamin D Status and Diabetes in a Clinic Based Sample of Hispanic Adults in Puerto Rico

Grisel Ramos-Trautmann¹, Lilliana González¹, Giselle Díaz-Luquis¹, Cynthia M. Pérez² and Cristina Palacios^{1*}

¹Nutrition Program, Graduate School of Public Health, Medical Sciences Campus, University of Puerto Rico, San Juan, Puerto Rico

²Epidemiology Program, Graduate School of Public Health, Medical Sciences Campus, University of Puerto Rico, San Juan, Puerto Rico

ABSTRACT

Background: Vitamin D deficiency is a public health problem around the world. Diabetes has been associated with vitamin D deficiency. We aimed to examine the association between the vitamin D status and diabetes in a clinic based sample of Hispanic adults in Puerto Rico.

Methods: Demographics and laboratory test results for serum 25(OH)D, Fasting Blood Glucose (FBG), and Haemoglobin A1C (HbA1c) were extracted from medical records. Vitamin D status was classified as deficient (<12 ng/ml); inadequate (12-20 ng/ml); insufficient (21-29 ng/ml) and optimal (≥ 30 ng/ml) using serum 25(OH)D levels.

Results: A total of 716 records were included in the analyses. Most were females (63.3%), with mean age of 54.1 \pm 14.9 y, mean BMI 30.1 \pm 6.3 kg/m² and mean serum 25(OH)D levels of 24.3 \pm 8.6 ng/ml. Most were classified as diabetics (41.1%). Those with diabetes had lower 25(OH)D levels compared to pre-diabetic and normal glucose status ($p < 0.05$). Serum 25(OH)D levels were inversely correlated to FBG and HbA1c in the total sample and in men ($p < 0.05$). After adjusting for age, gender, BMI and seasonality, there was a greater risk of diabetes, but not prediabetes, in those with serum 25(OH)D levels <30 ng/ml. This risk increased from 1.8 times in those with vitamin D insufficiency to 4.2 times in those with vitamin D deficiency (<12 ng/ml).

Conclusion: Diabetes risk significantly increased as serum 25(OH)D levels decreased in this group of Hispanic adults, underscoring the importance of routinely screening high risk individuals for vitamin D deficiency and offer supplementation to normalize serum levels.

KEYWORDS: Vitamin D status; 25(OH)D levels; Diabetes; Glucose parameters.

INTRODUCTION

Diabetes affects 25.8 million people of all ages in the United States (US), which represents 8.3% of the US population.¹ This is higher in Hispanics as evidenced by data from the Hispanic Community Health Study/Study of Latinos (HCHS/SOL), a population-based cohort study of Hispanic and Latino adults from diverse backgrounds, in which a diabetes prevalence of 16.7% in men and 17.2% in women was reported.^{2,3} Among the Hispanic groups studied in HCHS/SOL, Puerto Ricans had the highest prevalence of diabetes, particularly among women. In addition, results from the Behavioural Risk Factor Surveillance System in Puerto Rico (PR) also showed that the estimated prevalence of self-reported diabetes was >12%,⁴ which is among the highest compared to other states and territories of the US. Furthermore, a study in a representative sample of adults residing in the San Juan Metropolitan Area of PR showed that

the prevalence of diabetes was even higher, with an age-standardized prevalence of pre-diabetes and diabetes, detected by Fasting Plasma Glucose (FPG) and/or HbA1c, as 25.5% and 47.4%, respectively.⁵ Diabetes is considered one of the leading causes of death in US and PR,⁶ therefore, this is an important health disparity to address among Puerto Ricans.

Growing evidence suggests that diabetes risk is associated with vitamin D status. A recent meta-analysis of 21 prospective studies with a total of 76,220 participants (mostly non-Hispanic Whites) with about 5,000 incident cases of diabetes found a relative risk for diabetes of 0.62 (95% CI: 0.54-0.70) when comparing the highest with the lowest levels of 25(OH) D.⁷ For each 10 nmol/L increment in serum 25(OH)D levels, there was a reduction in the risk of diabetes of 4% ($p < 0.01$).⁷ In addition, when 25(OH)D levels were near 50 nmol/l, the risk of diabetes was significantly reduced. Another meta-analysis of 19 cross-sectional, 13 case-control, and 12 prospective studies suggested that low serum 25(OH)D levels is associated with glucose intolerance, β cell dysfunction, and insulin resistance.⁸

Vitamin D deficiency is a public health concern, with a high prevalence worldwide.⁹ In the US, 32% of the population has vitamin D levels below 20 ng/mL.¹⁰ There is also a health disparity in vitamin D status. Hispanics are at high risk of low vitamin D status.¹⁰⁻¹⁵ Among 358 Hispanic-American men, the highest prevalence of vitamin D deficiency was found among those born in PR (26%), compared to those from the Dominican Republic (21%), Central America (11%), and South America (9%).¹⁶ Recently, we found in a large sample of 2,293 adults in PR that 72% had vitamin D insufficiency (levels below 30 ng/ml) and 28% were vitamin D deficient.¹⁷

There is scarce information assessing the association between vitamin D status and glycemic in Hispanics. This is important to study as Hispanics have a higher prevalence of both diabetes and low vitamin D status, particularly Puerto Ricans. Therefore, to start addressing this gap in knowledge, we assessed whether there was an association between vitamin D and glycemic status in a clinic-based sample in Hispanic adults living in PR.

METHODS

The study was based on a retrospective, secondary analysis of medical records of patients attending the "Endocrinology, Diabetes and Metabolism Clinic" located in the municipality of Utuado in PR.

Participants

The study population consisted of non institutionalized individuals who attended this clinic for the first time between 2005 and 2013. A convenience sample was selected from those

records meeting the following inclusion criteria: aged 21 years and older; having test results for Fasting Blood Glucose (FBG), Oral Glucose Tolerance Test (OGTT), and/or Haemoglobin A1c (HbA1c); and serum 25(OH)D levels taken within 3 weeks of each other. Patients using any type of vitamin D supplementation regimens were excluded from the study, as there was a large variation in the doses reported (from 400 to 10,000 IU/d).

Data Collection

A data collection sheet was used to extract the information from medical records, which included data on demographics, anthropometrics, blood tests results for 25(OH)D levels and glucose metabolism markers, and prescribed hypoglycemic agents, which could potentially influence vitamin D metabolism.

Demographics and other health measures

Gender and age, calculated in years at the time of the medical visit, were recorded from the medical records. Prescribed hypoglycemic agents were recorded as physician's notes with names and doses used.

Anthropometrics

Body weight (pounds) and height (inches) were measured once by the clinic staff using a physician scale (Detecto, Model 338, MO, USA) while participants were wearing street clothes but no shoes. These measurements were converted to kg and meters. Body Mass Index (BMI) was calculated as the ratio of weight (kilograms) to the square of height (meters).

Blood tests

Laboratory test results dates were recorded from the medical records as month/day/year. All blood test results were taken after an overnight fast. Results from serum 25(OH)D levels were recorded in nanograms per millilitre (ng/mL) and these were analyzed by two methods. The main method used by 97% of the samples was immunoassay and only 3% were performed with Liquid Chromatography mass spectrometry (LC-MS/MS). Due to methodological differences between these methods,¹⁸ only blood test results analyzed by immunoassay were included. Serum 25(OH)D levels were measured by the local laboratory using a commercially available enzyme immunoassay kit (IDS 25-Hydroxy Vitamin D EIA kit). The blood tests were collected all year round and the date of the serum 25(OH)D was recorded to adjusted for seasonality. Since the half-life of 25(OH)D levels is 2-3 weeks,¹⁸ only blood tests results measured within this period were included in the analyses.

Although there is no universally accepted cut-offs for serum 25(OH)D levels, the Institute of Medicine (IOM)

established that serum 25(OH)D levels above 20 ng/ml (50 nmol/l) are optimal and sufficient to maintain bone health.¹⁹ However, the Endocrinology Society defines vitamin D deficiency as 25(OH)D levels below 20 ng/ml and adds a category for vitamin D insufficiency (levels 21-29 ng/ml; 52-72 nmol/l).²⁰ Furthermore, the Endocrinology Society also states that levels above 30 ng/ml (75 nmol/l) provide increased overall health benefits, such as reducing the risk of diabetes. Therefore, for the present study, vitamin D status was classified as: deficient if serum 25(OH)D levels were <12 ng/ml (30 nmol/l); inadequate if levels were 12-20 ng/ml (30-50 mmol/l); insufficient if levels were 21-29 ng/ml and optimal as levels ≥30 ng/ml.

The markers for glucose metabolism recorded were Fasting Blood Glucose (FBG), HbA1c, and 2-hour Plasma Glucose (2hPG). These were measured routinely by local laboratories; glucose levels were measured by Spectrophotometry and HbA1c by HPLC. Individuals were classified as having diabetes by using the following criteria: HbA1c ≥6.5%; FBG ≥126 mg/dl (≥7.0 mmol/l); and/or 2hPG ≥200 mg/dl (≥11.1 mmol/l)^{21,22} or if use of hypoglycemic agents was documented in the medical record.²³ The following criteria were used to classify pre diabetes: HbA1c 5.7-6.4%; FBG 100-125 mg/dl; or a 2hPG ≥140 (≥7.8 mmol/l) and <200 mg/dl (<11.1 mmol/l).²¹ Normal glucose status was defined as HbA1c <5.7%; FBG <100 mg/dl; and/or 2Hpg of <140 mg/dl (<7.8 mmol/l).²¹

This study was approved by the Institutional Review Board of the Medical Sciences Campus (MSC) of the University of Puerto Rico (UPR). In addition, the clinic director provided written authorization to access all medical records. To assure

confidentiality, all data collection sheets were stored in a locked room in the Nutrition Program of the MSC UPR Graduate School of Public Health.

STATISTICAL ANALYSIS

Baseline characteristics were summarized using mean and Standard Deviation (SD) for continuous variables and frequency distributions for categorical variables. Analysis of Covariance (ANCOVA) was used to assess differences in age-adjusted mean serum 25(OH)D levels across glycemic status (normal, pre diabetes, and diabetes). Bonferroni’s posthoc test was used for multiple comparisons. Pearson’s correlation coefficient was used to assess the correlation of serum 25(OH)D with glucose parameters. A multinomial logistic regression model was used to assess the association between vitamin D and glycemic status after adjustment for age, gender, BMI, and season. All analyses were performed using SPSS Statistical Package (SPSS version 21.0 for Windows, 2012, SPSS Inc., Chicago, IL.). Statistical significance was set at p<0.05.

RESULTS

From 1,379 consecutive medical records reviewed, 716 (52%) records met the inclusion criteria. A total of 89 records reported the use of vitamin D supplements and 574 did not have the levels of serum 25(OH)D or any of the glucose measures reported or were not done within 3 weeks of each other; therefore, these were excluded from the analyses. The descriptive characteristics of the study sample are presented in table 1.

Characteristic	Total			Males			Females		
	n	mean	SD	n	mean	SD	n	mean	SD
Age (y)	716	54.1 ± 14.9		263	53.8 ± 13.9		453	54.3 ± 15.4	
BMI (kg/m ²)	693	30.1 ± 6.3		255	30.4 ± 5.8		438	29.8 ± 6.5	
25(OH)D (ng/ml)	716	24.3 ± 8.6		263	25.5 ± 8.4		453	23.6 ± 8.6*	
Vitamin D Status (%)									
Deficient (<12 ng/ml)	39	5.4		10	3.8		29	6.4	
Inadequate (12-20 ng/ml)	192	26.8		63	24.0		129	28.5	
Insufficiency (21-29 ng/ml)	319	44.6		120	45.6		199	43.9	
Optimal (≥30 ng/ml)	166	23.2		70	26.6		96	21.2	
FBG (mg/dl)	697	122.8 ± 57.7		259	132.8 ± 64.6		438	117.0 ± 52.4*	
2h OGTT (mg/dl)	280	141.3 ± 68.2		100	154.1 ± 77.5		180	134.2 ± 61.5*	
HbA1c	547	6.9 ± 2.0		218	7.2 ± 2.2		329	6.8 ± 1.8*	
Glycemic Status (%)									
Normal	216	30.2		60	22.8		156	34.4	
Pre diabetes	201	28.1		77	29.3		124	27.4	
Diabetes	294	41.1		125	47.5		169	37.3	
Use of hypoglycemic medications									
No	278	38.8		115	43.7		163	36.0	
Yes	438	61.2		148	56.3		290	64.0	

*p<0.05 compared to males by t test. † Age-adjusted means.

Table 1: Descriptive statistics of the sample (Mean ± SD† or %)

Most participants were females (63.3%) and mean age was 54.1±14.9 years. Most patients were overweight (38.8%) or obese (46.3%), had suboptimal vitamin D status (76.8%), had undiagnosed diabetes according to laboratory criteria (55.8%), and 39.1% were taking hypoglycaemic medications. Serum 25(OH)D levels and FBG were significantly higher in men compared to women (p<0.05).

Figure 1 shows the difference in mean 25(OH)D levels by glucose status. Overall, individuals classified as normal glucose status had higher 25(OH)D levels than those with diabetes (p<0.001), whereas subjects with pre-diabetes had higher 25(OH)D levels than those with diabetes (p<0.01). No significant difference was observed between subjects with normal glucose and those with pre diabetes (p=0.78). When analyzed

separately by gender, those with normal glucose status had higher 25(OH)D levels than those with diabetes (p<0.001), but no significant differences were found in vitamin D levels between those with pre-diabetes and those with diabetes).

Correlations between 25(OH)D levels and glyceimic parameters are shown in table 2. There was a significant inverse correlation between 25(OH)D levels and FBG (p<0.001), whereas participants with lower 25(OH)D levels had higher FBS levels in the total sample and in males. Similarly, there was a significant inverse correlation between 25(OH)D levels and HbA1c levels overall and in men (p<0.001), whereas those with lower 25(OH)D levels had with higher HbA1c levels. No correlation was seen with 2h-OGTT.


Glyceimic Parameters	Total	Male	Female
FBG	- 0.10*	- 0.18*	- 0.09
2h OGTT	- 0.01	- 0.04	- 0.04
HbA1c	- 0.11*	- 0.23**	- 0.04

Table 2: Pearson's correlation† between 25(OH)D levels and glyceimic parameters

In the multinomial logistic regression model, decreased serum 25(OH)D levels were associated with diabetes after adjusting for age, gender, BMI, and seasonality (Table 3). However, this was not observed with pre-diabetes status.

DISCUSSION

In the present study we found that mean serum 25(OH)D levels were significantly lower in individuals with diabetes compared to those with normal glyceimic or pre diabetes. We also found a significant inverse correlation between serum



p<0.05 by ANCOVA, adjusting for age

Figure 1: Difference in 25(OH)D levels by glucose status, according to gender.

25(OH)D (ng/ml)	Odds of pre diabetes (95% CI)			
	Crude	P value	Adjusted*	P value
<12 ng/ml	2.459 (0.671, 9.014)	0.174	2.140 (0.556, 8.231)	0.268
12 – 20 ng/ml	1.365 (0.725, 2.571)	0.335	1.222 (0.616, 2.424)	0.567
21 – 30 ng/ml	1.296 (0.762, 2.203)	0.338	1.392 (0.787, 2.462)	0.256
>30 ng/ml	1.00		1.00	

25(OH)D (ng/ml)	Odds of diabetes (95% CI)			
	Crude	P value	Adjusted*	P value
<12 ng/ml	4.727 (1.566, 14.275)	0.006	4.195 (1.287, 13.666)	0.017
12 – 20 ng/ml	2.373 (1.419, 3.970)	0.001	2.422 (1.348, 4.354)	0.003
21 – 30 ng/ml	1.500 (0.963, 2.335)	0.073	1.773 (1.068, 2.942)	0.027
>30 ng/ml		1.00 (reference)		

*Adjusted for age, gender, BMI and seasonality.

Table 3: Multiple logistic regression of the association between 25(OH)D levels and risk of pre-diabetes and diabetes.

25(OH)D levels and glucose parameters, i.e., FBS and HbA1c, particularly in males. In addition, we found that risk of diabetes increased as levels of serum 25(OH)D levels decreased.

The inverse correlation between serum 25(OH)D levels and HbA1c found in the present study has also been reported in the US using data from the National Health and Nutrition Examination Survey in different racial/ethnic groups and in Caucasians from Italy.^{24,25} Also, studies have found an inverse correlation between serum 25(OH)D levels and FBS, 2h-OGTT, and HbA1c in Caucasians in Germany and Finland ($p < 0.001$).^{23,26} We were unable to detect a significant negative correlation between 25(OH)D and 2 hours OGTT; this could be explained by the fact that insulin resistance was not reported in the medical records to make corrections in this variable, which is known to affect glucose metabolism.

The present study also found an increase diabetes risk as serum 25(OH)D levels decreased, which is consistent with the meta-analysis of Italian, British, Turkish, and Austrians cohort studies reported by Pittas et al.⁸ Similarly, studies in individuals from southern Spain and from Greece have also found a higher incidence of diabetes in individuals with lower vitamin D status.^{27,28} Furthermore, a study in Sweden found that individuals with pre diabetes and diabetes had lower vitamin D status compared to those with normal glycemia,²⁹ similar to results from the present study.

A possible mechanism for the relation between vitamin D status and diabetes is the presence of specific vitamin D receptors in pancreatic beta cells, which improves the function of insulin secretion.³⁰ A recent review suggests that vitamin D deficiency may lead to variations in insulin secretion, glucose intolerance, and diabetes, through a direct action of the vitamin D receptors activation or indirectly through calcemic hormones and also *via* inflammation.³¹ Vitamin D may lower insulin resistance by its effect on phosphorus and calcium metabolism and by its effect on the regulation of the insulin receptor gene.³²

Differences in the recommendations from the IOM and the US Endocrine Society's Practice Guidelines for the classification of vitamin D status reflect different goals and views on current evidence. Using the cut-off points established by the IOM, there are less individuals considered to have inadequate vitamin D status. The IOM levels were established with an endpoint in bone health while the Endocrine Society's cut-off points are more related to optimizing health for preventing endocrine related disorders, such as diabetes. Our results support the views of the Endocrine Society, as our study found that those with serum 25(OH)D levels >30 ng/ml had significantly lower risk of diabetes. Therefore, it is recommended that these cut-off points be used when considering the clinical management of patients with diabetes in this population of Puerto Ricans.

The present study provided the opportunity to describe the association between serum 25(OH)D levels and diabetes using a large sample of Hispanic adults in PR. The use of serum 25(OH)D as a measure of vitamin D status is an important strength because this measures total vitamin D (D2 and D3) status, which reflects vitamin D intake as well as vitamin D synthesized in the skin upon exposure to sun. However, our study has some limitations that merit discussion. First, the temporal relationship between vitamin D and glycemic status cannot be assessed in this cross sectional study. Second, this study was done at a single endocrinology clinic, thus our data are unlikely to be representative of the Puerto Rican population. Third, data on known and suspected risk factors for diabetes were not readily available (such as physical activity, family history of diabetes, central obesity, hypertension, dyslipidemia, and blood pressure), thus their potential confounding effects were not assessed. Longitudinal studies should be performed to confirm our findings in a larger and representative sample of Hispanics.

CONCLUSION

In conclusion, low vitamin D levels significantly increased the odds of diabetes in this sample of Hispanic adults in PR. Given that diabetes is one of the leading causes of death in US and PR and that Puerto Ricans have the highest prevalence of both diabetes and low vitamin D status compared to the US and other territories, it is important that health professionals regularly screen for vitamin D status and offer interventional strategies to correct vitamin D deficiency. These results may have significant public health implications for defining potential intervention and management of diabetes that may be easily implemented in the clinical setting in this high risk population.

NEW CONTRIBUTION TO THE LITERATURE

Low vitamin D status has been significantly associated with diabetes in other populations, mainly Caucasians; however, there is limited information in Hispanics, a group with a high prevalence of both diabetes and low vitamin D status. The present study confirms this association in a group of Hispanic adults in PR and adds to the available evidence for routinely screening this population for vitamin D deficiency.

ACKNOWLEDGMENT

This study was supported in part by Award Number 8G12MD007600 from the National Institute on Minority Health and Health Disparities. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. We are grateful to Dr. Juan J. Maldonado-Rivera for his cooperation in the access to the medical records of his clinic.

REFERENCES

1. Barnes PM, Ward BW, Freeman G, Schiller JS. Early release of selected estimates based on data from the January-March 2011. National Health Interview Survey. National Center for Health Statistics. Available at: <http://www.cdc.gov/nchs/nhis.htm>. Accessed 2013.
2. Daviglius ML, Talavera GA, Aviles-Santa ML, et al. Prevalence of major cardiovascular risk factors and cardiovascular diseases among hispanic/latino individuals of diverse backgrounds in the united states. *JAMA*. 2012; 308(17): 1775-1784. doi: [10.1001/jama.2012.14517](https://doi.org/10.1001/jama.2012.14517)
3. Miranda JJ, Herrera VM, Chirinos JA, et al. Major cardiovascular risk factors in latin america: A comparison with the United States. The Latin American Consortium of Studies in Obesity (LASO). *PLoS One*. 2013; 8(1): e54056. doi: [10.1371/journal.pone.0054056](https://doi.org/10.1371/journal.pone.0054056)
4. Center for Disease Control and Prevention (CDC). Behavioral Risk Factor Surveillance System Survey (BRFSS) data. Available at: <http://www.cdc.gov/BRFSS/>. Accessed 2009.
5. Perez CM, Guzman M, Ortiz AP, et al. Prevalence of the metabolic syndrome in san Juan, Puerto Rico. *Ethn Dis*. 2008; 18(4): 434-441.
6. Center for Disease Control and Prevention (CDC). Deaths: Final data for 2011. Atlanta, GA: 2011; National Vital Statistics Report (NVSr).
7. Song Y, Wang L, Pittas AG, et al. Blood 25-hydroxy vitamin D levels and incident type 2 diabetes: A meta-analysis of prospective studies. *Diabetes Care*. 2013; 36(5): 1422-1428. doi: [10.2337/dc12-0962](https://doi.org/10.2337/dc12-0962)
8. Pittas AG, Lau J, Hu FB, Dawson-Hughes B. The role of vitamin D and calcium in type 2 diabetes. A systematic review and meta-analysis. *J Clin Endocrinol Metab*. 2007; 92(6): 2017-2029. doi: [http://dx.doi.org/10.1210/jc.2007-0298](https://doi.org/10.1210/jc.2007-0298)
9. Palacios C, Gonzalez L. Is vitamin D deficiency a major global public health problem? *J Steroid Biochem Mol Biol*. 2014; 144(Pt A): 138-145. doi: [10.1016/j.jsbmb.2013.11.003](https://doi.org/10.1016/j.jsbmb.2013.11.003)
10. Ganji V, Zhang X, Tangpricha V. Serum 25-hydroxyvitamin D concentrations and prevalence estimates of hypovitaminosis D in the U.S. population based on assay-adjusted data. *J Nutr*. 2012; 142(3): 498-507. doi: [10.3945/jn.111.151977](https://doi.org/10.3945/jn.111.151977)
11. Looker AC, Dawson-Hughes B, Calvo MS, Gunter EW, Sahyoun NR. Serum 25-hydroxyvitamin D status of adolescents and adults in two seasonal subpopulations from NHANES III. *Bone*. 2002; 30(5): 771-777. doi: [http://dx.doi.org/10.1016/S8756-3282\(02\)00692-0](https://doi.org/10.1016/S8756-3282(02)00692-0)
12. Jacobs ET, Alberts DS, Foote JA, et al. Vitamin D insufficiency in southern Arizona. *Am J Clin Nutr*. 2008; 87(3): 608-613.
13. Oliveri B, Plantalech L, Bagur A, et al. High prevalence of vitamin D insufficiency in healthy elderly people living at home in argentina. *Eur J Clin Nutr*. 2004; 58(2): 337-342. doi: [10.1038/sj.ejcn.1601786](https://doi.org/10.1038/sj.ejcn.1601786)
14. Reasner CA 2nd, Dunn JF, Fetchick DA, et al. Alteration of vitamin D metabolism in mexican-americans. *J Bone Miner Res*. 1990; 5(1): 13-17. doi: [10.1002/jbmr.5650050105](https://doi.org/10.1002/jbmr.5650050105)
15. Levis S, Gomez A, Jimenez C, et al. Vitamin d deficiency and seasonal variation in an adult south florida population. *J Clin Endocrinol Metab*. 2005; 90(3): 1557-1562. doi: [http://dx.doi.org/10.1210/jc.2004-0746](https://doi.org/10.1210/jc.2004-0746)
16. Araujo AB, Travison TG, Esche GR, Holick MF, Chen TC, McKinlay JB. Serum 25-hydroxyvitamin D and bone mineral density among Hispanic men. *Osteoporos Int*. 2009; 20(2): 245-255. doi: [10.1007/s00198-008-0652-9](https://doi.org/10.1007/s00198-008-0652-9)
17. Suarez-Martinez EB, Perez CM, Cruz SK, Khorsandi S, Chardon C, Ferder L. Importance of vitamin D and vitamin D levels status in puerto ricans. *J Health Care Poor Underserved*. 2013; 24(4 Suppl): 38-47. doi: [10.1353/hpu.2014.0000](https://doi.org/10.1353/hpu.2014.0000)
18. Holick MF. Vitamin D status: Measurement, interpretation, and clinical application. *Ann Epidemiol*. 2009; 19(2): 73-78. doi: [10.1016/j.annepidem.2007.12.001](https://doi.org/10.1016/j.annepidem.2007.12.001)
19. Institute of Medicine. Dietary Reference Intakes for Calcium and Vitamin D. Washington, DC: The National Academy Press; 2011.
20. Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Evaluation, treatment, and prevention of vitamin d deficiency: An endocrine society clinical practice guideline. *J Clin Endocrinol Metab*. 2011; 96(7): 1911-1930. doi: [10.1210/jc.2011-0385](https://doi.org/10.1210/jc.2011-0385)
21. American Diabetes Association (ADA). Diagnosis and classification of diabetes mellitus. *Diabetes care*. 2013; 36(Suppl 1): S67-S74.
22. Malkani S, Mordes JP. Implications of using hemoglobin A1C for diagnosing diabetes mellitus. *Am J Med*. 2011; 124(5): 395-401. doi: [10.1016/j.amjmed.2010.11.025](https://doi.org/10.1016/j.amjmed.2010.11.025)
23. O'Hartaigh B, Neil Thomas G, Silbernagel G, et al. Association of 25-hydroxyvitamin D with type 2 diabetes among patients undergoing coronary angiography: Cross-sectional findings from the LUDwigshafen risk and cardiovascular health

(LURIC) study. *Clin Endocrinol (Oxf)*. 2013; 79(2): 192-198. doi: [10.1111/cen.12024](https://doi.org/10.1111/cen.12024)

24. Kositsawat J, Freeman VL, Gerber BS, Geraci S. Association of A1C levels with vitamin D status in U.S. adults: Data from the national health and nutrition examination survey. *Diabetes Care*. 2010; 33(6): 1236-1238. doi: [10.2337/dc09-2150](https://doi.org/10.2337/dc09-2150)

25. Zoppini G, Galletti A, Targher G, et al. Glycated haemoglobin is inversely related to serum vitamin D levels in type 2 diabetic patients. *PLoS One*. 2013; 8(12): e82733. doi: [10.1371/journal.pone.0082733](https://doi.org/10.1371/journal.pone.0082733)

26. Hurskainen AR, Virtanen JK, Tuomainen TP, Nurmi T, Vuolteenaho S. Association of serum 25-hydroxyvitamin D with type 2 diabetes and markers of insulin resistance in a general older population in Finland. *Diabetes Metab Res Rev*. 2012; 28(5): 418-423. doi: [10.1002/dmrr.2286](https://doi.org/10.1002/dmrr.2286)

27. Gonzalez-Molero I, Rojo-Martinez G, Morcillo S, et al. Vitamin D and incidence of diabetes: A prospective cohort study. *Clin Nutr*. 2012; 31(4): 571-573. doi: [10.1016/j.clnu.2011.12.001](https://doi.org/10.1016/j.clnu.2011.12.001)

28. Kostoglou-Athanassiou I, Athanassiou P, Gkountouvas A, Kaldrymides P. Vitamin D and glycemic control in diabetes mellitus type 2. *Ther Adv Endocrinol Metab*. 2013; 4(4): 122-128. doi: [10.1177/2042018813501189](https://doi.org/10.1177/2042018813501189)

29. Deleskog A, Hilding A, Brismar K, Hamsten A, Efendic S, Ostenson CG. Low serum 25-hydroxyvitamin D level predicts progression to type 2 diabetes in individuals with prediabetes but not with normal glucose tolerance. *Diabetologia*. 2012; 55(6): 1668-1678. doi: [10.1007/s00125-012-2529-x](https://doi.org/10.1007/s00125-012-2529-x)

30. Pittas AG, Harris SS, Stark PC, Dawson-Hughes B. The effects of calcium and vitamin D supplementation on blood glucose and markers of inflammation in nondiabetic adults. *Diabetes Care*. 2007; 30(4): 980-986. doi: [10.2337/dc06-1994](https://doi.org/10.2337/dc06-1994)

31. Chagas CE, Borges MC, Martini LA, Rogero MM. Focus on vitamin D, inflammation and type 2 diabetes. *Nutrients*. 2012; 4(1): 52-67. doi: [10.3390/nu4010052](https://doi.org/10.3390/nu4010052)

32. Maestro B, Molero S, Bajo S, Davila N, Calle C. Transcriptional activation of the human insulin receptor gene by 1,25-dihydroxyvitamin D(3). *Cell Biochem Funct*. 2002; 20(3): 227-232. doi: [10.1002/cbf.951](https://doi.org/10.1002/cbf.951)