

Opinion

*Corresponding author

Han Huang, MD

Associate Professor
Department of Anesthesiology
West China Second University Hospital
Sichuan University
20#, 3rd Segment, Ren Min Nan Lu
Chengdu 610041, Sichuan, China
Tel. 86-28-8550-3753
Fax: 86-28-8550-3752
E-mail: han.huang@scu.edu.cn

Volume 2 : Issue 2

Article Ref. #: 1000AOJ2115

Article History

Received: December 11th, 2017

Accepted: December 21st, 2017

Published: December 27th, 2017

Citation

Zhou J, Huang H. Mind the gap between the bench and the bed: The general anesthetics-induced neurotoxicity in the real world. *Res Pract Anesthesiol Open J*. 2017; 2(2): 40-41. doi: [10.17140/RPAOJ-2-115](https://doi.org/10.17140/RPAOJ-2-115)

Copyright

©2017 Huang H. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mind the Gap Between the Bench and the Bed: The General Anesthetics-Induced Neurotoxicity in the Real World

Jieshu Zhou, MD¹; Han Huang, MD^{1,2*}

¹Training Anaesthetist, Department of Anesthesiology and Key Laboratory of Birth Defects and Related Diseases of Women and Children (Sichuan University), Ministry of Education, West China Second University Hospital, Sichuan University, Chengdu Shi, Sichuan Sheng, China

²Translational Neuroscience Centre, West China Hospital, Sichuan University, Chengdu Shi, Sichuan Sheng, China

From the very beginning of Morton's successful demonstration of surgical anesthesia, reversibility has been identified as one of the unique characteristics of anesthesia. However, recent studies just reveal the opposite; general anesthetics (GAs) produce long-term, if not permanent, effects in the human brain, especially in the immature brain, at clinically relevant concentrations for clinical relevant durations, so-called the neurotoxicity of GAs on immature brains.¹ The neurotoxicity has been demonstrated with all types of GAs, including volatile anesthetics, benzodiazepine, propofol, and N-methyl-d-aspartate (NMDA) antagonists.² Although, the underlying mechanisms have not been well-illustrated, it has been generally accepted that the neuroinflammation plays a vital role in the pathogenesis of GAs-induced neurotoxicity. Due to huge impact of this subject (animal studies showed that both fetal and early post-natal exposure to GAs caused neurotoxicity), FDA issued a change in labeling regarding the safe use of anesthetic and sedative agents³ and suggest delaying pediatric surgery if possible, to avoid repeated or lengthy exposure to GAs in children under the age of 3 or in pregnant women during their third trimester.

However, most of the adverse results came from animal or *in vitro* studies, while most of the human trial reveal negative results, which indicate that single exposure to GAs at early days in their life caused no neurotoxicity in children.⁴⁻⁶ Even with multiple exposures, the differences are generally very small.⁷

Clearly, there is a huge difference between the animal and human studies designs. In animal studies, subjects were exposed to GAs alone, without any surgical manipulation, in order to observe the isolated effect of GAs on neurodevelopment. However, in the real world, it is unlikely for our young patients to receive GAs alone, without any surgical procedures. On the other hand, most of the surgical procedures cannot be performed without anesthesia. Therefore, young children receiving minor surgery (such as inguinal hernias, circumcisions, cystoscopies, and pyloromyotomies) were included for these clinical trials,⁸ for which it is possible to perform without anesthesia or with light sedation only.

The majority of pediatric surgeries cannot be performed without anesthesia deep enough. So, it is of very limited clinical significance to compare between with and without anesthesia. As we mentioned above, all the currently available GAs have been reported to produce neurotoxicity in developing brains. Therefore, from clinical perspective, it is more reasonable for us to compare different types of GAs, such as inhaled *versus* intravenous, to identify the one with least neurotoxicity, if there is one.

Another factor needs to be considered is that the primary diseases which require surgical treatment. It is evident that different diseases produce different effects on neurodevelopment.⁹ So, it is equally important to restrict the types of diseases in the future trial.

In summary, the causal relationship between early-life exposure to GAs and neurodevelopment impairment has not been proved with human evidence, which remains one of the most intensively investigated filed in our specialty. But considering the convincing results from massive preclinical studies, it may be the time for us, the anesthesiologists, to accept the concept that anesthesia is not completely reverse and it is highly like to produce a permanent effect in developing the brain. However, we should not dwell on whether GA is neurotoxic or not. No matter how toxic it could be, most of the surgeries cannot be performed without proper anesthesia, because the untreated pain caused much more harm to children and their developing brains. Instead, it is urgent for us to find out the least toxic GA among current available GAs. Like the surgeons, who never stop performing surgery in the fear of an ugly scar, instead, they are just trying their best to make it smaller. We, anesthesiologists, should not be demanding of the neurotoxicity of GAs on developing brain while ignoring the huge benefit provided by our excellent care.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

REFERENCES

1. Vutskits L, Xie Z. Lasting impact of general anaesthesia on the brain: Mechanisms and relevance. *Nat Rev Neurosci*. 2016; 17: 705-717. doi: [10.1038/nrn.2016.128](https://doi.org/10.1038/nrn.2016.128)
2. Lin EP, Lee JR, Lee CS, Deng M, Loepke AW. Do anesthetics harm the developing human brain? An integrative analysis of animal and human studies. *Neurotoxicol Teratol*. 2017; 60: 117-128. doi: [10.1016/j.ntt.2016.10.008](https://doi.org/10.1016/j.ntt.2016.10.008)
3. FDA Drug Safety Communication: FDA review results in new warnings about using general anesthetics and sedation drugs in young children and pregnant women. Web site. <http://www.fda.gov/Drugs/DrugSafety/ucm532356.htm>. Accessed December 10, 2017.
4. Glatz P, Sandin RH, Pedersen NL, Bonamy AK, Eriksson LI, Granath F. Association of anesthesia and surgery during childhood with long-term academic performance. *JAMA Pediatr*. 2017; 171: e163470. doi: [10.1001/jamapediatrics.2016.3470](https://doi.org/10.1001/jamapediatrics.2016.3470)
5. Sun LS, Li G, Miller TL, et al. Association between a single general anesthesia exposure before age 36 months and neurocognitive outcomes in later childhood. *JAMA*. 2016; 315: 2312-2320. doi: [10.1001/jama.2016.6967](https://doi.org/10.1001/jama.2016.6967)
6. Davidson AJ, Disma N, de Graaff JC, et al. Neurodevelopmental outcome at 2 years of age after general anaesthesia and awake-regional anaesthesia in infancy (GAS): An international multicentre, randomised controlled trial. *Lancet*. 2016; 387: 239-250. doi: [10.1016/S0140-6736\(15\)00608-X](https://doi.org/10.1016/S0140-6736(15)00608-X)
7. Graham MR, Brownell M, Chateau DG, Dragan RD, Burchill C, Fransoo RR. Neurodevelopmental assessment in kindergarten in children exposed to general anesthesia before the age of 4 years: A retrospective matched cohort study. *Anesthesiology*. 2016; 125: 667-677. doi: [10.1097/ALN.0000000000001245](https://doi.org/10.1097/ALN.0000000000001245)
8. Bong CL, Allen JC, Kim JT. The effects of exposure to general anesthesia in infancy on academic performance at age 12. *Anesth Analg*. 2013; 117: 1419-1428. doi: [10.1213/ANE.0b013e318299a7c2](https://doi.org/10.1213/ANE.0b013e318299a7c2)
9. Armstrong FD. Neurodevelopment and chronic illness: Mechanisms of disease and treatment. *Ment Retard Dev Disabil Res Rev*. 2006; 12(3): 168-173. doi: [10.1002/mrdd.20114](https://doi.org/10.1002/mrdd.20114)