

Short Communication

*Corresponding author

Manxue Jia, MD, PhD

Research Scientist

Aaron Diamond AIDS Research Center

Affiliate of the Rockefeller University

New York 10016, USA

E-mail: mjia@adarc.org

Volume 3 : Issue 1

Article Ref. #: 1000HARTO3120

Article History

Received: June 6th, 2016

Accepted: June 13th, 2016

Published: June 13th, 2016

Citation

Jia M. Natural killer (NK) cells and human immunodeficiency virus (HIV) infection. *HIV/AIDS Res Treat Open J.* 2016; 3(1): 13-14. doi: [10.17140/HARTOJ-3-120](https://doi.org/10.17140/HARTOJ-3-120)

Copyright

©2016 Jia M. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Natural Killer (NK) Cells and Human Immunodeficiency Virus (HIV) Infection

Manxue Jia, MD, PhD*

Aaron Diamond AIDS Research Center, Affiliate of the Rockefeller University, New York 10016, USA

ABSTRACT

Human Immunodeficiency Virus (HIV) infection has become a serious problem of public health. Recently, it is observed cellular immunity is involved in HIV infection and disease progression, especially the Natural Killer (NK) cells, which are important part of innate immunity. In order to clearly understand the mechanism of HIV infection, we try to figure out the function of NK cells during HIV infection. The article reviews the function of NK cells and the effect of NK cells in HIV infection.

KEYWORDS: Natural Killer (NK); Human Immunodeficiency Virus (HIV) infection; Virus.

The human immunodeficiency virus (HIV) directly targets and devastates the host's immune system, leading to serious infection. Recently, research found innate immunity played very important role in the host's response against viruses, and increasing data indicate that innate immune responses play a key role in the development of effective vaccine-induced immune responses. In particular, Natural Killer (NK) cells represent important early effector cells of the antiviral innate immune defense, which was discovered from 1975 by Herberman.¹ NK cells, which account for up to 15% of Peripheral Blood Lymphocytes (PBL), are classified into three subsets, CD3^{neg}CD16^{pos}CD56^{dim}, which is composed of 80%~90% NK cells; CD3^{neg}CD56^{bright}, which is composed of 2~10% NK cells; and CD3^{neg}CD16^{pos}CD56^{neg}, which is composed of 5~10% NK cells.² NK cells can lyse virally infected cells without prior sensitization and participate in the regulation of innate and adaptive immune responses. The ability of NK cells mediating protection against viral infection depends on the relative abundance of each subset. Data from our laboratory and others have demonstrated that the clinical progression of HIV is correlated with changes in the distribution of NK cell sub-populations, namely a decrease in CD16^{pos}CD56^{dim} subsets and an increase in CD16^{pos}CD56^{neg} subsets.

The effects of NK subsets rely on the composition of receptors on the NK cell surface, including inhibitory NK receptors (iNKR) and activating receptors.³ These receptors recognize relative ligands of target cells, and induce activating or inhibitory signal. During infections, the binding of NK cell receptors to ligands on the target cell can cause it to overcome inhibitory signals and become activated, culminating in the killing of the target cell. NK cells can be activated by the binding of its receptors (such as CD16) to the Fc portion of antibodies attached to the surface of target cells. We also found increased expression of the activating receptor NKG2C, NKG2C and inhibitory receptor NKG2A in patients with deteriorating clinical status. The ratio of NKG2A to NKG2C receptors decreased with advanced clinical progression of disease. Therefore, increased activation of NK cells (through modulated expression of NKG2A compared to NKG2C) may be a marker for identifying the clinical course of HIV-1 infection. Epidemiological studies have shown that particular NK Killer cell immunoglobulin-like receptor (KIR) genes expressed in conjunction with their HLA ligands are associated with significantly slower HIV-1 disease progression and lower viral set-point.^{4,5} NK cells expressing protective KIR genotypes are associated with protection from infection, can significantly inhibit HIV-1 replication *in vitro*.⁶⁻⁹ We observed the inhibitory killer immunoglobulin-like receptor CD158a (KIR2DL1) to be up-regulated in HIV-positive individuals.⁶

More recently, scientists showed that NK cells are involved in several other pathways to combat HIV infection, such as through secreting degranulated perforin and granzyme B to kill target cells; the Fas-FasL pathway to induce lysis of infected cells; production of cytokines to regulate immunity; and antibody-dependent cell-mediated cytotoxicity (ADCC) to lysis infected cells.⁶ NK cells could kill target cells and inhibit HIV-1 replication, at least partly due to antibody-dependent cell-mediated cytotoxicity (ADCC). The mechanism of ADCC against HIV-infected cells is: Fc receptor-positive effector cells bind to gp120 or gp41-expressing HIV-infected target cells *via* gp120 or gp41 specific antibodies of certain Immunoglobulin G (IgG) isotypes and mediate their killing. NK cell is the important effector cell inducing ADCC. However, the roles of NK cells against infection are still under debate, and mechanisms are needed intensive research. Consistent with some other studies, our study also showed that ADCC responses were associated with slower progression of HIV infection. In addition, NK cells are protective against HIV disease progression through combinations of killer cell immunoglobulin-like receptors (KIR) and specific human leucocyte antigen class I (HLA-1) ligands; the studies have shown that particular HLA alleles appear to be associated with stemming the progression of HIV-1 infection.¹⁰ We also demonstrated that ADCC responses are associated negatively with the clinical progression of HIV-1 infection and correlated positively with infected time and specific HLA alleles (HLA-A*30/B*13/Cw*06) in long-term survivors in a uniformly infected cohort.⁶ Taken together, these studies suggest that NK cells might play an important role in controlling HIV-1 replication, and that the virus can evade NK cell mediated immune pressure by selecting for sequence polymorphisms in areas targeted by KIR⁺ NK cells.

This review briefly summarized the function of NK cells immunity against HIV infection and disease progression, which may provide helpful data into understanding of HIV-1 pathogenesis and immune mechanisms, facilitating anti-virus drugs development and vaccine evaluation.

REFERENCES

1. Herberman RB, Nunn ME, Lavrin DH. Natural cytotoxic reactivity of mouse lymphoid cells against syngeneic acid allogeneic tumors. I. Distribution of reactivity and specificity. *Int J Cancer*. 1975; 16(2): 216-229.
2. Mavilio D, Lombardo G, Benjamin J, et al. Characterization of CD56-/CD16+ natural killer (NK) cells: A highly dysfunctiona lNK subset expanded in HIV-infected viremic individuals. *Proc Natl Acad Sci USA*. 2005; 102(8): 2886-2891. doi: [10.1073/pnas.0409872102](https://doi.org/10.1073/pnas.0409872102)
3. Jia M, Li D, Shao Y. Advance research on relationship between NK cells and human immunodeficiency virus infection. *Chinese Tropical Medicine*. 2012; 12(4): 537-541.
4. Vivian JP, Duncan RC, Berry R, et al. Killer cell immunoglobulin-like receptor 3DL1-mediated recognition of human leukocyte antigen B. *Nature*. 2011; 479(7373): 401-405. doi: [10.1038/nature10517](https://doi.org/10.1038/nature10517)
5. Fadda L, O'Connor GM, Kumar S, et al. Common HIV-1 peptide variants mediate differential binding of KIR3DL1 to HLA-Bw4 molecules. *J Virol*. 2011; 85(12): 5970-5974. doi: [10.1128/JVI.00412-11](https://doi.org/10.1128/JVI.00412-11)
6. Jia M, Li D, He X, et al. Impaired natural killer cell-induced antibody-dependent cell-mediated cytotoxicity is associated with HIV-1 disease progression. *Clin Exp Immunol*. 2013; 171(1): 107-116. doi: [10.1111/j.1365-2249.2012.04672.x](https://doi.org/10.1111/j.1365-2249.2012.04672.x)
7. Brackenridge S, Evans EJ, Toebes M, et al. An early HIV mutation within an HLA-B*57-restricted T cell epitope abrogates binding to the killer inhibitory receptor 3DL1. *J Virol*. 2011; 85(11): 5415-5422. doi: [10.1128/JVI.00238-11](https://doi.org/10.1128/JVI.00238-11)
8. Thananchai H, Makadzange T, Maenaka K, et al. Reciprocal recognition of an HLA-Cw4-restricted HIV-1 gp120 epitope by CD8+ T cells and NK cells. *AIDS*. 2009; 23(2): 189-193. doi: [10.1097/QAD.0b013e32831fb55a](https://doi.org/10.1097/QAD.0b013e32831fb55a)
9. Yu XG, Addo MM, Perkins BA, et al. Differences in the expressed HLA class I alleles effect the differential clustering of HIV type 1-specific T cell responses in infected Chinese and caucasians. *AIDS Res Hum Retroviruses*. 2004; 20(5): 557-564. doi: [10.1089/088922204323087813](https://doi.org/10.1089/088922204323087813)
10. Boulet S, Kleyman M, Kim JY, et al. A combined genotype of KIR3DL1 high expressing alleles and HLA-B*57 is associated with a reduced risk of HIV infection. *AIDS*. 2008; 22(12): 1487-1491. doi: [10.1097/QAD.0b013e3282ffde7e](https://doi.org/10.1097/QAD.0b013e3282ffde7e)