

Illustration

Necrotizing Fasciitis: Fatal Evolution of Acral Melanoma

El Marfi Abdelhafid, MD^{1*}; Samia Mrabat, MD²; Kaoutar Laamari, MD²; Mohammed El Idrissi, PhD¹; Abdelhalim El Ibrahimi, PhD¹; Abdelmajid El Mrini, PhD¹; Zakia Douhi, MD²; Sara Elloudi, PhD²; Hanane Baybay, PhD²; Mernissi F. Zahra, PhD²

¹Department of Traumatology and Orthopedic Surgery, University Hospital Hassan II, Fes, Morocco

²Department of Dermatology, University Hospital Hassan II, Fes, Morocco

*Corresponding author

Abdelhafid El Marfi, MD

Department of Traumatology and Orthopedic Surgery, University Hospital Hassan II, Fes, Morocco; Tel. +212 678 33 98 79; Fax. 05 35 61 37 29;

E-mail: hafid.elmarfi@gmail.com

Article information

Received: November 7th, 2019; **Revised:** December 21st, 2019; **Accepted:** January 18th, 2020; **Published:** January 28th, 2020

Cite this article

El Marfi A, Mrabat S, Laamari K, et al. Necrotizing fasciitis: Fatal evolution of acral melanoma. *Orthop Res Traumatol Open J.* 2020; 5(1): 1-2.

doi: [10.17140/ORTOJ-5-116](https://doi.org/10.17140/ORTOJ-5-116)

Necrotizing fasciitis (NF) is a life-threatening soft tissue infection characterized by a rapidly spreading infection of the subcutaneous tissue and in particular the fascia. Early diagnosis and radical surgical debridement are required. Surgery is indicated if NF is clinically suspected and must not be delayed by diagnosis in order to reduce mortality. We report a case of NF following surgery of an acral melanoma.^{1,2}

A 73-year-old patient, with a history of type 2 diabetes with a poor glycemic balance, was diagnosed with acral non metastatic pigmented melanoma of the fifth toe (Figure 1) and underwent surgery to excise the tumor. He was seen 10-days later with a painful and acute red leg along with signs of sepsis (fever, hypotension, tachycardia). Dermatological examination found an ill-limited oedematous erythematous leg with large phlyctenes and necrotic areas extending from the back of the foot to the left leg (Figure 2). Palpation of the leg found crepitus and hypoesthetic zones. The member's X-ray showed an infiltration of the soft parts of the leg

with the presence of air bubbles (Figure 3). As the diagnosis of necrotizing fasciitis was highly suspected, the patient underwent surgery (Figure 4) for an discharge aponevrotomy. He died hours later due to a septic shock.



Figure 4. An Intraoperative Picture of the Discharge Aponevrotomy



CONSENT

The authors have received written informed consent from the patient.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

REFERENCES

1. Leiblein M, Marzi I, Sander AL, Barker JH, Ebert F, Frank J. Necrotizing fasciitis: Treatment concepts and clinical results. *Eur J Trauma Emerg Surg.* 2018; 44(2): 279-290. doi: [10.1007/s00068-017-0792-8](https://doi.org/10.1007/s00068-017-0792-8)
2. Hakkarainen TW, Kopari NM, Pham TN, Evans HL. Necrotizing soft tissue infections: Review and current concepts in treatment, systems of care, and outcomes. *Curr Probl Surg.* 2014; 51(8): 344–362. doi: [10.1067/j.cpsurg.2014.06.001](https://doi.org/10.1067/j.cpsurg.2014.06.001)