

Editorial

Corresponding author

Sarah Dee Geiger, PhD, MS

Visiting Assistant Professor
College of Health and Human
Sciences

Northern Illinois University
255 Wirtz Hall, DeKalb
IL 60115, USA

E-mail: Geiger@niu.edu

Volume 1 : Issue 1

Article Ref. #: 1000PNNOJ1e001

Article History

Received: May 1st, 2014

Accepted: May 2nd, 2014

Published: May 2nd, 2014

Citation

Geiger SD. Pediatric exposures to persistent environmental chemicals. *Pediatr Neonatal Nurs Open J*. 2014; 1(1): e1-e2. doi: [10.17140/PNNOJ-1-e001](https://doi.org/10.17140/PNNOJ-1-e001)

Copyright

©2014 Geiger SD. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Pediatric Exposures to Persistent Environmental Chemicals

Sarah Dee Geiger*

College of Health and Human Sciences, Northern Illinois University, 255 Wirtz Hall, DeKalb, IL 60115, USA

Traditional risk factors are responsible for about 70% of the population attributable risk of Cardiovascular Disease (CVD).¹ Common environmental exposures are known to be responsible for some portion of the remaining 30%. Therefore, it is important to study chemicals like the class known as Perfluoroalkyl Chemicals (PFCs) or Perfluoroalkyl Substances (PSAs). Two of the most highly studied PFCs are Perfluorooctanoic Acid (PFOA or C8) and Perfluorooctane Sulfonate (PFOS or C8S). PFCs are persistent in the environment and associations have been shown with a whole host of negative health outcomes in laboratory animals, including endocrine-disrupting properties as well as developmental effects.² To make matters worse, PFCs have been detected in the blood of >98% of the US population³ and epidemiological studies have demonstrated associations between PFOA and PFOS, and many negative health outcomes such as cancer,⁴ CVD,⁵ osteoarthritis,⁶ hyperuricemia,⁷ pregnancy-induced hypertension,⁸ endocrine disruption,⁹ dyslipidemia,^{10,11} and reproductive effects,^{12,13} often times even at baseline levels typical of the general population's exposure level.

Children are an ideal sample for studies examining the relationship between common environmental exposures and health outcomes because as a group they are largely devoid of cumulative lifestyle risk factors typically experienced by adults; this phenomenon results in associations less subject to confounding. Perhaps more importantly, pediatrics populations are one of our most vulnerable and should be studied with particular rigor in terms of health effects resulting from virtually unavoidable environmental exposures.

In this field of research, the trajectory tends to move from animal studies to occupational epidemiological studies to highly exposed community-based, to population-based, and finally to pediatrics, before possibly moving on to longitudinal study design. Although associations between PFCs and health outcomes among children have not yet been extensively studied, there is a small and growing body of existing literature in this area.

Intermediate cardiovascular disease outcomes among children are known risk factors for earlier onset of more severe forms of CVD, as well as other types of intermediate CVD that tend to cluster together.¹⁴ In this context, Frisbee et al., using C8 Health Project data (n=12,476) noted a significant positive association among highly exposed children between PFCs and total cholesterol and Low-Density Lipoprotein Cholesterol (LDL-C).¹⁵ Another study by Geiger et al. confirmed the associations among children using nationally representative US data, showing an overall positive, significant association between both PFOA and PFOS, and total cholesterol and LDL-C.¹⁶ Geiger et al. also examined associations between PFCs and serum uric acid levels and results indicated a strong, significant relationship independent even after complex multivariate adjustment.¹⁷ Finally, a cross-sectional study by Hoffman et al.¹⁸ found a modest association between PFOA (OR 1.12; 95% confidence interval [CI]: 1.01-1.23) and PFOS (OR 1.03; 95% CI: 1.01-1.05) and attention deficit hyperactivity disorder among children (n=571).

Needless to say, more research, particularly of longitudinal study design, is needed on child exposure to this class of chemicals as well as other persistent organic pollutants such as Polychlorinated Biphenyls (PCBs) as well as semi-persistent organic pollutants such as Bisphenol-A (BPA).

REFERENCES

1. Willett WC. Balancing life-style and genomics research for disease prevention. *Science*. 2002; 26: 296(5568): 695-698. doi: [10.1126/science.1071055](https://doi.org/10.1126/science.1071055)
2. American Heart Association. Heart Disease and Stroke Statistics: 2006 Update Dallas, TX. Website: <https://circ.ahajournals.org/content/113/6/e85.extract>. 2006; Accessed May 27, 2009.
3. Calafat AM, Wong LY, Kuklennyik Z, Reidy JA, Needham LL. Polyfluoroalkyl chemicals in the U.S. population: data from the National Health and Nutrition Examination Survey (NHANES) 2003-2004 and comparisons with NHANES 1999-2000. *Environ Health Perspect*. 2007; 115(11): 1596-1602. doi: [10.1289/ehp.10598](https://doi.org/10.1289/ehp.10598)
4. Eriksen KT, Sorensen M, McLaughlin JK et al. Perfluorooctanoate and perfluorooctanesulfonate plasma levels and risk of cancer in the general Danish population. *J Natl Cancer Inst*. 2009; 15: 101(8): 605-609. doi: [10.1093/jnci/djp041](https://doi.org/10.1093/jnci/djp041)
5. Sakr CJ, Symons JM, Kreckmann KH, Leonard RC. Ischaemic heart disease mortality study among workers with occupational exposure to ammonium perfluorooctanoate. *Occup Environ Med*. 2009; 66(10): 699-703. doi: [10.1136/oem.2008.041582](https://doi.org/10.1136/oem.2008.041582)
6. Innes KE, Ducatman AM, Luster MI, Shankar A. Association of osteoarthritis with serum levels of the environmental contaminants perfluorooctanoate and perfluorooctane sulfonate in a large Appalachian population. *Am J Epidemiol*. 2011; 15: 174(4): 440-450. doi: [10.1093/aje/kwr107](https://doi.org/10.1093/aje/kwr107)
7. Shankar A, Xiao J, Ducatman A. Perfluoroalkyl chemicals and elevated serum uric acid in US adults. *Clin Epidemiol*. 2011; 3: 251-258. doi: [10.2147/CLEP.S21677](https://doi.org/10.2147/CLEP.S21677)
8. C8 Science Panel. Probable Link Evaluation of Pregnancy Induced Hypertension and Preeclampsia. Website: http://www.c8sciencepanel.org/pdfs/Probable_Link_C8_PIH_5Dec2011.pdf. 2012; Accessed February 1, 2012.
9. Olsen GW, Zobel LR. Assessment of lipid, hepatic, and thyroid parameters with serum perfluorooctanoate (PFOA) concentrations in fluorochemical production workers. *Int Arch Occup Environ Health*. 2007; 81(2): 231-246. doi: [10.1007/s00420-007-0213-0](https://doi.org/10.1007/s00420-007-0213-0)
10. Sakr CJ, Kreckmann KH, Green JW, Gillies PJ, Reynolds JL, Leonard RC. Cross-sectional study of lipids and liver enzymes related to a serum biomarker of exposure (ammonium perfluorooctanoate or APFO) as part of a general health survey in a cohort of occupationally exposed workers. *J Occup Environ Med*. 2007; 49(10): 1086-1096.
11. Frisbee SJ, Shankar A, Knox SS et al. Perfluorooctanoic acid, perfluorooctanesulfonate, and serum lipids in children and adolescents: results from the C8 Health Project. *Arch Pediatr Adolesc Med*. 2010; 164(9): 860-869. doi: [10.1001/archpediatrics.2010.163](https://doi.org/10.1001/archpediatrics.2010.163)
12. Sakr CJ, Kreckmann KH, Green JW, Gillies PJ, Reynolds JL, Leonard RC. Cross-sectional study of lipids and liver enzymes related to a serum biomarker of exposure (ammonium perfluorooctanoate or APFO) as part of a general health survey in a cohort of occupationally exposed workers. *J Occup Environ Med*. 2007; 49(10): 1086-1096.
13. Joensen UN, Bossi R, Leffers H, et al. Do perfluoroalkyl compounds impair human semen quality? *Environ Health Perspect*. 2009; 117(6): 923-927. doi: [10.1289/ehp.0800517](https://doi.org/10.1289/ehp.0800517)
14. Daniels SR, Greer FR. Lipid screening and cardiovascular health in childhood. *Pediatrics*. 2008; 122(1): 198-208. doi: [10.1542/peds.2008-1349](https://doi.org/10.1542/peds.2008-1349)
15. Frisbee SJ, Shankar A, Knox SS, et al. Perfluorooctanoic acid, perfluorooctanesulfonate, and serum lipids in children and adolescents: results from the C8 Health Project. *Arch Pediatr Adolesc Med*. 2010; 164(9): 860-869.
16. Geiger SD, Xiao J, Ducatman A, et al. The association between perfluoroalkyl chemicals and serum lipid levels in adolescents. *Chemosphere*. 2014; 98: 78-83.
17. Geiger SD, Xiao J, Shankar A. Positive association between perfluoroalkyl chemicals and hyperuricemia in children. *American Journal of Epidemiology*. 2013; 177(11): 1255-1262. doi: [10.1093/aje/kws392](https://doi.org/10.1093/aje/kws392)
18. Hoffman K, Webster TF, Weisskopf MG, Weinberg J, Vieira VM. Exposure to polyfluoroalkyl chemicals and attention deficit/hyperactivity disorder in U.S. children 12- 15 years of age. *Environ Health Perspect*. 2010; 118(12): 1762-1767. doi: [10.1289/ehp.1001898](https://doi.org/10.1289/ehp.1001898)